

Table 2.11C Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	874	926	920	900	852	846
12	20	19	12	15	9	9
13	33	30	26	23	19	17
14	35	34	30	28	22	19
15	47	48	38	40	29	32
16	57	52	50	45	35	36
17	63	59	54	52	44	43
18	98	104	87	91	70	75
19	95	101	85	90	71	76
20	106	121	88	94	79	74
21	102	100	85	82	71	70
22	94	109	79	91	70	83
23	91	95	74	79	68	70
24	95	102	74	81	66	71
25	94	105	78	80	72	72
26-29	370	393	287	315	264	286
30-34	421	426	299	300	271	263
35-39	397	416	284	281	259	254
40-44	444	429	309	274	276	242
45-49	476	469	299	311	280	293
50-54	626	610	436	384	408	366
55-59	626	525	380	305	357	284
60-64	541	540	288	269	272	253
65 or Older	466	445	318	338	304	320

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.11D Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.34	0.36	0.36	0.35	0.34	0.33
12	0.51	0.50	0.32	0.39	0.24	0.25
13	0.75	0.66	0.60	0.53	0.44	0.39
14	0.83	0.76	0.73	0.62	0.55	0.45
15	0.93	0.96	0.80	0.85	0.68	0.70
16	1.04	0.99	0.97	0.90	0.76	0.75
17	1.08	1.11	1.03	1.04	0.89	0.90
18	1.19	1.18	1.19	1.22	1.09	1.17
19	1.22	1.25	1.25	1.28	1.19	1.30
20	1.24	1.31	1.29	1.31	1.32	1.27
21	1.37	1.27	1.33	1.31	1.29	1.25
22	1.13	1.20	1.18	1.33	1.17	1.30
23	1.15	1.23	1.27	1.37	1.25	1.32
24	1.20	1.17	1.34	1.36	1.31	1.31
25	1.20	1.13	1.39	1.28	1.40	1.29
26-29	1.06	1.14	1.26	1.24	1.23	1.19
30-34	1.00	1.06	1.15	1.16	1.10	1.07
35-39	1.10	1.10	1.14	1.19	1.12	1.12
40-44	1.00	1.16	1.14	1.17	1.08	1.08
45-49	0.93	1.05	1.05	1.14	1.02	1.11
50-54	1.37	1.30	1.56	1.40	1.50	1.36
55-59	1.43	1.45	1.61	1.41	1.55	1.33
60-64	1.71	1.62	1.62	1.35	1.55	1.27
65 or Older	1.20	1.11	0.82	0.84	0.78	0.80

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.11P Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2011 Versus 2010

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0080		0.0213		0.0299	
12	0.4840		0.4054		0.9086	
13	0.1021		0.0909		0.1605	
14	0.0770		0.0092		0.0888	
15	0.0045		0.0450		0.0436	
16	0.3120		0.3585		0.9560	
17	0.1238		0.5009		0.8014	
18	0.9017		0.8952		0.9116	
19	0.0096		0.0489		0.0365	
20	0.1059		0.2120		0.1561	
21	0.3409		0.4209		0.3974	
22	0.3136		0.0964		0.3994	
23	0.8284		0.4602		0.5048	
24	0.4737		0.7522		0.2892	
25	0.3433		0.7355		0.3932	
26-29	0.0138		0.3228		0.1581	
30-34	0.1914		0.5967		0.6127	
35-39	0.6111		0.8741		0.8864	
40-44	0.5397		0.1127		0.1030	
45-49	0.0329		0.7510		0.6778	
50-54	0.8673		0.8968		0.3764	
55-59	0.5065		0.0965		0.0810	
60-64	0.9691		0.6593		0.5317	
65 or Older	0.7177		0.5742		0.7404	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.12C Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	882	927	838	859	780	816
12	17	17	11	13	7	9
13	29	24	23	20	17	13
14	31	31	26	25	20	17
15	42	44	33	35	25	27
16	51	49	43	41	32	33
17	60	54	50	46	39	38
18	90	91	77	79	64	67
19	88	93	76	80	63	67
20	96	106	78	84	68	69
21	93	92	75	73	66	65
22	89	103	71	85	64	75
23	88	90	66	74	60	66
24	90	93	70	73	62	65
25	88	100	69	72	65	64
26-29	355	374	262	287	244	259
30-34	402	414	269	269	252	237
35-39	377	402	256	251	227	229
40-44	421	414	275	229	262	217
45-49	456	448	262	286	250	271
50-54	617	593	409	361	391	351
55-59	612	510	342	262	323	244
60-64	524	526	257	242	243	226
65 or Older	470	458	277	307	266	293

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.12D Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.35	0.36	0.33	0.33	0.31	0.32
12	0.43	0.44	0.27	0.34	0.19	0.24
13	0.66	0.55	0.55	0.45	0.41	0.32
14	0.75	0.70	0.65	0.57	0.50	0.40
15	0.89	0.91	0.73	0.76	0.60	0.61
16	1.02	0.96	0.89	0.84	0.70	0.69
17	1.09	1.06	0.98	0.96	0.81	0.82
18	1.25	1.23	1.18	1.23	1.05	1.13
19	1.28	1.30	1.25	1.30	1.17	1.24
20	1.22	1.28	1.24	1.31	1.24	1.23
21	1.39	1.30	1.33	1.28	1.25	1.23
22	1.14	1.22	1.21	1.30	1.19	1.25
23	1.26	1.28	1.28	1.32	1.24	1.28
24	1.28	1.28	1.34	1.30	1.30	1.25
25	1.27	1.21	1.40	1.33	1.35	1.30
26-29	1.05	1.21	1.21	1.22	1.16	1.17
30-34	1.02	1.12	1.10	1.11	1.05	1.00
35-39	1.09	1.21	1.10	1.13	1.04	1.07
40-44	1.07	1.16	1.10	1.06	1.07	1.02
45-49	0.99	1.07	0.99	1.11	0.96	1.06
50-54	1.42	1.34	1.53	1.37	1.47	1.34
55-59	1.53	1.49	1.51	1.28	1.45	1.20
60-64	1.76	1.67	1.49	1.23	1.41	1.16
65 or Older	1.21	1.14	0.71	0.76	0.68	0.73

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.12P Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2011 Versus 2010

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.0028		0.0321		0.0308	
12	0.6140		0.4632		0.7915	
13	0.0405		0.0178		0.0746	
14	0.0974		0.0443		0.1211	
15	0.0195		0.0157		0.0396	
16	0.7647		0.8915		0.6679	
17	0.2051		0.5135		0.7056	
18	0.5353		0.6456		0.6899	
19	0.0189		0.2290		0.1013	
20	0.3066		0.6387		0.8357	
21	0.8646		0.3966		0.4037	
22	0.2789		0.5186		0.2291	
23	0.7227		0.3194		0.1678	
24	0.1383		0.1981		0.1704	
25	0.5158		0.8915		0.3257	
26-29	0.0051		0.3219		0.0562	
30-34	0.3348		0.7201		0.3950	
35-39	0.7208		0.9140		0.8100	
40-44	0.8868		0.0533		0.0852	
45-49	0.0382		0.8558		0.9226	
50-54	0.8156		0.9837		0.5264	
55-59	0.3362		0.0439		0.0348	
60-64	0.9096		0.6277		0.5147	
65 or Older	0.6334		0.5123		0.6150	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.13C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	700	752	351	326	314	278
12	9	9	6	7	4	3
13	17	15	14	11	10	9
14	18	17	16	12	10	7
15	25	26	21	23	13	14
16	29	28	23	23	16	15
17	31	31	25	27	17	17
18	48	55	42	43	34	30
19	51	45	37	33	27	19
20	55	46	38	32	33	25
21	52	49	32	32	22	24
22	50	52	34	34	26	29
23	47	43	28	27	24	21
24	45	47	27	29	21	20
25	55	51	32	31	26	26
26-29	185	194	104	97	92	72
30-34	196	224	106	96	88	84
35-39	238	226	132	110	127	103
40-44	228	241	91	95	85	89
45-49	233	238	105	81	96	76
50-54	294	275	95	102	79	78
55-59	245	240	96	92	93	80
60-64	200	217	87	89	69	81
65 or Older	316	326	129	123	120	117

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.13D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.28	0.29	0.14	0.13	0.12	0.11
12	0.24	0.24	0.16	0.20	0.11	0.08
13	0.40	0.36	0.33	0.27	0.24	0.22
14	0.45	0.39	0.41	0.27	0.25	0.17
15	0.56	0.57	0.47	0.53	0.30	0.32
16	0.67	0.60	0.53	0.50	0.37	0.35
17	0.70	0.70	0.58	0.60	0.41	0.40
18	0.92	1.00	0.83	0.83	0.70	0.59
19	1.01	0.95	0.77	0.73	0.58	0.44
20	1.03	0.98	0.76	0.72	0.69	0.56
21	1.04	0.98	0.68	0.68	0.48	0.53
22	1.01	0.98	0.75	0.69	0.59	0.61
23	1.02	0.96	0.66	0.66	0.56	0.52
24	1.03	1.01	0.66	0.68	0.51	0.48
25	1.15	1.08	0.75	0.71	0.62	0.62
26-29	0.96	0.99	0.61	0.55	0.55	0.42
30-34	0.86	0.99	0.52	0.47	0.44	0.42
35-39	0.99	1.00	0.64	0.54	0.62	0.51
40-44	0.96	1.01	0.44	0.45	0.41	0.42
45-49	0.90	0.97	0.46	0.37	0.42	0.35
50-54	1.19	1.09	0.43	0.44	0.36	0.34
55-59	1.19	1.19	0.49	0.48	0.48	0.42
60-64	1.16	1.09	0.52	0.49	0.42	0.44
65 or Older	0.81	0.81	0.33	0.31	0.31	0.29

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.13P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2011 Versus 2010

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.5970		0.0750		0.0585	
12	0.9823		0.5941		0.9782	
13	0.3205		0.1500		0.3629	
14	0.2359		0.0036		0.0734	
15	0.2704		0.4187		0.7254	
16	0.4597		0.9194		0.8094	
17	0.5557		0.6583		0.4099	
18	0.2977		0.3351		0.2319	
19	0.0243		0.0114		0.0007	
20	0.2838		0.4354		0.0295	
21	0.5012		0.3976		0.9137	
22	0.6228		0.7192		0.5075	
23	0.0366		0.1965		0.1126	
24	0.7038		0.2894		0.1495	
25	0.9953		0.5471		0.6662	
26-29	0.1700		0.5285		0.1117	
30-34	0.2762		0.6800		0.9651	
35-39	0.0097		0.1697		0.1638	
40-44	0.8116		0.4075		0.3348	
45-49	0.2680		0.3376		0.7603	
50-54	0.6463		0.4684		0.9264	
55-59	0.1761		0.8812		0.5256	
60-64	0.5548		0.7662		0.4487	
65 or Older	0.1469		0.9564		0.9175	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.14C Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	929	920	510	504	363	373
12	8	12	5	10	4	5
13	17	18	13	14	9	10
14	23	21	20	18	10	10
15	30	34	24	29	16	19
16	38	36	31	31	19	20
17	41	43	34	35	24	25
18	73	75	65	63	42	42
19	72	70	60	57	39	39
20	75	92	57	65	42	43
21	82	78	56	56	36	35
22	74	76	52	53	36	38
23	69	69	46	46	35	33
24	70	80	44	53	33	36
25	73	76	45	49	30	29
26-29	289	301	168	192	106	144
30-34	302	324	166	152	110	108
35-39	292	320	151	139	103	105
40-44	334	294	171	156	109	101
45-49	318	309	152	143	115	87
50-54	441	421	191	187	142	142
55-59	394	361	184	155	125	124
60-64	341	344	130	135	102	98
65 or Older	429	417	136	170	110	137

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.14D Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.37	0.36	0.20	0.20	0.14	0.14
12	0.22	0.33	0.14	0.27	0.11	0.14
13	0.40	0.43	0.31	0.32	0.23	0.23
14	0.56	0.48	0.48	0.40	0.25	0.24
15	0.66	0.74	0.55	0.64	0.39	0.43
16	0.79	0.74	0.67	0.66	0.43	0.44
17	0.85	0.88	0.75	0.73	0.53	0.57
18	1.17	1.15	1.10	1.03	0.82	0.76
19	1.20	1.17	1.10	1.05	0.81	0.82
20	1.27	1.40	1.03	1.12	0.83	0.83
21	1.40	1.30	1.09	1.08	0.81	0.73
22	1.24	1.28	1.08	1.05	0.81	0.79
23	1.26	1.26	0.97	1.02	0.80	0.76
24	1.26	1.35	0.98	1.07	0.76	0.78
25	1.32	1.28	0.99	1.06	0.74	0.68
26-29	1.20	1.28	0.93	0.98	0.61	0.77
30-34	1.08	1.15	0.77	0.73	0.53	0.54
35-39	1.11	1.25	0.69	0.69	0.51	0.53
40-44	1.11	1.13	0.77	0.72	0.51	0.49
45-49	1.08	1.10	0.65	0.63	0.50	0.40
50-54	1.57	1.48	0.82	0.80	0.63	0.61
55-59	1.64	1.55	0.92	0.79	0.63	0.64
60-64	1.71	1.53	0.77	0.72	0.61	0.53
65 or Older	1.10	1.04	0.35	0.42	0.28	0.34

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.14P Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2011 Versus 2010

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.1910		0.0386		0.2401	
12	0.0343		0.0222		0.4577	
13	0.8646		0.7279		0.8974	
14	0.2077		0.1507		0.7435	
15	0.2618		0.4873		0.6936	
16	0.3945		1.0000		0.6729	
17	0.9917		0.3936		0.1347	
18	0.1572		0.0839		0.2589	
19	0.0047		0.3151		0.6920	
20	0.8284		0.4050		0.9364	
21	0.7452		0.3460		0.8606	
22	0.1456		0.2281		0.7693	
23	0.0990		0.6915		0.8527	
24	0.4992		0.1129		0.9760	
25	0.8438		0.8179		0.0845	
26-29	0.0134		0.9676		0.1085	
30-34	0.6711		0.7995		1.0000	
35-39	0.9755		0.0172		0.1645	
40-44	0.0287		0.1584		0.1353	
45-49	0.3174		0.3875		0.0109	
50-54	0.8456		0.7347		0.9283	
55-59	0.8925		0.2062		0.6098	
60-64	0.9112		0.8992		0.8204	
65 or Older	0.2160		0.3183		0.3333	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.15C Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	715	684	967	984	990	1,006
12	20	22	17	17	11	12
13	39	39	33	32	20	16
14	44	44	37	37	24	24
15	60	64	54	57	36	36
16	67	67	61	63	42	41
17	74	73	70	66	51	47
18	115	126	108	120	85	94
19	106	121	99	114	84	96
20	118	139	112	135	96	117
21	130	117	128	115	113	106
22	112	135	109	132	95	117
23	109	107	106	99	96	89
24	104	119	99	115	92	105
25	112	120	108	115	98	103
26-29	415	446	387	416	355	371
30-34	455	468	419	431	388	371
35-39	472	492	444	450	408	410
40-44	493	484	457	454	399	407
45-49	504	509	463	451	393	385
50-54	675	690	597	609	537	546
55-59	682	601	584	520	514	459
60-64	634	593	518	532	452	462
65 or Older	444	376	501	488	486	486

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.15D Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2010 and 2011

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.28	0.27	0.38	0.38	0.39	0.39
12	0.51	0.57	0.43	0.45	0.27	0.31
13	0.87	0.82	0.76	0.70	0.48	0.38
14	0.96	0.91	0.85	0.80	0.59	0.55
15	1.04	1.08	1.02	1.05	0.77	0.77
16	1.06	1.00	1.06	1.06	0.90	0.83
17	1.08	1.12	1.11	1.14	1.01	0.97
18	1.07	1.21	1.20	1.29	1.20	1.29
19	0.99	1.13	1.08	1.27	1.24	1.39
20	0.98	1.08	1.06	1.27	1.27	1.40
21	0.82	0.94	0.97	1.02	1.29	1.26
22	0.80	0.84	1.00	1.02	1.23	1.23
23	0.63	0.77	0.87	1.08	1.21	1.24
24	0.77	0.82	0.98	1.04	1.24	1.33
25	0.69	0.73	0.93	0.98	1.19	1.21
26-29	0.69	0.75	1.01	1.04	1.20	1.20
30-34	0.68	0.74	0.95	1.00	1.07	1.09
35-39	0.83	0.76	1.05	1.03	1.24	1.21
40-44	0.72	0.72	1.01	1.01	1.11	1.21
45-49	0.60	0.76	1.00	1.07	1.09	1.19
50-54	0.94	0.93	1.46	1.25	1.61	1.40
55-59	1.08	1.07	1.72	1.62	1.81	1.71
60-64	1.26	1.23	1.84	1.64	1.87	1.75
65 or Older	1.14	0.94	1.29	1.21	1.25	1.21

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.15P Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2011 Versus 2010

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	0.4099		0.6550		0.9707	
12	0.9393		0.7117		0.9720	
13	0.1759		0.0900		0.0244	
14	0.0218		0.1280		0.0944	
15	0.2506		0.0543		0.5378	
16	0.9719		0.8697		0.7592	
17	0.7892		0.4301		0.4226	
18	0.0020		0.0383		0.1727	
19	0.0654		0.0405		0.1338	
20	0.2219		0.4222		0.9655	
21	0.0701		0.0937		0.3259	
22	0.6373		0.8202		0.6363	
23	0.3591		0.0691		0.6285	
24	0.9197		0.6939		0.5618	
25	0.4332		0.8793		0.7635	
26-29	0.5311		0.8414		0.9795	
30-34	0.5687		1.0000		0.3698	
35-39	0.4199		0.5070		0.9395	
40-44	0.2222		0.7834		0.2913	
45-49	0.0049		0.1409		0.0654	
50-54	0.9759		0.6896		0.3496	
55-59	0.3569		0.6654		0.9919	
60-64	0.6817		0.6207		0.8029	
65 or Older	0.1044		0.5811		0.2267	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.16C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2010 and 2011

Age Category	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	990	1,006	752	734	422	423
12	11	12	5	8	1	1
13	20	16	11	10	3	1
14	24	24	17	16	7	4
15	36	36	27	25	13	10
16	42	41	32	30	14	14
17	51	47	41	35	19	16
18	85	94	68	72	32	37
19	84	96	69	71	40	37
20	96	117	77	102	50	56
21	113	106	88	85	52	46
22	95	117	73	90	45	42
23	96	89	72	74	39	38
24	92	105	70	80	38	39
25	98	103	69	81	41	35
26-29	355	371	260	268	135	145
30-34	388	371	284	265	139	152
35-39	408	410	256	247	128	115
40-44	399	407	246	250	136	109
45-49	393	385	250	246	136	128
50-54	537	546	307	294	166	151
55-59	514	459	278	267	152	115
60-64	452	462	225	235	98	123
65 or Older	486	486	263	254	122	110

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.16D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Standard Errors of Percentages, 2010 and 2011

Age Category	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	0.39	0.39	0.30	0.29	0.17	0.16
12	0.27	0.31	0.13	0.22	0.03	0.03
13	0.48	0.38	0.26	0.23	0.07	0.03
14	0.59	0.55	0.42	0.38	0.17	0.10
15	0.77	0.77	0.62	0.56	0.30	0.24
16	0.90	0.83	0.73	0.65	0.33	0.31
17	1.01	0.97	0.85	0.76	0.44	0.37
18	1.20	1.29	1.10	1.16	0.62	0.71
19	1.24	1.39	1.18	1.21	0.83	0.76
20	1.27	1.40	1.25	1.55	0.99	1.06
21	1.29	1.26	1.42	1.39	1.07	0.94
22	1.23	1.23	1.23	1.40	0.96	0.90
23	1.21	1.24	1.30	1.37	0.91	0.87
24	1.24	1.33	1.31	1.36	0.88	0.85
25	1.19	1.21	1.28	1.40	0.97	0.81
26-29	1.20	1.20	1.20	1.20	0.75	0.77
30-34	1.07	1.09	1.08	1.12	0.66	0.75
35-39	1.24	1.21	1.07	1.04	0.62	0.55
40-44	1.11	1.21	0.97	1.05	0.62	0.51
45-49	1.09	1.19	0.97	0.99	0.59	0.57
50-54	1.61	1.40	1.26	1.14	0.74	0.64
55-59	1.81	1.71	1.25	1.26	0.76	0.59
60-64	1.87	1.75	1.29	1.20	0.59	0.66
65 or Older	1.25	1.21	0.68	0.63	0.31	0.27

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.16P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Age Category	Alcohol Use (2010)	Alcohol Use (2011)	Binge Alcohol Use (2010)	Binge Alcohol Use (2011)	Heavy Alcohol Use (2010)	Heavy Alcohol Use (2011)
TOTAL	0.9707		0.2901		0.0256	
12	0.9720		0.1052		0.8182	
13	0.0244		0.3559		0.0533	
14	0.0944		0.1658		0.1709	
15	0.5378		0.1480		0.5880	
16	0.7592		0.6816		0.9963	
17	0.4226		0.8491		0.5820	
18	0.1727		0.4713		0.6256	
19	0.1338		0.0154		0.0532	
20	0.9655		0.8360		0.1621	
21	0.3259		0.2894		0.1496	
22	0.6363		0.7790		0.2234	
23	0.6285		0.5747		0.9126	
24	0.5618		0.9533		0.9299	
25	0.7635		0.8130		0.0106	
26-29	0.9795		0.8757		0.6746	
30-34	0.3698		0.4643		0.9580	
35-39	0.9395		0.2113		0.0526	
40-44	0.2913		0.9040		0.1104	
45-49	0.0654		0.6986		0.2915	
50-54	0.3496		0.5442		0.8174	
55-59	0.9919		0.5270		0.4786	
60-64	0.8029		0.6892		0.2029	
65 or Older	0.2267		0.3932		0.9241	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.