

Table 7.22C GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	118	106	106	119	155	121
AGE GROUP						
12 to 17	8	7	5	8	6	5
18 to 25	37	30	22	25	27	20
26 or Older	111	100	100	117	150	120
GENDER						
Male	88	87	87	98	130	101
Female	70	59	55	63	68	63
PAST YEAR USE						
TOTAL	31	13	31	26	47	37
AGE GROUP						
12 to 17	5	3	3	5	2	4
18 to 25	10	10	8	10	9	8
26 or Older	28	11	30	24	46	35
GENDER						
Male	29	12	30	24	43	36
Female	7	4	9	9	19	6
PAST MONTH USE						
TOTAL	18	9	18	9	29	28
AGE GROUP						
12 to 17	3	3	1	3	*	1
18 to 25	4	8	3	3	3	3
26 or Older	17	*	17	*	29	28
GENDER						
Male	18	9	17	4	23	28
Female	*	*	*	8	18	*

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.22D GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.05	0.04	0.04	0.05	0.06	0.05
AGE GROUP						
12 to 17	0.03	0.03	0.02	0.03	0.02	0.02
18 to 25	0.11	0.09	0.07	0.08	0.08	0.06
26 or Older	0.06	0.05	0.05	0.06	0.08	0.06
GENDER						
Male	0.07	0.07	0.07	0.08	0.11	0.08
Female	0.06	0.05	0.04	0.05	0.05	0.05
PAST YEAR USE						
TOTAL	0.01	0.01	0.01	0.01	0.02	0.01
AGE GROUP						
12 to 17	0.02	0.01	0.01	0.02	0.01	0.02
18 to 25	0.03	0.03	0.02	0.03	0.03	0.02
26 or Older	0.01	0.01	0.02	0.01	0.02	0.02
GENDER						
Male	0.02	0.01	0.02	0.02	0.03	0.03
Female	0.01	0.00	0.01	0.01	0.01	0.00
PAST MONTH USE						
TOTAL	0.01	0.00	0.01	0.00	0.01	0.01
AGE GROUP						
12 to 17	0.01	0.01	0.00	0.01	*	0.00
18 to 25	0.01	0.03	0.01	0.01	0.01	0.01
26 or Older	0.01	*	0.01	*	0.01	0.01
GENDER						
Male	0.01	0.01	0.01	0.00	0.02	0.02
Female	*	*	*	0.01	0.01	*

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.22P GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2011 Versus 2006-2010

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.0578	0.9481	0.3428	0.5979	0.0916	
AGE GROUP						
12 to 17	0.0210	0.6540	0.6558	0.1569	0.5522	
18 to 25	0.0000	0.0000	0.0290	0.0004	0.0211	
26 or Older	0.7746	0.2423	0.1563	0.7885	0.1940	
GENDER						
Male	0.6974	0.5422	0.1616	0.9610	0.1328	
Female	0.0035	0.4344	0.6742	0.3760	0.3658	
PAST YEAR USE						
TOTAL	0.5643	0.3417	0.9122	0.8751	0.2835	
AGE GROUP						
12 to 17	0.4610	0.3836	0.5394	0.6073	0.0857	
18 to 25	0.0632	0.1047	0.9883	0.5053	0.5314	
26 or Older	0.9666	0.1489	0.9604	0.6406	0.2707	
GENDER						
Male	0.7729	0.3048	0.7904	0.7152	0.6092	
Female	0.1473	0.7931	0.5302	0.4114	0.0870	
PAST MONTH USE						
TOTAL	0.6824	0.5314	0.6665	0.3838	0.4936	
AGE GROUP						
12 to 17	0.2244	0.4855	0.9138	0.2552	*	
18 to 25	0.4861	0.0428	0.6696	0.6835	0.4726	
26 or Older	0.5298	*	0.6273	*	0.5316	
GENDER						
Male	0.6030	0.4944	0.5320	0.2198	0.9439	
Female	*	*	*	*	*	

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

**Table 7.23C Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
Standard Errors of Numbers in Thousands, 2006-2011**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	165	170	180	220	257	232
AGE GROUP						
12 to 17	34	37	31	32	37	34
18 to 25	98	97	95	112	117	113
26 or Older	112	116	139	164	204	194
GENDER						
Male	123	125	127	174	197	161
Female	96	102	118	113	139	158
PAST YEAR USE						
TOTAL	92	99	102	135	128	134
AGE GROUP						
12 to 17	27	28	25	25	29	27
18 to 25	66	69	65	82	84	86
26 or Older	53	58	71	90	76	91
GENDER						
Male	62	72	74	107	97	93
Female	60	61	65	70	67	92
PAST MONTH USE						
TOTAL	43	45	55	65	73	65
AGE GROUP						
12 to 17	14	17	13	13	16	12
18 to 25	32	37	31	36	46	41
26 or Older	24	15	42	53	49	47
GENDER						
Male	33	32	41	57	57	45
Female	27	30	33	35	37	42

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

**Table 7.23D Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
Standard Errors of Percentages, 2006-2011**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.07	0.07	0.07	0.09	0.10	0.09
AGE GROUP						
12 to 17	0.14	0.15	0.12	0.13	0.15	0.13
18 to 25	0.30	0.30	0.29	0.33	0.34	0.33
26 or Older	0.06	0.06	0.07	0.08	0.10	0.10
GENDER						
Male	0.10	0.10	0.10	0.14	0.16	0.13
Female	0.08	0.08	0.09	0.09	0.11	0.12
PAST YEAR USE						
TOTAL	0.04	0.04	0.04	0.05	0.05	0.05
AGE GROUP						
12 to 17	0.11	0.11	0.10	0.10	0.12	0.11
18 to 25	0.20	0.21	0.20	0.24	0.25	0.25
26 or Older	0.03	0.03	0.04	0.05	0.04	0.05
GENDER						
Male	0.05	0.06	0.06	0.09	0.08	0.07
Female	0.05	0.05	0.05	0.05	0.05	0.07
PAST MONTH USE						
TOTAL	0.02	0.02	0.02	0.03	0.03	0.03
AGE GROUP						
12 to 17	0.05	0.07	0.05	0.05	0.06	0.05
18 to 25	0.10	0.11	0.09	0.11	0.13	0.12
26 or Older	0.01	0.01	0.02	0.03	0.03	0.02
GENDER						
Male	0.03	0.03	0.03	0.05	0.05	0.04
Female	0.02	0.02	0.03	0.03	0.03	0.03

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

**Table 7.23P Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
P Values from Tests of Differences of Percentages, 2011 Versus 2006-2010**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.0000	0.0000	0.0000	0.0000	0.6886	
AGE GROUP						
12 to 17	0.1495	0.3539	0.8942	0.6808	0.2279	
18 to 25	0.0000	0.0000	0.0000	0.0601	0.9294	
26 or Older	0.0000	0.0000	0.0000	0.0001	0.4617	
GENDER						
Male	0.0000	0.0000	0.0000	0.3189	0.0922	
Female	0.0000	0.0000	0.0000	0.0000	0.0049	
PAST YEAR USE						
TOTAL	0.0000	0.0000	0.0000	0.2222	0.4654	
AGE GROUP						
12 to 17	0.6349	0.6500	0.4902	0.8896	0.6939	
18 to 25	0.0000	0.0000	0.0000	0.3105	0.8591	
26 or Older	0.0000	0.0000	0.0010	0.3928	0.1386	
GENDER						
Male	0.0000	0.0000	0.0000	0.9819	0.9595	
Female	0.0003	0.0000	0.0003	0.0432	0.2690	
PAST MONTH USE						
TOTAL	0.0007	0.0002	0.0188	0.6638	0.2717	
AGE GROUP						
12 to 17	0.7618	0.6350	0.2562	0.9164	0.7161	
18 to 25	0.0072	0.0180	0.0148	0.7946	0.8004	
26 or Older	0.0205	0.0006	0.4067	0.4343	0.2675	
GENDER						
Male	0.0008	0.0007	0.0162	0.7901	0.4523	
Female	0.1727	0.0697	0.3117	0.6950	0.3892	

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

**Table 7.24C Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
Standard Errors of Numbers in Thousands, 2006-2011**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	206	216	233	265	289	264
AGE GROUP						
12 to 17	24	24	21	25	21	22
18 to 25	61	54	54	55	62	54
26 or Older	190	207	221	254	279	258
GENDER						
Male	143	149	149	185	213	166
Female	140	162	181	178	193	204
PAST YEAR USE						
TOTAL	138	140	151	149	184	138
AGE GROUP						
12 to 17	18	16	16	18	15	17
18 to 25	42	38	35	37	36	32
26 or Older	130	133	144	141	177	133
GENDER						
Male	102	98	95	106	142	86
Female	93	98	112	103	114	104
PAST MONTH USE						
TOTAL	85	54	58	83	133	72
AGE GROUP						
12 to 17	10	7	6	7	8	7
18 to 25	17	16	18	18	16	16
26 or Older	81	51	55	79	132	71
GENDER						
Male	70	27	38	49	113	53
Female	41	45	44	66	73	49

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

**Table 7.24D Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
Standard Errors of Percentages, 2006-2011**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.08	0.09	0.09	0.11	0.11	0.10
AGE GROUP						
12 to 17	0.10	0.09	0.08	0.10	0.09	0.09
18 to 25	0.19	0.17	0.16	0.16	0.18	0.16
26 or Older	0.10	0.11	0.12	0.13	0.14	0.13
GENDER						
Male	0.12	0.12	0.12	0.15	0.17	0.13
Female	0.11	0.13	0.14	0.14	0.15	0.15
PAST YEAR USE						
TOTAL	0.06	0.06	0.06	0.06	0.07	0.05
AGE GROUP						
12 to 17	0.07	0.06	0.06	0.07	0.06	0.07
18 to 25	0.13	0.12	0.11	0.11	0.11	0.09
26 or Older	0.07	0.07	0.07	0.07	0.09	0.07
GENDER						
Male	0.09	0.08	0.08	0.09	0.12	0.07
Female	0.07	0.08	0.09	0.08	0.09	0.08
PAST MONTH USE						
TOTAL	0.03	0.02	0.02	0.03	0.05	0.03
AGE GROUP						
12 to 17	0.04	0.03	0.02	0.03	0.03	0.03
18 to 25	0.05	0.05	0.05	0.05	0.05	0.05
26 or Older	0.04	0.03	0.03	0.04	0.07	0.04
GENDER						
Male	0.06	0.02	0.03	0.04	0.09	0.04
Female	0.03	0.04	0.03	0.05	0.06	0.04

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

**Table 7.24P Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
P Values from Tests of Differences of Percentages, 2011 Versus 2006-2010**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.0000	0.0388	0.2173	0.5078	0.2878	
AGE GROUP						
12 to 17	0.2343	0.5084	0.8902	0.5718	0.2473	
18 to 25	0.1244	0.7428	0.1157	0.0096	0.0252	
26 or Older	0.0000	0.0252	0.1134	0.9244	0.4866	
GENDER						
Male	0.0353	0.2653	0.1835	0.2541	0.0395	
Female	0.0001	0.0814	0.6337	0.9022	0.6230	
PAST YEAR USE						
TOTAL	0.2593	0.2691	0.2673	0.1357	0.0913	
AGE GROUP						
12 to 17	0.3564	0.9919	0.6534	0.7995	0.2965	
18 to 25	0.0008	0.0213	0.0137	0.0012	0.1178	
26 or Older	0.8912	0.5647	0.5354	0.4280	0.1311	
GENDER						
Male	0.0663	0.0688	0.1622	0.0198	0.0204	
Female	0.8523	0.9042	0.7466	0.9357	0.9342	
PAST MONTH USE						
TOTAL	0.4612	0.3002	0.7500	0.2157	0.0288	
AGE GROUP						
12 to 17	0.3907	0.6168	0.5028	0.9652	0.9224	
18 to 25	0.6693	0.3544	0.0594	0.0846	0.4373	
26 or Older	0.5637	0.2097	0.4544	0.3698	0.0359	
GENDER						
Male	0.1230	0.2776	0.9251	0.1410	0.0392	
Female	0.3920	0.6672	0.5825	0.7046	0.3709	

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.25C Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	283	260	274	258	288	254
AGE GROUP						
12 to 17	40	41	36	36	39	37
18 to 25	77	74	71	83	80	79
26 or Older	264	244	260	235	269	242
GENDER						
Male	229	203	223	193	236	215
Female	151	163	169	163	157	145
PAST YEAR USE						
TOTAL	99	85	98	111	162	111
AGE GROUP						
12 to 17	30	29	26	25	27	25
18 to 25	38	37	31	35	37	40
26 or Older	87	72	91	100	158	100
GENDER						
Male	76	55	81	75	147	86
Female	64	66	56	80	70	71
PAST MONTH USE						
TOTAL	51	46	53	60	67	60
AGE GROUP						
12 to 17	13	15	14	13	12	11
18 to 25	15	19	16	17	21	15
26 or Older	47	39	49	57	63	56
GENDER						
Male	45	34	43	37	52	43
Female	26	31	31	47	43	42

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.25D Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.11	0.10	0.11	0.10	0.11	0.10
AGE GROUP						
12 to 17	0.16	0.16	0.15	0.15	0.16	0.15
18 to 25	0.24	0.23	0.22	0.25	0.23	0.23
26 or Older	0.14	0.13	0.14	0.12	0.14	0.12
GENDER						
Male	0.19	0.17	0.18	0.16	0.19	0.17
Female	0.12	0.13	0.13	0.13	0.12	0.11
PAST YEAR USE						
TOTAL	0.04	0.03	0.04	0.04	0.06	0.04
AGE GROUP						
12 to 17	0.12	0.12	0.10	0.10	0.11	0.10
18 to 25	0.12	0.11	0.10	0.11	0.11	0.12
26 or Older	0.05	0.04	0.05	0.05	0.08	0.05
GENDER						
Male	0.06	0.05	0.07	0.06	0.12	0.07
Female	0.05	0.05	0.04	0.06	0.05	0.05
PAST MONTH USE						
TOTAL	0.02	0.02	0.02	0.02	0.03	0.02
AGE GROUP						
12 to 17	0.05	0.06	0.05	0.05	0.05	0.05
18 to 25	0.05	0.06	0.05	0.05	0.06	0.04
26 or Older	0.02	0.02	0.03	0.03	0.03	0.03
GENDER						
Male	0.04	0.03	0.04	0.03	0.04	0.03
Female	0.02	0.02	0.02	0.04	0.03	0.03

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.25P Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2011 Versus 2006-2010

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.2541	0.4332	0.8176	0.8090	0.2052	
AGE GROUP						
12 to 17	0.0001	0.0101	0.0748	0.2853	0.1492	
18 to 25	0.7714	0.2368	0.9762	0.3670	0.5205	
26 or Older	0.4910	0.3811	1.0000	0.4567	0.3639	
GENDER						
Male	0.2299	0.2738	0.6459	0.2312	0.2452	
Female	0.8012	0.8628	0.3267	0.2368	0.6622	
PAST YEAR USE						
TOTAL	0.6180	0.2308	0.5194	0.6396	0.2435	
AGE GROUP						
12 to 17	0.0001	0.0105	0.0434	0.1969	0.1512	
18 to 25	0.7738	0.8044	0.4790	0.6677	0.6273	
26 or Older	0.4487	0.0342	0.3171	0.4853	0.2729	
GENDER						
Male	0.3034	0.3782	0.9621	0.5615	0.0762	
Female	0.6839	0.4243	0.3043	0.9408	0.4360	
PAST MONTH USE						
TOTAL	0.5574	0.7693	0.7952	0.7583	0.3855	
AGE GROUP						
12 to 17	0.0104	0.0157	0.1240	0.0604	0.2583	
18 to 25	0.4115	0.0983	0.4425	0.1634	0.2008	
26 or Older	0.7892	0.3249	0.7394	0.6320	0.7616	
GENDER						
Male	0.1757	0.6662	0.5548	0.8608	0.3665	
Female	0.4452	0.9878	0.7670	0.7971	0.7709	

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.26C Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	145	140	136	151	166	152
AGE GROUP						
12 to 17	14	12	12	12	15	13
18 to 25	50	47	40	38	39	41
26 or Older	133	129	126	145	160	147
GENDER						
Male	123	115	108	124	152	120
Female	69	68	73	76	64	90
PAST YEAR USE						
TOTAL	38	31	28	32	36	49
AGE GROUP						
12 to 17	8	7	9	8	11	11
18 to 25	14	19	14	16	22	26
26 or Older	34	24	23	27	26	40
GENDER						
Male	35	27	25	23	33	39
Female	14	16	11	23	14	28
PAST MONTH USE						
TOTAL	18	11	7	24	22	26
AGE GROUP						
12 to 17	4	5	5	6	5	9
18 to 25	6	6	6	9	8	9
26 or Older	16	8	*	21	20	22
GENDER						
Male	14	10	6	13	21	12
Female	10	4	4	20	8	23

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Ketamine is also known as "Special K" or "Super K."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.26D Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.06	0.06	0.05	0.06	0.07	0.06
AGE GROUP						
12 to 17	0.05	0.05	0.05	0.05	0.06	0.05
18 to 25	0.15	0.14	0.12	0.11	0.11	0.12
26 or Older	0.07	0.07	0.07	0.07	0.08	0.07
GENDER						
Male	0.10	0.10	0.09	0.10	0.12	0.10
Female	0.05	0.05	0.06	0.06	0.05	0.07
PAST YEAR USE						
TOTAL	0.02	0.01	0.01	0.01	0.01	0.02
AGE GROUP						
12 to 17	0.03	0.03	0.04	0.03	0.05	0.04
18 to 25	0.04	0.06	0.04	0.05	0.07	0.07
26 or Older	0.02	0.01	0.01	0.01	0.01	0.02
GENDER						
Male	0.03	0.02	0.02	0.02	0.03	0.03
Female	0.01	0.01	0.01	0.02	0.01	0.02
PAST MONTH USE						
TOTAL	0.01	0.00	0.00	0.01	0.01	0.01
AGE GROUP						
12 to 17	0.02	0.02	0.02	0.03	0.02	0.04
18 to 25	0.02	0.02	0.02	0.03	0.02	0.03
26 or Older	0.01	0.00	*	0.01	0.01	0.01
GENDER						
Male	0.01	0.01	0.00	0.01	0.02	0.01
Female	0.01	0.00	0.00	0.02	0.01	0.02

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Ketamine is also known as "Special K" or "Super K."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.26P Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2011 Versus 2006-2010

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.9574	0.6354	0.4281	0.9403	0.7634	
AGE GROUP						
12 to 17	0.7407	0.3235	0.4609	0.6196	0.7550	
18 to 25	0.0000	0.0000	0.0081	0.0614	0.3287	
26 or Older	0.0314	0.0417	0.1207	0.6000	0.9419	
GENDER						
Male	0.9127	0.7073	0.3311	0.8868	0.2703	
Female	0.8019	0.7228	0.9629	0.7253	0.1707	
PAST YEAR USE						
TOTAL	0.0515	0.0494	0.0018	0.0288	0.1528	
AGE GROUP						
12 to 17	0.1005	0.0872	0.2035	0.1958	0.9046	
18 to 25	0.0117	0.4519	0.0054	0.0575	0.6617	
26 or Older	0.5771	0.1413	0.0756	0.2701	0.1465	
GENDER						
Male	0.1512	0.1077	0.0151	0.0184	0.3537	
Female	0.1355	0.2208	0.0299	0.5502	0.1813	
PAST MONTH USE						
TOTAL	0.4264	0.1427	0.0440	0.8685	0.5322	
AGE GROUP						
12 to 17	0.1582	0.1920	0.1344	0.4490	0.2598	
18 to 25	0.2208	0.3314	0.3085	0.8768	0.6226	
26 or Older	0.9727	0.4514	*	0.6045	0.9149	
GENDER						
Male	0.9343	0.5385	0.1292	0.7495	0.9280	
Female	0.3247	0.1702	0.1451	0.9945	0.2928	

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Ketamine is also known as "Special K" or "Super K."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.27C DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	62	71	74	79	101	84
AGE GROUP						
12 to 17	9	10	10	10	10	10
18 to 25	29	24	28	33	36	36
26 or Older	54	65	68	73	94	74
GENDER						
Male	55	66	60	71	87	70
Female	29	23	42	33	53	50
PAST YEAR USE						
TOTAL	21	16	27	29	35	32
AGE GROUP						
12 to 17	4	6	8	8	8	7
18 to 25	15	12	14	19	16	24
26 or Older	14	7	21	21	31	19
GENDER						
Male	21	14	23	19	28	27
Female	5	6	14	22	21	16
PAST MONTH USE						
TOTAL	7	*	18	20	28	17
AGE GROUP						
12 to 17	*	*	4	4	7	3
18 to 25	5	5	9	4	7	8
26 or Older	*	*	15	19	27	14
GENDER						
Male	6	4	17	8	22	8
Female	4	*	*	18	18	15

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: DMT is also known as dimethyltryptamine.

NOTE: AMT is also known as alpha-methyltryptamine.

NOTE: 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy".

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.27D DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.03	0.03	0.03	0.03	0.04	0.03
AGE GROUP						
12 to 17	0.04	0.04	0.04	0.04	0.04	0.04
18 to 25	0.09	0.07	0.09	0.10	0.10	0.10
26 or Older	0.03	0.03	0.04	0.04	0.05	0.04
GENDER						
Male	0.05	0.05	0.05	0.06	0.07	0.06
Female	0.02	0.02	0.03	0.03	0.04	0.04
PAST YEAR USE						
TOTAL	0.01	0.01	0.01	0.01	0.01	0.01
AGE GROUP						
12 to 17	0.02	0.02	0.03	0.03	0.03	0.03
18 to 25	0.05	0.04	0.04	0.06	0.05	0.07
26 or Older	0.01	0.00	0.01	0.01	0.02	0.01
GENDER						
Male	0.02	0.01	0.02	0.02	0.02	0.02
Female	0.00	0.00	0.01	0.02	0.02	0.01
PAST MONTH USE						
TOTAL	0.00	*	0.01	0.01	0.01	0.01
AGE GROUP						
12 to 17	*	*	0.02	0.02	0.03	0.01
18 to 25	0.02	0.01	0.03	0.01	0.02	0.02
26 or Older	*	*	0.01	0.01	0.01	0.01
GENDER						
Male	0.00	0.00	0.01	0.01	0.02	0.01
Female	0.00	*	*	0.01	0.01	0.01

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: DMT is also known as dimethyltryptamine.

NOTE: AMT is also known as alpha-methyltryptamine.

NOTE: 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy".

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.27P DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2011 Versus 2006-2010

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.0007	0.0008	0.0046	0.1317	0.2648	
AGE GROUP						
12 to 17	0.1521	0.1376	0.2703	0.6828	0.8839	
18 to 25	0.0006	0.0000	0.0000	0.0290	0.4638	
26 or Older	0.0546	0.1202	0.2474	0.5448	0.1356	
GENDER						
Male	0.0013	0.0035	0.0005	0.0972	0.5565	
Female	0.2455	0.0880	0.8903	0.8716	0.2763	
PAST YEAR USE						
TOTAL	0.0001	0.0000	0.0076	0.1240	0.5583	
AGE GROUP						
12 to 17	0.0160	0.0880	0.5503	0.8061	0.8093	
18 to 25	0.0001	0.0000	0.0001	0.0222	0.0202	
26 or Older	0.3755	0.0824	0.9326	0.9882	0.3623	
GENDER						
Male	0.0004	0.0000	0.0049	0.0047	0.1862	
Female	0.0706	0.0581	0.5713	0.3396	0.4202	
PAST MONTH USE						
TOTAL	0.1563	*	0.9684	0.6478	0.2827	
AGE GROUP						
12 to 17	*	*	0.8950	0.3845	0.1630	
18 to 25	0.3310	0.1218	0.6861	0.1612	0.7141	
26 or Older	*	*	0.8019	0.4018	0.3402	
GENDER						
Male	0.1804	0.0329	0.4395	0.7168	0.3100	
Female	0.4188	*	*	0.4999	0.6194	

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: DMT is also known as dimethyltryptamine.

NOTE: AMT is also known as alpha-methyltryptamine.

NOTE: 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy".

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.28C Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	92	110	129	159	189	175
AGE GROUP						
12 to 17	20	23	27	28	28	25
18 to 25	62	68	75	89	95	92
26 or Older	59	78	98	120	155	139
GENDER						
Male	81	91	104	142	159	142
Female	34	54	62	63	99	97
PAST YEAR USE						
TOTAL	48	68	70	72	77	52
AGE GROUP						
12 to 17	16	18	23	22	22	19
18 to 25	38	45	47	47	42	39
26 or Older	22	43	43	46	59	29
GENDER						
Male	41	55	55	58	73	43
Female	21	34	37	37	28	28
PAST MONTH USE						
TOTAL	18	26	24	28	28	20
AGE GROUP						
12 to 17	7	9	10	9	9	8
18 to 25	16	21	21	15	17	18
26 or Older	*	11	8	22	20	7
GENDER						
Male	17	22	22	19	27	17
Female	7	11	8	20	9	10

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.28D Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.04	0.04	0.05	0.06	0.07	0.07
AGE GROUP						
12 to 17	0.08	0.09	0.11	0.12	0.12	0.10
18 to 25	0.19	0.21	0.23	0.26	0.28	0.27
26 or Older	0.03	0.04	0.05	0.06	0.08	0.07
GENDER						
Male	0.07	0.08	0.09	0.12	0.13	0.11
Female	0.03	0.04	0.05	0.05	0.08	0.07
PAST YEAR USE						
TOTAL	0.02	0.03	0.03	0.03	0.03	0.02
AGE GROUP						
12 to 17	0.06	0.07	0.09	0.09	0.09	0.07
18 to 25	0.12	0.14	0.14	0.14	0.12	0.11
26 or Older	0.01	0.02	0.02	0.02	0.03	0.01
GENDER						
Male	0.03	0.05	0.05	0.05	0.06	0.03
Female	0.02	0.03	0.03	0.03	0.02	0.02
PAST MONTH USE						
TOTAL	0.01	0.01	0.01	0.01	0.01	0.01
AGE GROUP						
12 to 17	0.03	0.03	0.04	0.04	0.04	0.03
18 to 25	0.05	0.06	0.06	0.04	0.05	0.05
26 or Older	*	0.01	0.00	0.01	0.01	0.00
GENDER						
Male	0.01	0.02	0.02	0.02	0.02	0.01
Female	0.01	0.01	0.01	0.02	0.01	0.01

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.28P Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: *P* Values from Tests of Differences of Percentages, 2011 Versus 2006-2010

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.0000	0.0000	0.0000	0.0007	0.8964	
AGE GROUP						
12 to 17	0.0000	0.0132	0.5822	0.1956	0.3532	
18 to 25	0.0000	0.0000	0.0000	0.0394	0.9733	
26 or Older	0.0000	0.0000	0.0000	0.0016	0.8909	
GENDER						
Male	0.0000	0.0000	0.0000	0.1654	0.3234	
Female	0.0000	0.0000	0.0000	0.0000	0.1514	
PAST YEAR USE						
TOTAL	0.3065	0.0454	0.0000	0.0000	0.0003	
AGE GROUP						
12 to 17	0.0111	0.4715	0.0523	0.1743	0.3254	
18 to 25	0.3206	0.0062	0.0001	0.0000	0.0161	
26 or Older	0.0347	0.6643	0.1601	0.0306	0.0089	
GENDER						
Male	0.9650	0.0561	0.0001	0.0000	0.0003	
Female	0.0372	0.3732	0.0236	0.0762	0.5068	
PAST MONTH USE						
TOTAL	0.8102	0.0300	0.0331	0.0700	0.3952	
AGE GROUP						
12 to 17	0.3455	0.7261	0.0902	0.2588	0.5122	
18 to 25	0.3287	0.0483	0.0773	0.5981	0.7051	
26 or Older	*	0.3781	0.7699	0.1160	0.5854	
GENDER						
Male	0.3694	0.0317	0.0143	0.1107	0.2015	
Female	0.2093	0.5240	0.8736	0.3633	0.3716	

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.