

Table 7.29C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2011

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL										
TOBACCO PRODUCTS¹	832	803	794	845	850	844	871	831	852	846
Cigarettes	795	789	768	777	807	809	810	797	780	816
Smokeless Tobacco	279	284	245	267	284	283	312	291	314	278
Cigars	350	331	323	354	352	337	366	361	363	373
Pipe Tobacco	166	131	147	146	167	170	147	151	147	144
ALCOHOL	989	919	957	975	968	1,019	983	957	990	1,006
Binge Alcohol Use ²	732	682	698	701	746	770	735	783	752	734
Heavy Alcohol Use ²	392	380	382	379	427	429	414	406	422	423
MALE										
TOBACCO PRODUCTS¹	572	573	570	576	585	598	628	602	626	581
Cigarettes	527	545	537	515	545	577	565	565	536	537
Smokeless Tobacco	264	256	234	241	273	252	296	283	302	264
Cigars	327	305	290	324	320	312	333	336	321	336
Pipe Tobacco	139	127	139	141	160	168	131	144	137	135
ALCOHOL	656	599	626	621	610	666	632	651	675	674
Binge Alcohol Use ²	540	517	513	529	559	585	586	605	577	567
Heavy Alcohol Use ²	346	321	319	321	354	374	357	349	353	355
FEMALE										
TOBACCO PRODUCTS¹	513	507	497	543	511	511	558	505	535	542
Cigarettes	501	488	491	538	503	493	553	498	526	532
Smokeless Tobacco	100	126	65	87	76	113	75	51	78	82
Cigars	114	123	122	112	131	116	114	130	153	142
Pipe Tobacco	90	35	43	42	46	30	55	37	54	46
ALCOHOL	644	600	664	637	645	676	673	687	635	684
Binge Alcohol Use ²	415	382	390	402	399	395	398	441	441	441
Heavy Alcohol Use ²	177	174	174	163	169	171	192	193	201	207

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2011.

Table 7.29D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Percentages, 2002-2011

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL										
TOBACCO PRODUCTS¹	0.35	0.34	0.33	0.35	0.35	0.34	0.35	0.33	0.34	0.33
Cigarettes	0.34	0.33	0.32	0.32	0.33	0.33	0.32	0.32	0.31	0.32
Smokeless Tobacco	0.12	0.12	0.10	0.11	0.12	0.11	0.12	0.12	0.12	0.11
Cigars	0.15	0.14	0.13	0.15	0.14	0.14	0.15	0.14	0.14	0.14
Pipe Tobacco	0.07	0.06	0.06	0.06	0.07	0.07	0.06	0.06	0.06	0.06
ALCOHOL	0.42	0.39	0.40	0.40	0.39	0.41	0.39	0.38	0.39	0.39
Binge Alcohol Use ²	0.31	0.29	0.29	0.29	0.30	0.31	0.29	0.31	0.30	0.29
Heavy Alcohol Use ²	0.17	0.16	0.16	0.16	0.17	0.17	0.17	0.16	0.17	0.16
MALE										
TOBACCO PRODUCTS¹	0.50	0.50	0.49	0.49	0.49	0.50	0.52	0.49	0.51	0.47
Cigarettes	0.46	0.47	0.46	0.44	0.46	0.48	0.47	0.46	0.43	0.43
Smokeless Tobacco	0.23	0.22	0.20	0.20	0.23	0.21	0.24	0.23	0.25	0.21
Cigars	0.29	0.27	0.25	0.27	0.27	0.26	0.27	0.27	0.26	0.27
Pipe Tobacco	0.12	0.11	0.12	0.12	0.13	0.14	0.11	0.12	0.11	0.11
ALCOHOL	0.58	0.52	0.54	0.53	0.51	0.55	0.52	0.53	0.55	0.54
Binge Alcohol Use ²	0.48	0.45	0.44	0.45	0.47	0.49	0.48	0.49	0.47	0.45
Heavy Alcohol Use ²	0.30	0.28	0.27	0.27	0.30	0.31	0.29	0.29	0.29	0.28
FEMALE										
TOBACCO PRODUCTS¹	0.42	0.41	0.40	0.43	0.40	0.40	0.43	0.39	0.41	0.41
Cigarettes	0.41	0.40	0.40	0.43	0.40	0.39	0.43	0.38	0.40	0.40
Smokeless Tobacco	0.08	0.10	0.05	0.07	0.06	0.09	0.06	0.04	0.06	0.06
Cigars	0.09	0.10	0.10	0.09	0.10	0.09	0.09	0.10	0.12	0.11
Pipe Tobacco	0.07	0.03	0.03	0.03	0.04	0.02	0.04	0.03	0.04	0.03
ALCOHOL	0.53	0.49	0.54	0.51	0.51	0.53	0.52	0.53	0.49	0.51
Binge Alcohol Use ²	0.34	0.31	0.31	0.32	0.32	0.31	0.31	0.34	0.34	0.33
Heavy Alcohol Use ²	0.15	0.14	0.14	0.13	0.13	0.13	0.15	0.15	0.15	0.16

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2011.

Table 7.29P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: P Values from Tests of Differences of Percentages, 2011 Versus 2002-2010

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL										
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0062	0.0299	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0040	0.0308	
Smokeless Tobacco	0.4870	0.7566	0.1286	0.7947	0.3758	0.7460	0.0816	0.1627	0.0585	
Cigars	0.0394	0.0431	0.0003	0.0031	0.0042	0.0590	0.1488	0.1237	0.2401	
Pipe Tobacco	0.5247	0.0587	0.4217	0.3886	0.2019	0.9564	0.3447	0.9439	0.8042	
ALCOHOL	0.1514	0.0017	0.0073	0.9484	0.1389	0.2895	0.7310	0.8518	0.9707	
Binge Alcohol Use ²	0.5920	0.9506	0.7955	0.9955	0.3479	0.1019	0.0834	0.0130	0.2901	
Heavy Alcohol Use ²	0.0124	0.0058	0.0007	0.0570	0.0023	0.0023	0.0005	0.0042	0.0256	
MALE										
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0012	0.0746	0.0315	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0019	0.1292	0.0713	
Smokeless Tobacco	0.4926	0.9679	0.1922	0.8032	0.2008	0.5955	0.0424	0.0820	0.0519	
Cigars	0.0020	0.0246	0.0000	0.0002	0.0029	0.0102	0.0239	0.1047	0.2828	
Pipe Tobacco	0.6343	0.2765	0.8487	0.2005	0.0709	0.5606	0.2902	0.6354	0.8822	
ALCOHOL	0.4171	0.4435	0.8199	0.0695	0.7307	0.8875	0.2298	0.3006	0.4468	
Binge Alcohol Use ²	0.0640	0.1420	0.0874	0.3941	0.0424	0.0108	0.0137	0.0141	0.1773	
Heavy Alcohol Use ²	0.0009	0.0092	0.0013	0.0133	0.0010	0.0017	0.0010	0.0181	0.0569	
FEMALE										
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0003	0.0001	0.0001	0.0165	0.0098	0.0258	0.3996	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0036	0.0027	0.0055	0.1714	
Smokeless Tobacco	0.8360	0.4757	0.2926	0.8166	0.1603	0.5420	0.4073	0.1378	0.6831	
Cigars	0.0315	0.9420	0.3960	0.1543	0.8378	0.1636	0.0505	0.9536	0.7643	
Pipe Tobacco	0.6554	0.0042	0.0388	0.2534	0.1118	0.0070	0.9898	0.1632	0.8332	
ALCOHOL	0.0032	0.0000	0.0000	0.0962	0.0119	0.1325	0.0897	0.4275	0.4196	
Binge Alcohol Use ²	0.1514	0.0416	0.0710	0.2301	0.2517	0.4803	0.6293	0.3499	0.8566	
Heavy Alcohol Use ²	0.4512	0.2308	0.1424	0.7093	0.5120	0.4340	0.1703	0.1083	0.2564	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2011.

Table 7.30C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Numbers in Thousands, 2002-2011

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL										
TOBACCO PRODUCTS¹	81	80	80	78	72	76	69	71	69	68
Cigarettes	74	73	76	71	66	67	61	64	63	60
Smokeless Tobacco	27	30	32	29	31	36	30	32	30	30
Cigars	47	44	44	46	41	45	41	39	36	41
Pipe Tobacco	14	18	19	17	18	19	17	21	17	18
ALCOHOL	80	82	81	81	82	87	79	79	80	77
Binge Alcohol Use ²	67	67	73	65	69	69	57	62	62	55
Heavy Alcohol Use ²	31	33	36	32	33	37	30	30	28	25
MALE										
TOBACCO PRODUCTS¹	59	56	55	59	54	57	50	52	52	53
Cigarettes	53	47	52	53	46	48	42	44	45	46
Smokeless Tobacco	26	29	30	28	29	34	28	30	28	28
Cigars	37	36	37	38	34	37	34	33	31	31
Pipe Tobacco	13	17	15	14	15	16	12	18	14	14
ALCOHOL	55	57	59	57	57	61	53	58	57	53
Binge Alcohol Use ²	46	46	52	48	47	51	41	48	44	42
Heavy Alcohol Use ²	24	23	27	27	26	28	23	23	23	22
FEMALE										
TOBACCO PRODUCTS¹	51	52	53	50	47	48	46	46	43	43
Cigarettes	49	50	52	48	46	46	44	44	40	40
Smokeless Tobacco	9	6	11	8	10	8	10	9	12	10
Cigars	26	23	25	24	24	22	21	25	20	24
Pipe Tobacco	9	7	13	11	10	11	12	11	10	12
ALCOHOL	58	60	59	58	59	61	56	52	54	53
Binge Alcohol Use ²	46	46	50	43	49	45	42	40	43	37
Heavy Alcohol Use ²	19	23	23	20	21	23	20	19	17	15

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2011.

Table 7.30D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Percentages, 2002-2011

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL										
TOBACCO PRODUCTS¹	0.33	0.32	0.32	0.31	0.29	0.30	0.28	0.29	0.28	0.27
Cigarettes	0.30	0.29	0.30	0.28	0.26	0.27	0.25	0.26	0.26	0.24
Smokeless Tobacco	0.11	0.12	0.13	0.12	0.12	0.14	0.12	0.13	0.12	0.12
Cigars	0.19	0.17	0.18	0.18	0.16	0.18	0.16	0.16	0.15	0.16
Pipe Tobacco	0.06	0.07	0.08	0.07	0.07	0.08	0.07	0.09	0.07	0.07
ALCOHOL	0.32	0.33	0.32	0.32	0.32	0.34	0.32	0.32	0.33	0.31
Binge Alcohol Use ²	0.27	0.27	0.29	0.26	0.27	0.27	0.23	0.25	0.25	0.22
Heavy Alcohol Use ²	0.12	0.13	0.14	0.13	0.13	0.15	0.12	0.12	0.11	0.10
MALE										
TOBACCO PRODUCTS¹	0.47	0.44	0.43	0.46	0.41	0.45	0.39	0.42	0.42	0.41
Cigarettes	0.42	0.37	0.40	0.41	0.36	0.37	0.33	0.35	0.36	0.36
Smokeless Tobacco	0.21	0.23	0.23	0.22	0.22	0.26	0.22	0.24	0.23	0.22
Cigars	0.30	0.28	0.28	0.29	0.26	0.29	0.27	0.27	0.25	0.25
Pipe Tobacco	0.10	0.13	0.11	0.11	0.12	0.12	0.10	0.14	0.11	0.11
ALCOHOL	0.43	0.45	0.45	0.44	0.44	0.47	0.41	0.46	0.46	0.42
Binge Alcohol Use ²	0.37	0.36	0.40	0.37	0.36	0.40	0.32	0.38	0.35	0.33
Heavy Alcohol Use ²	0.19	0.18	0.21	0.21	0.20	0.22	0.18	0.18	0.18	0.17
FEMALE										
TOBACCO PRODUCTS¹	0.42	0.42	0.43	0.40	0.38	0.39	0.38	0.38	0.36	0.35
Cigarettes	0.40	0.41	0.42	0.39	0.37	0.37	0.36	0.37	0.34	0.33
Smokeless Tobacco	0.08	0.05	0.09	0.07	0.08	0.07	0.08	0.08	0.11	0.08
Cigars	0.21	0.19	0.20	0.19	0.19	0.18	0.17	0.21	0.17	0.20
Pipe Tobacco	0.07	0.06	0.10	0.09	0.08	0.09	0.10	0.09	0.09	0.10
ALCOHOL	0.48	0.49	0.48	0.47	0.47	0.49	0.46	0.43	0.45	0.43
Binge Alcohol Use ²	0.38	0.38	0.41	0.35	0.39	0.37	0.35	0.33	0.36	0.30
Heavy Alcohol Use ²	0.16	0.19	0.19	0.16	0.17	0.18	0.16	0.15	0.14	0.12

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2011.

Table 7.30P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: P Values from Tests of Differences of Percentages, 2011 Versus 2002-2010

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL										
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0468	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0004	0.0785	
Smokeless Tobacco	0.5281	0.8661	0.2634	0.9639	0.0866	0.0382	0.3486	0.0917	0.1062	
Cigars	0.0000	0.0000	0.0000	0.0021	0.0024	0.0006	0.0877	0.0068	0.3516	
Pipe Tobacco	0.0591	0.3879	0.8237	0.2858	0.4954	0.9117	0.8682	0.0621	0.3652	
ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0018	0.0007	0.4804	
Binge Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.1690	
Heavy Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0019	0.0002	0.2621	
MALE										
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0432	0.0001	0.2252	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0003	0.0008	0.0787	0.0269	0.4482	
Smokeless Tobacco	0.5932	0.6666	0.1728	0.8367	0.0560	0.0191	0.3161	0.0951	0.2124	
Cigars	0.0000	0.0000	0.0000	0.0004	0.0013	0.0000	0.0067	0.0101	0.8108	
Pipe Tobacco	0.5467	0.3502	0.6606	0.7263	0.5796	0.4892	0.9363	0.0045	1.0000	
ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0678	0.0014	0.4731	
Binge Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0149	0.0002	0.3265	
Heavy Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0001	0.0002	0.0433	0.0453	0.3561	
FEMALE										
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0101	0.1066	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0050	0.0734	
Smokeless Tobacco	0.7300	0.0303	0.6433	0.4582	0.8544	0.6086	0.9925	0.7300	0.2950	
Cigars	0.3359	0.3665	0.1524	0.6433	0.3976	0.8403	0.4163	0.3369	0.2135	
Pipe Tobacco	0.0274	0.0020	0.3995	0.0406	0.0557	0.2592	0.8756	0.4415	0.1523	
ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0062	0.1018	0.7530	
Binge Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	0.0113	0.3409	
Heavy Alcohol Use ²	0.0001	0.0000	0.0000	0.0009	0.0006	0.0030	0.0232	0.0009	0.5800	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2011.

Table 7.31C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Numbers in Thousands, 2002-2011

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL										
TOBACCO PRODUCTS¹	150	151	161	155	160	162	153	169	166	167
Cigarettes	150	151	159	152	156	159	149	162	160	162
Smokeless Tobacco	59	56	60	64	65	63	68	78	81	70
Cigars	83	84	97	92	94	92	96	99	101	100
Pipe Tobacco	24	25	30	37	33	32	34	40	41	48
ALCOHOL	164	160	164	166	168	170	163	174	169	184
Binge Alcohol Use ²	161	155	166	171	169	169	167	182	168	189
Heavy Alcohol Use ²	112	114	120	120	125	114	124	128	127	125
MALE										
TOBACCO PRODUCTS¹	100	106	106	110	109	116	109	121	121	113
Cigarettes	99	109	105	107	108	116	106	121	113	111
Smokeless Tobacco	56	55	58	60	62	61	63	73	75	65
Cigars	73	72	83	79	78	81	80	81	84	89
Pipe Tobacco	22	21	27	33	30	30	29	35	33	39
ALCOHOL	111	100	110	101	108	110	109	117	112	119
Binge Alcohol Use ²	110	105	113	111	115	116	118	130	121	126
Heavy Alcohol Use ²	87	91	88	96	95	94	95	99	101	87
FEMALE										
TOBACCO PRODUCTS¹	101	93	102	101	102	97	101	98	96	105
Cigarettes	102	92	101	101	98	95	100	96	97	102
Smokeless Tobacco	8	11	10	12	13	13	14	18	20	14
Cigars	38	43	44	41	44	43	47	45	49	50
Pipe Tobacco	12	12	12	16	11	14	17	15	23	26
ALCOHOL	102	114	104	112	112	114	107	110	109	123
Binge Alcohol Use ²	99	100	96	104	103	110	102	105	101	116
Heavy Alcohol Use ²	55	59	60	56	63	60	65	61	59	72

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2011.

Table 7.31D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Percentages, 2002-2011

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL										
TOBACCO PRODUCTS¹	0.48	0.48	0.50	0.48	0.49	0.50	0.47	0.50	0.49	0.49
Cigarettes	0.48	0.47	0.49	0.47	0.48	0.49	0.45	0.48	0.47	0.47
Smokeless Tobacco	0.19	0.18	0.19	0.20	0.20	0.19	0.20	0.23	0.24	0.20
Cigars	0.27	0.26	0.30	0.28	0.29	0.28	0.29	0.29	0.30	0.29
Pipe Tobacco	0.08	0.08	0.09	0.11	0.10	0.10	0.10	0.12	0.12	0.14
ALCOHOL	0.53	0.50	0.51	0.51	0.51	0.52	0.49	0.52	0.50	0.54
Binge Alcohol Use ²	0.52	0.49	0.52	0.53	0.52	0.52	0.51	0.54	0.49	0.55
Heavy Alcohol Use ²	0.36	0.36	0.37	0.37	0.38	0.35	0.38	0.38	0.37	0.36
MALE										
TOBACCO PRODUCTS¹	0.64	0.67	0.66	0.67	0.66	0.70	0.66	0.72	0.70	0.66
Cigarettes	0.64	0.69	0.65	0.66	0.66	0.70	0.64	0.71	0.65	0.65
Smokeless Tobacco	0.36	0.34	0.36	0.37	0.38	0.37	0.38	0.43	0.43	0.38
Cigars	0.47	0.45	0.51	0.48	0.48	0.49	0.48	0.48	0.49	0.52
Pipe Tobacco	0.14	0.13	0.17	0.20	0.18	0.18	0.17	0.21	0.19	0.23
ALCOHOL	0.71	0.63	0.68	0.62	0.66	0.67	0.66	0.69	0.65	0.69
Binge Alcohol Use ²	0.71	0.66	0.70	0.68	0.70	0.70	0.71	0.77	0.70	0.73
Heavy Alcohol Use ²	0.56	0.57	0.54	0.59	0.58	0.57	0.57	0.59	0.59	0.51
FEMALE										
TOBACCO PRODUCTS¹	0.65	0.59	0.64	0.63	0.63	0.59	0.62	0.59	0.57	0.61
Cigarettes	0.66	0.58	0.63	0.63	0.60	0.58	0.61	0.58	0.58	0.60
Smokeless Tobacco	0.05	0.07	0.06	0.08	0.08	0.08	0.08	0.11	0.12	0.08
Cigars	0.25	0.27	0.27	0.26	0.27	0.26	0.29	0.27	0.29	0.29
Pipe Tobacco	0.08	0.08	0.08	0.10	0.07	0.08	0.10	0.09	0.14	0.15
ALCOHOL	0.66	0.72	0.65	0.69	0.69	0.70	0.65	0.66	0.65	0.72
Binge Alcohol Use ²	0.64	0.63	0.60	0.65	0.64	0.68	0.62	0.63	0.60	0.68
Heavy Alcohol Use ²	0.35	0.37	0.37	0.35	0.39	0.37	0.39	0.36	0.35	0.42

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2011.

Table 7.31P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: P Values from Tests of Differences of Percentages, 2011 Versus 2002-2010

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL										
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	0.0033	0.0016	0.0253	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.0004	0.1714	
Smokeless Tobacco	0.0306	0.0048	0.0653	0.2618	0.4829	0.4996	0.7399	0.0244	0.0014	
Cigars	0.6895	0.1533	0.0000	0.0047	0.0018	0.0126	0.1363	0.1078	0.2458	
Pipe Tobacco	0.0000	0.0000	0.0002	0.0260	0.0025	0.0001	0.0087	0.6381	0.8545	
ALCOHOL	0.7781	0.3596	0.7986	0.7665	0.0711	0.4408	0.5605	0.1377	0.2848	
Binge Alcohol Use ²	0.1367	0.0168	0.0606	0.0047	0.0013	0.0068	0.0614	0.0090	0.3236	
Heavy Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0019	0.0037	
MALE										
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0001	0.0103	0.1766	0.0177	0.1312	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0001	0.0231	0.2131	0.0304	0.8305	
Smokeless Tobacco	0.0786	0.0069	0.1172	0.2908	0.5479	0.4720	0.8295	0.0499	0.0037	
Cigars	0.2445	0.0471	0.0000	0.0007	0.0001	0.0005	0.0270	0.0267	0.2193	
Pipe Tobacco	0.0006	0.0000	0.0574	0.2638	0.1111	0.0133	0.0448	0.6341	0.5108	
ALCOHOL	0.0471	0.0001	0.0843	0.0009	0.0051	0.0425	0.3349	0.0080	0.0053	
Binge Alcohol Use ²	0.0002	0.0000	0.0002	0.0000	0.0001	0.0006	0.0481	0.0014	0.0547	
Heavy Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	0.0072	
FEMALE										
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0080	0.0032	0.0198	0.1085	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	0.0017	0.0211	
Smokeless Tobacco	0.0041	0.1709	0.0276	0.3309	0.2266	0.6890	0.3315	0.1175	0.3496	
Cigars	0.1957	0.4999	0.9394	0.6402	0.5590	0.1425	0.3537	0.4228	0.8710	
Pipe Tobacco	0.0002	0.0001	0.0000	0.0110	0.0003	0.0007	0.0680	0.0666	0.5759	
ALCOHOL	0.0137	0.0218	0.0311	0.0062	0.9466	0.3689	0.9163	0.6545	0.2070	
Binge Alcohol Use ²	0.0957	0.1169	0.2794	0.1394	0.3515	0.5534	0.4257	0.5322	0.4785	
Heavy Alcohol Use ²	0.1584	0.0566	0.0961	0.0978	0.0002	0.0029	0.0202	0.3578	0.1787	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2011.

Table 7.32C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2011

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL										
TOBACCO PRODUCTS¹	797	748	752	798	799	801	849	784	813	805
Cigarettes	754	738	718	726	760	763	785	753	747	771
Smokeless Tobacco	270	274	236	256	267	268	303	274	302	266
Cigars	331	332	311	335	332	312	352	341	340	352
Pipe Tobacco	160	126	142	141	162	166	141	141	141	135
ALCOHOL	953	887	933	936	914	986	958	931	946	978
Binge Alcohol Use ²	702	639	654	650	697	714	703	751	715	699
Heavy Alcohol Use ²	355	349	356	347	392	402	391	377	398	396
MALE										
TOBACCO PRODUCTS¹	553	546	552	554	566	575	613	574	599	560
Cigarettes	506	521	512	493	527	551	546	538	514	514
Smokeless Tobacco	252	247	226	232	258	238	286	267	290	252
Cigars	311	302	280	309	305	290	325	318	306	321
Pipe Tobacco	132	124	135	136	156	164	128	136	134	129
ALCOHOL	630	577	610	598	583	640	616	631	645	665
Binge Alcohol Use ²	527	492	492	507	541	562	565	585	552	552
Heavy Alcohol Use ²	323	306	307	302	333	355	340	326	337	343
FEMALE										
TOBACCO PRODUCTS¹	488	477	469	515	477	486	543	485	516	520
Cigarettes	475	460	466	508	472	469	539	481	509	510
Smokeless Tobacco	99	125	63	85	74	111	73	49	72	80
Cigars	99	117	107	98	116	104	101	116	140	130
Pipe Tobacco	89	32	39	37	43	26	50	32	47	36
ALCOHOL	628	582	654	615	614	658	656	668	620	663
Binge Alcohol Use ²	393	347	361	370	356	364	382	422	427	406
Heavy Alcohol Use ²	158	155	160	146	147	157	181	181	190	184

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2011.

Table 7.32D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Percentages, 2002-2011

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL										
TOBACCO PRODUCTS ¹	0.44	0.41	0.41	0.43	0.43	0.42	0.44	0.40	0.42	0.41
Cigarettes	0.42	0.41	0.39	0.39	0.40	0.40	0.41	0.39	0.38	0.39
Smokeless Tobacco	0.15	0.15	0.13	0.14	0.14	0.14	0.16	0.14	0.15	0.13
Cigars	0.18	0.18	0.17	0.18	0.18	0.16	0.18	0.18	0.17	0.18
Pipe Tobacco	0.09	0.07	0.08	0.08	0.09	0.09	0.07	0.07	0.07	0.07
ALCOHOL	0.53	0.49	0.51	0.51	0.49	0.52	0.50	0.48	0.48	0.49
Binge Alcohol Use ²	0.39	0.35	0.36	0.35	0.37	0.38	0.37	0.39	0.37	0.35
Heavy Alcohol Use ²	0.20	0.19	0.19	0.19	0.21	0.21	0.20	0.19	0.20	0.20
MALE										
TOBACCO PRODUCTS ¹	0.65	0.63	0.63	0.62	0.63	0.63	0.67	0.62	0.64	0.59
Cigarettes	0.59	0.60	0.59	0.56	0.59	0.61	0.59	0.58	0.55	0.54
Smokeless Tobacco	0.29	0.29	0.26	0.26	0.29	0.26	0.31	0.29	0.31	0.27
Cigars	0.36	0.35	0.32	0.35	0.34	0.32	0.35	0.34	0.33	0.34
Pipe Tobacco	0.15	0.14	0.15	0.15	0.17	0.18	0.14	0.15	0.14	0.14
ALCOHOL	0.74	0.67	0.70	0.67	0.65	0.70	0.67	0.68	0.69	0.70
Binge Alcohol Use ²	0.62	0.57	0.56	0.57	0.60	0.62	0.61	0.63	0.59	0.58
Heavy Alcohol Use ²	0.38	0.35	0.35	0.34	0.37	0.39	0.37	0.35	0.36	0.36
FEMALE										
TOBACCO PRODUCTS ¹	0.52	0.50	0.49	0.53	0.49	0.49	0.54	0.48	0.51	0.50
Cigarettes	0.51	0.49	0.49	0.53	0.48	0.47	0.54	0.48	0.50	0.49
Smokeless Tobacco	0.11	0.13	0.07	0.09	0.08	0.11	0.07	0.05	0.07	0.08
Cigars	0.11	0.12	0.11	0.10	0.12	0.11	0.10	0.12	0.14	0.13
Pipe Tobacco	0.10	0.03	0.04	0.04	0.04	0.03	0.05	0.03	0.05	0.03
ALCOHOL	0.67	0.61	0.68	0.64	0.63	0.66	0.66	0.66	0.61	0.64
Binge Alcohol Use ²	0.42	0.37	0.38	0.38	0.36	0.37	0.38	0.42	0.42	0.39
Heavy Alcohol Use ²	0.17	0.16	0.17	0.15	0.15	0.16	0.18	0.18	0.19	0.18

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2011.

Table 7.32P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: P Values from Tests of Differences of Percentages, 2011 Versus 2002-2010

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL										
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0001	0.0000	0.0000	0.0002	0.0008	0.0713	0.1136	
Cigarettes	0.0000	0.0000	0.0001	0.0000	0.0000	0.0001	0.0006	0.0459	0.0771	
Smokeless Tobacco	0.1682	0.2963	0.2234	0.9603	0.3368	0.7653	0.0700	0.5052	0.3536	
Cigars	0.1150	0.2417	0.0852	0.0472	0.0706	0.3685	0.3289	0.3962	0.3313	
Pipe Tobacco	0.4453	0.8280	0.8234	0.0836	0.0232	0.3472	0.8685	0.9651	0.6725	
ALCOHOL	0.1055	0.0003	0.0032	0.9634	0.0463	0.1761	0.6187	0.7509	0.7679	
Binge Alcohol Use ²	0.7539	0.1977	0.3324	0.2679	0.7057	0.4937	0.2570	0.1059	0.5015	
Heavy Alcohol Use ²	0.4674	0.4566	0.1602	0.8089	0.3333	0.1579	0.0391	0.0728	0.1783	
MALE										
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0001	0.0000	0.0000	0.0002	0.0026	0.3207	0.0817	
Cigarettes	0.0000	0.0000	0.0001	0.0002	0.0000	0.0001	0.0038	0.3568	0.0765	
Smokeless Tobacco	0.1981	0.4837	0.2785	0.9725	0.1789	0.6228	0.0370	0.2632	0.2709	
Cigars	0.0118	0.1840	0.0149	0.0077	0.0699	0.1983	0.1152	0.4003	0.4499	
Pipe Tobacco	0.6794	0.8959	0.8012	0.1037	0.0256	0.2697	0.5975	0.9883	0.7516	
ALCOHOL	0.6520	0.9333	0.8906	0.1747	0.8419	0.5598	0.2932	0.7538	0.8826	
Binge Alcohol Use ²	0.4129	0.9012	0.6089	0.6347	0.3461	0.1003	0.0403	0.1222	0.4088	
Heavy Alcohol Use ²	0.0867	0.4094	0.1791	0.6777	0.0969	0.0604	0.0382	0.1638	0.2695	
FEMALE										
TOBACCO PRODUCTS¹	0.0010	0.0015	0.1034	0.0115	0.0173	0.0841	0.0617	0.1018	0.6700	
Cigarettes	0.0004	0.0021	0.0532	0.0051	0.0075	0.0446	0.0332	0.0420	0.4015	
Smokeless Tobacco	0.5568	0.2967	0.5121	0.9959	0.2275	0.6092	0.5016	0.0502	0.4194	
Cigars	0.0305	0.9437	0.1706	0.1102	0.8023	0.3692	0.1063	0.9606	0.5551	
Pipe Tobacco	0.3818	0.7461	0.9473	0.5234	0.6693	0.4525	0.3308	0.7430	0.7541	
ALCOHOL	0.0045	0.0000	0.0000	0.1567	0.0056	0.1393	0.0739	0.4139	0.5088	
Binge Alcohol Use ²	0.1113	0.0151	0.0160	0.1676	0.0339	0.2206	0.3086	0.5382	0.9121	
Heavy Alcohol Use ²	0.0926	0.9328	0.5955	0.1720	0.2783	0.5761	0.6182	0.2703	0.4666	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2011.

Table 7.33C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2011

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL										
TOBACCO PRODUCTS¹	829	784	793	839	838	838	872	822	840	845
Cigarettes	790	774	765	772	799	803	809	790	772	814
Smokeless Tobacco	276	283	243	264	281	280	311	288	310	275
Cigars	346	333	323	351	348	331	363	358	358	369
Pipe Tobacco	163	130	145	145	165	169	146	149	147	143
ALCOHOL	972	909	950	969	954	1,011	977	945	974	1,006
Binge Alcohol Use ²	730	675	699	701	742	766	737	784	748	736
Heavy Alcohol Use ²	389	376	383	378	425	426	414	404	421	423
MALE										
TOBACCO PRODUCTS¹	565	561	568	576	581	594	630	597	616	580
Cigarettes	521	537	532	514	542	573	565	564	529	535
Smokeless Tobacco	260	255	233	239	271	249	295	281	299	261
Cigars	322	306	291	324	316	306	332	332	318	337
Pipe Tobacco	136	127	138	141	159	168	131	143	136	135
ALCOHOL	643	590	616	616	599	658	626	644	666	678
Binge Alcohol Use ²	540	517	511	529	561	585	588	606	573	568
Heavy Alcohol Use ²	346	320	318	322	352	373	358	349	352	355
FEMALE										
TOBACCO PRODUCTS¹	509	498	494	541	506	505	553	501	532	539
Cigarettes	497	481	488	537	497	487	548	494	524	529
Smokeless Tobacco	99	126	64	87	75	112	74	51	76	82
Cigars	110	121	119	109	128	112	113	127	150	140
Pipe Tobacco	90	35	41	40	45	28	54	35	53	45
ALCOHOL	640	593	662	635	641	673	667	679	634	677
Binge Alcohol Use ²	411	375	389	401	395	392	399	442	444	443
Heavy Alcohol Use ²	175	170	173	162	168	169	192	192	201	207

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2011.

Table 7.33D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Percentages, 2002-2011

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL										
TOBACCO PRODUCTS¹	0.39	0.37	0.37	0.39	0.38	0.38	0.39	0.36	0.37	0.36
Cigarettes	0.38	0.36	0.36	0.35	0.36	0.36	0.36	0.35	0.34	0.35
Smokeless Tobacco	0.13	0.13	0.11	0.12	0.13	0.13	0.14	0.13	0.14	0.12
Cigars	0.16	0.16	0.15	0.16	0.16	0.15	0.16	0.16	0.16	0.16
Pipe Tobacco	0.08	0.06	0.07	0.07	0.07	0.08	0.06	0.07	0.06	0.06
ALCOHOL	0.46	0.43	0.44	0.44	0.43	0.45	0.43	0.42	0.42	0.43
Binge Alcohol Use ²	0.35	0.32	0.32	0.32	0.34	0.34	0.33	0.34	0.33	0.32
Heavy Alcohol Use ²	0.19	0.18	0.18	0.17	0.19	0.19	0.18	0.18	0.18	0.18
MALE										
TOBACCO PRODUCTS¹	0.56	0.55	0.55	0.55	0.55	0.55	0.58	0.54	0.56	0.52
Cigarettes	0.52	0.53	0.51	0.49	0.51	0.53	0.52	0.51	0.48	0.48
Smokeless Tobacco	0.26	0.25	0.22	0.23	0.25	0.23	0.27	0.26	0.27	0.23
Cigars	0.32	0.30	0.28	0.31	0.30	0.29	0.31	0.30	0.29	0.30
Pipe Tobacco	0.13	0.12	0.13	0.13	0.15	0.16	0.12	0.13	0.12	0.12
ALCOHOL	0.64	0.58	0.59	0.59	0.56	0.61	0.58	0.59	0.60	0.61
Binge Alcohol Use ²	0.53	0.51	0.49	0.50	0.53	0.55	0.54	0.55	0.52	0.51
Heavy Alcohol Use ²	0.34	0.31	0.31	0.31	0.33	0.35	0.33	0.32	0.32	0.32
FEMALE										
TOBACCO PRODUCTS¹	0.47	0.45	0.44	0.48	0.44	0.44	0.47	0.43	0.45	0.45
Cigarettes	0.45	0.44	0.44	0.48	0.44	0.42	0.47	0.42	0.44	0.44
Smokeless Tobacco	0.09	0.11	0.06	0.08	0.07	0.10	0.06	0.04	0.06	0.07
Cigars	0.10	0.11	0.11	0.10	0.11	0.10	0.10	0.11	0.13	0.12
Pipe Tobacco	0.08	0.03	0.04	0.04	0.04	0.02	0.05	0.03	0.04	0.04
ALCOHOL	0.59	0.54	0.59	0.56	0.56	0.58	0.57	0.58	0.54	0.56
Binge Alcohol Use ²	0.38	0.34	0.35	0.36	0.35	0.34	0.34	0.38	0.37	0.37
Heavy Alcohol Use ²	0.16	0.15	0.16	0.14	0.15	0.15	0.16	0.16	0.17	0.17

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2011.

Table 7.33P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: P Values from Tests of Differences of Percentages, 2011 Versus 2002-2010

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL										
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0171	0.0482	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0085	0.0466	
Smokeless Tobacco	0.4016	0.6919	0.1070	0.8405	0.4507	0.9050	0.0943	0.2207	0.0817	
Cigars	0.1077	0.1168	0.0027	0.0070	0.0093	0.1248	0.1956	0.2052	0.2026	
Pipe Tobacco	0.6700	0.0746	0.4393	0.3061	0.1668	0.9714	0.3548	0.8572	0.7186	
ALCOHOL	0.1033	0.0008	0.0037	0.9822	0.1045	0.2364	0.6912	0.9788	0.9236	
Binge Alcohol Use ²	0.9561	0.5730	0.7177	0.7436	0.6364	0.2003	0.1367	0.0295	0.3614	
Heavy Alcohol Use ²	0.0250	0.0136	0.0021	0.0939	0.0046	0.0049	0.0008	0.0082	0.0322	
MALE										
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0016	0.1477	0.0452	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0025	0.1797	0.0892	
Smokeless Tobacco	0.4015	0.9429	0.1588	0.8371	0.2536	0.7709	0.0502	0.1159	0.0678	
Cigars	0.0069	0.0672	0.0002	0.0006	0.0060	0.0300	0.0421	0.1647	0.2774	
Pipe Tobacco	0.6997	0.2397	0.8367	0.1984	0.0729	0.5917	0.2973	0.8829	0.8863	
ALCOHOL	0.4304	0.4438	0.8591	0.0562	0.7562	0.8317	0.2291	0.4325	0.5432	
Binge Alcohol Use ²	0.1105	0.2150	0.1679	0.4733	0.0733	0.0222	0.0175	0.0291	0.2197	
Heavy Alcohol Use ²	0.0014	0.0125	0.0024	0.0194	0.0015	0.0029	0.0014	0.0249	0.0701	
FEMALE										
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0032	0.0002	0.0004	0.0311	0.0179	0.0407	0.4806	
Cigarettes	0.0000	0.0000	0.0005	0.0000	0.0001	0.0080	0.0052	0.0097	0.2206	
Smokeless Tobacco	0.8080	0.3799	0.3179	0.8795	0.1603	0.5735	0.4017	0.1262	0.5685	
Cigars	0.0145	0.7861	0.2366	0.1137	0.9697	0.1590	0.0638	0.8091	0.6189	
Pipe Tobacco	0.8712	0.0244	0.0482	0.4802	0.2177	0.0113	0.9761	0.2155	0.5799	
ALCOHOL	0.0014	0.0000	0.0000	0.0649	0.0070	0.1049	0.0730	0.3830	0.3870	
Binge Alcohol Use ²	0.0554	0.0094	0.0150	0.1100	0.0979	0.3153	0.4244	0.4592	0.7824	
Heavy Alcohol Use ²	0.2970	0.4402	0.2720	0.5341	0.6958	0.5946	0.2292	0.1830	0.2748	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2011.