

Table 2.11A Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	173,654	175,260	82,829	80,846	69,497 ^a	66,879
12	159	130	80	78	27	26
13	353	324	213	184	101	81
14	661	643	433	391	199	189
15	1,015	934	707	676	331	355
16	1,416 ^a	1,275	1,051	942	619	538
17	1,750	1,617	1,316	1,189	877 ^a	762
18	2,550	2,297	2,077	1,874	1,501	1,333
19	2,480	2,450	1,983	1,941	1,457	1,468
20	2,835	2,731	2,188	2,137	1,675	1,678
21	3,007	2,897	2,232	2,145	1,791	1,654
22	3,105	3,071	2,303	2,138	1,834	1,681
23	3,046	3,092	2,083	2,188	1,666	1,801
24	2,973	3,154	2,052	2,205	1,677	1,700
25	2,929	2,957	1,973	1,969	1,564	1,566
26-29	12,548	12,541	8,094	8,156	6,741	6,502
30-34	14,583	15,385	8,385	9,205	7,082	7,814
35-39	14,135	13,465	7,394	6,768	6,297	5,701
40-44	14,617	15,693	7,159	7,262	6,219	6,258
45-49	15,557	14,977	7,475	6,869	6,727	6,029
50-54	17,262	17,810	7,392	7,118	6,603	6,141
55-59	15,231	14,664	6,128	5,475	5,560	4,718
60-64	11,861	13,072	3,853	4,034	3,452	3,490
65 or Older	29,580	30,080	6,257	5,899	5,497	5,392

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.11B Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	66.8	66.8	31.9 ^a	30.8	26.7 ^b	25.5
12	3.9	3.5	2.0	2.1	0.7	0.7
13	8.8	7.7	5.3	4.4	2.5	1.9
14	15.7	14.7	10.3	8.9	4.7	4.3
15	24.2	22.6	16.9	16.4	7.9	8.6
16	33.7 ^b	29.8	25.0 ^a	22.0	14.7 ^a	12.6
17	40.9	39.2	30.8	28.8	20.5	18.5
18	54.2 ^b	48.5	44.1 ^b	39.6	31.9 ^a	28.2
19	58.3	58.8	46.6	46.6	34.2	35.2
20	65.8	63.6	50.8	49.8	38.9	39.1
21	68.1	68.4	50.5	50.6	40.5	39.0
22	69.8	68.6	51.8 ^a	47.8	41.2 ^a	37.6
23	69.7	70.6	47.6	50.0	38.1	41.1
24	72.8	71.8	50.2	50.2	41.1	38.7
25	73.4	71.9	49.4	47.9	39.2	38.1
26-29	73.1	72.9	47.2	47.4	39.3	37.8
30-34	73.8	76.4	42.4 ^a	45.7	35.9	38.8
35-39	72.7	70.7	38.0	35.5	32.4	29.9
40-44	72.4	73.1	35.5	33.8	30.8	29.1
45-49	72.6	74.9	34.9	34.4	31.4	30.2
50-54	73.8	77.2	31.6	30.8	28.2	26.6
55-59	74.0	71.4	29.8	26.6	27.0 ^a	23.0
60-64	70.8	73.5	23.0	22.7	20.6	19.6
65 or Older	70.7	69.2	14.9	13.6	13.1	12.4

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.12A Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	161,057	162,138	67,802	66,256	57,525	55,778
12	125 ^a	80	66	54	21	16
13	258	237	156	130	72	55
14	506	491	317	274	139	115
15	833	733	542	500	250	252
16	1,159 ^a	1,020	806 ^a	681	465 ^a	372
17	1,457	1,342	1,062 ^a	922	688 ^a	587
18	2,224 ^b	1,868	1,697 ^b	1,409	1,181 ^a	992
19	2,142	2,118	1,588	1,594	1,181	1,188
20	2,512	2,434	1,846	1,789	1,383	1,391
21	2,685	2,547	1,914	1,772	1,478	1,336
22	2,809	2,789	1,977	1,844	1,559	1,468
23	2,751	2,799	1,788	1,829	1,442	1,489
24	2,717	2,868	1,738	1,887	1,445	1,463
25	2,753	2,719	1,647	1,626	1,340	1,329
26-29	11,802	11,707	6,868	6,897	5,725	5,623
30-34	13,666	14,453	7,242	7,863	6,308	6,689
35-39	13,206	12,467	6,019	5,720	5,187	4,876
40-44	13,545	14,461	5,481	5,795	4,899	5,018
45-49	14,264	13,874	6,035	5,442	5,577 ^a	4,882
50-54	16,068	16,726	6,180	5,720	5,742	5,323
55-59	14,368	13,975	4,818	4,456	4,433	4,036
60-64	11,174	12,237	3,173	3,234	2,828	2,915
65 or Older	28,031	28,193	4,845	4,817	4,183	4,363

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.12B Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	61.9	61.8	26.1	25.3	22.1 ^a	21.3
12	3.1 ^a	2.1	1.6	1.4	0.5	0.4
13	6.4	5.6	3.9	3.1	1.8	1.3
14	12.0	11.2	7.5	6.2	3.3	2.6
15	19.9	17.7	12.9	12.1	6.0	6.1
16	27.6 ^b	23.9	19.2 ^b	15.9	11.1 ^b	8.7
17	34.1	32.5	24.8 ^a	22.3	16.1	14.2
18	47.3 ^b	39.5	36.1 ^b	29.8	25.1 ^b	21.0
19	50.3	50.8	37.3	38.3	27.7	28.5
20	58.3	56.7	42.8	41.7	32.1	32.4
21	60.8	60.1	43.3	41.8	33.4	31.5
22	63.2	62.3	44.4	41.2	35.1	32.8
23	62.9	63.9	40.9	41.8	33.0	34.0
24	66.5	65.3	42.5	43.0	35.4	33.3
25	68.9	66.1	41.2	39.5	33.6	32.3
26-29	68.8	68.0	40.0	40.1	33.4	32.7
30-34	69.2	71.8	36.7	39.0	31.9	33.2
35-39	67.9	65.5	31.0	30.0	26.7	25.6
40-44	67.1	67.3	27.2	27.0	24.3	23.4
45-49	66.6	69.4	28.2	27.2	26.0	24.4
50-54	68.7	72.5	26.4	24.8	24.5	23.1
55-59	69.8	68.0	23.4	21.7	21.5	19.6
60-64	66.7	68.8	19.0	18.2	16.9	16.4
65 or Older	67.0	64.9	11.6	11.1	10.0	10.0

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.13A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	45,936	46,139	12,040	11,866	9,036	8,813
12	43	45	17	30	9	10
13	111	91	63	45	27	15
14	187	187	117	126	57	63
15	279	269	163	193	86	91
16	444	416	286	292	154	138
17	522	493	325	325	182	171
18	717	726	441	489	247	306
19	779	834	445	476	236	281
20	818	834	368	428	228	262
21	927	835	403	428	255	266
22	901	1,047	414	394	275	237
23	968	967	370	353	229	234
24	872	999	327	335	223	232
25	909	823	341	328	219	201
26-29	3,870	3,623	1,257	1,131	936	774
30-34	4,287	4,364	1,034	1,243	786	962
35-39	4,873	4,294	1,171	982	926	761
40-44	5,148	5,314	1,381	1,114	1,204	1,037
45-49	4,578	4,806	928	1,021	823	902
50-54	4,017 ^a	4,930	600	777	524	603
55-59	3,447	2,858	490	535	434	500
60-64	2,234	2,521	393	219	354	179
65 or Older	5,005	4,865	706	603	623	590

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.13B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	17.7	17.6	4.6	4.5	3.5	3.4
12	1.1	1.2	0.4	0.8	0.2	0.3
13	2.8	2.2	1.6	1.1	0.7	0.3
14	4.4	4.3	2.8	2.9	1.4	1.4
15	6.7	6.5	3.9	4.7	2.1	2.2
16	10.6	9.7	6.8	6.8	3.7	3.2
17	12.2	11.9	7.6	7.9	4.3	4.1
18	15.2	15.3	9.4	10.3	5.3	6.5
19	18.3	20.0	10.5	11.4	5.5	6.8
20	19.0	19.4	8.5	10.0	5.3	6.1
21	21.0	19.7	9.1	10.1	5.8	6.3
22	20.3 ^a	23.4	9.3	8.8	6.2	5.3
23	22.1	22.1	8.5	8.1	5.2	5.3
24	21.4	22.7	8.0	7.6	5.4	5.3
25	22.8	20.0	8.5	8.0	5.5	4.9
26-29	22.6	21.1	7.3	6.6	5.5	4.5
30-34	21.7	21.7	5.2	6.2	4.0	4.8
35-39	25.1	22.5	6.0	5.2	4.8	4.0
40-44	25.5	24.7	6.8 ^a	5.2	6.0	4.8
45-49	21.4 ^a	24.0	4.3	5.1	3.8	4.5
50-54	17.2 ^a	21.4	2.6	3.4	2.2	2.6
55-59	16.7	13.9	2.4	2.6	2.1	2.4
60-64	13.3	14.2	2.3	1.2	2.1 ^a	1.0
65 or Older	12.0	11.2	1.7	1.4	1.5	1.4

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.14A Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	88,596	89,117	25,647	24,933	13,417	12,438
12	26	33	15	18	4	5
13	110	97	67	65	27	13
14	235	202	178 ^b	118	62 ^b	29
15	410 ^a	338	285	242	80	93
16	710 ^a	586	499 ^a	415	212	184
17	899	813	640	566	261	252
18	1,455 ^a	1,258	1,111	979	574	495
19	1,418	1,404	1,009	996	488	455
20	1,684	1,611	1,068	1,024	520	513
21	1,805	1,754	973	960	498	477
22	1,841	1,873	982	869	492 ^a	386
23	1,802	1,899	811	876	399	463
24	1,782	1,876	795	820	391	384
25	1,702	1,704	771	690	353	300
26-29	7,498	7,724	2,844	3,116	1,485	1,465
30-34	8,371	8,916	2,553	2,835	1,223	1,430
35-39	7,819	7,255	2,086	1,713	1,134 ^a	812
40-44	7,456	8,012	1,787	1,789	911	972
45-49	7,567	7,302	1,766	1,714	946	927
50-54	8,080	9,140	1,780	1,926	1,006	939
55-59	7,650	6,789	1,733	1,232	1,194 ^b	582
60-64	5,889	6,342	849	1,061	467	640
65 or Older	12,387	12,187	1,045	909	690	621

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.14B Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	34.1	34.0	9.9	9.5	5.2 ^a	4.7
12	0.6	0.9	0.4	0.5	0.1	0.1
13	2.7	2.3	1.7	1.5	0.7	0.3
14	5.6	4.6	4.2 ^b	2.7	1.5 ^b	0.7
15	9.8	8.2	6.8	5.8	1.9	2.3
16	16.9 ^b	13.7	11.9 ^a	9.7	5.0	4.3
17	21.0	19.7	14.9	13.7	6.1	6.1
18	30.9 ^b	26.6	23.6 ^a	20.7	12.2	10.5
19	33.3	33.7	23.7	23.9	11.5	10.9
20	39.1	37.5	24.8	23.8	12.1	11.9
21	40.8	41.4	22.0	22.6	11.3	11.3
22	41.4	41.9	22.1	19.4	11.1 ^a	8.6
23	41.2	43.4	18.6	20.0	9.1	10.6
24	43.6	42.7	19.5	18.7	9.6	8.7
25	42.6	41.5	19.3	16.8	8.8	7.3
26-29	43.7	44.9	16.6	18.1	8.7	8.5
30-34	42.4	44.3	12.9	14.1	6.2	7.1
35-39	40.2	38.1	10.7	9.0	5.8 ^a	4.3
40-44	37.0	37.3	8.9	8.3	4.5	4.5
45-49	35.3	36.5	8.2	8.6	4.4	4.6
50-54	34.5 ^a	39.6	7.6	8.3	4.3	4.1
55-59	37.2	33.0	8.4	6.0	5.8 ^b	2.8
60-64	35.2	35.7	5.1	6.0	2.8	3.6
65 or Older	29.6	28.0	2.5	2.1	1.6	1.4

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.15A Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	214,013	213,794	173,573	173,966	135,502	136,868
12	268	227	175	129	48	41
13	527	503	371	341	132	126
14	1,060	977	826	717	366	306
15	1,598 ^a	1,450	1,290	1,179	570	501
16	2,072	2,061	1,735	1,678	870	763
17	2,543	2,451	2,158	2,089	1,232	1,140
18	3,207	3,097	2,882	2,751	1,879	1,661
19	3,148	3,096	2,822	2,762	1,902	1,894
20	3,494	3,388	3,113	3,046	2,296	2,219
21	3,904	3,746	3,729	3,597	3,079	2,941
22	4,042	4,025	3,820	3,757	3,095	3,132
23	3,975	4,021	3,642	3,694	2,998	3,052
24	3,757	4,011	3,413	3,674	2,827 ^a	3,095
25	3,667	3,763	3,356	3,427	2,752	2,745
26-29	15,734	15,662	14,199	14,205	11,496	11,878
30-34	17,729	18,041	15,523	15,860	12,295	12,808
35-39	17,325	16,897	14,942	14,258	12,119	11,473
40-44	18,153	19,313	15,494	16,345	12,131	13,043
45-49	19,556 ^a	17,858	15,854	14,693	12,643	11,784
50-54	21,107	21,234	17,154	17,259	13,701	13,827
55-59	18,281	18,135	13,934	13,481	10,941	10,792
60-64	14,890	15,390	11,212	11,355	8,888	9,524
65 or Older	33,976	34,448	21,927 ^a	23,671	17,242	18,123

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.15B Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	82.3 ^a	81.5	66.7	66.3	52.1	52.2
12	6.6	6.1	4.3	3.4	1.2	1.1
13	13.1	11.9	9.2	8.1	3.3	3.0
14	25.2 ^a	22.3	19.7 ^b	16.3	8.7	7.0
15	38.1 ^a	35.1	30.8	28.5	13.6	12.1
16	49.3	48.2	41.3	39.2	20.7 ^a	17.9
17	59.4	59.4	50.4	50.6	28.8	27.6
18	68.2	65.4	61.3	58.1	39.9 ^b	35.1
19	73.9	74.3	66.3	66.3	44.7	45.5
20	81.1	78.9	72.2	70.9	53.3	51.7
21	88.3	88.4	84.4	84.9	69.7	69.4
22	90.9	89.9	85.9	83.9	69.6	70.0
23	90.9	91.9	83.3	84.4	68.6	69.7
24	92.0	91.3	83.6	83.6	69.2	70.5
25	91.8	91.6	84.0	83.4	68.9	66.8
26-29	91.7	91.0	82.8	82.5	67.0	69.0
30-34	89.8	89.6	78.6	78.8	62.2	63.6
35-39	89.1	88.7	76.9	74.9	62.3	60.2
40-44	90.0	89.9	76.8	76.1	60.1	60.7
45-49	91.3	89.3	74.0	73.5	59.0	58.9
50-54	90.2	92.0	73.3	74.8	58.5	59.9
55-59	88.8	88.3	67.7	65.6	53.2	52.5
60-64	88.9	86.6	67.0	63.9	53.1	53.6
65 or Older	81.2	79.3	52.4	54.5	41.2	41.7

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.16A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Numbers in Thousands, 2012 and 2013

Age Category	Alcohol Use (2012)	Alcohol Use (2013)	Binge Alcohol Use (2012)	Binge Alcohol Use (2013)	Heavy Alcohol Use (2012)	Heavy Alcohol Use (2013)
TOTAL	135,502	136,868	59,685	60,064	17,021	16,541
12	48	41	20	12	1	4
13	132	126	53	55	15	5
14	366	306	169	136	13	15
15	570	501	285	249	37	43
16	870	763	495	419	101	73
17	1,232	1,140	778	685	159	152
18	1,879	1,661	1,225	1,053	310	284
19	1,902	1,894	1,306	1,312	472 ^a	356
20	2,296	2,219	1,521	1,478	543	480
21	3,079	2,941	2,093	1,914	716 ^a	585
22	3,095	3,132	2,061	2,042	659	625
23	2,998	3,052	1,930	1,895	604	591
24	2,827 ^a	3,095	1,809	1,883	514	577
25	2,752	2,745	1,716	1,609	567 ^a	440
26-29	11,496	11,878	6,463	6,890	1,704	1,934
30-34	12,295	12,808	6,662	7,092	1,761	2,124
35-39	12,119	11,473	5,623	5,636	1,586	1,425
40-44	12,131	13,043	5,536	5,512	1,616	1,567
45-49	12,643	11,784	5,364	5,151	1,531	1,387
50-54	13,701	13,827	5,320	5,314	1,537	1,294
55-59	10,941	10,792	3,445	3,275	1,033	810
60-64	8,888	9,524	2,397	2,513	721	844
65 or Older	17,242	18,123	3,413	3,938	820	925

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.16B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Percentages, 2012 and 2013

Age Category	Alcohol Use (2012)	Alcohol Use (2013)	Binge Alcohol Use (2012)	Binge Alcohol Use (2013)	Heavy Alcohol Use (2012)	Heavy Alcohol Use (2013)
TOTAL	52.1	52.2	23.0	22.9	6.5	6.3
12	1.2	1.1	0.5	0.3	0.0	0.1
13	3.3	3.0	1.3	1.3	0.4	0.1
14	8.7	7.0	4.0	3.1	0.3	0.3
15	13.6	12.1	6.8	6.0	0.9	1.0
16	20.7 ^a	17.9	11.8 ^a	9.8	2.4	1.7
17	28.8	27.6	18.2	16.6	3.7	3.7
18	39.9 ^b	35.1	26.0 ^a	22.3	6.6	6.0
19	44.7	45.5	30.7	31.5	11.1 ^a	8.5
20	53.3	51.7	35.3	34.4	12.6	11.2
21	69.7	69.4	47.4	45.2	16.2	13.8
22	69.6	70.0	46.3	45.6	14.8	14.0
23	68.6	69.7	44.1	43.3	13.8	13.5
24	69.2	70.5	44.3	42.9	12.6	13.1
25	68.9	66.8	43.0 ^a	39.2	14.2 ^a	10.7
26-29	67.0	69.0	37.7	40.0	9.9	11.2
30-34	62.2	63.6	33.7	35.2	8.9	10.5
35-39	62.3	60.2	28.9	29.6	8.2	7.5
40-44	60.1	60.7	27.4	25.7	8.0	7.3
45-49	59.0	58.9	25.0	25.8	7.1	6.9
50-54	58.5	59.9	22.7	23.0	6.6	5.6
55-59	53.2	52.5	16.7	15.9	5.0	3.9
60-64	53.1	53.6	14.3	14.1	4.3	4.7
65 or Older	41.2	41.7	8.2	9.1	2.0	2.1

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.