

Table 6.11C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	233	232	270	267	257	270	217	245
Marijuana and Hashish	209	209	239	244	224	239	172	207
Cocaine	88	96	84	58	57	58	48	45
Crack	63	35	58	26	9	38	38	27
Heroin	22	13	19	21	22	22	34	26
Hallucinogens	58	57	63	57	27	101	21	35
LSD	26	21	14	29	5	5	6	8
PCP	11	1	4	11	3	*	*	10
Ecstasy	45	45	50	42	25	66	12	13
Inhalants	22	31	53	25	22	36	19	34
Nonmedical Use of Psychotherapeutics ^{3,4}	110	124	135	129	130	119	133	143
Pain Relievers	92	108	105	104	97	90	115	117
OxyContin®	29	28	16	38	13	18	27	64
Tranquilizers	67	59	89	69	84	45	81	62
Stimulants ⁴	45	60	62	65	54	56	39	49
Methamphetamine ⁴	25	42	53	40	28	43	33	44
Sedatives	12	20	36	15	24	44	21	44
ILLICIT DRUGS OTHER THAN MARIJUANA²	142	147	164	145	142	165	142	154

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.11D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	1.12	1.17	0.57	0.57	0.33	0.34	0.17	0.19
Marijuana and Hashish	1.05	1.09	0.52	0.53	0.29	0.30	0.14	0.16
Cocaine	0.51	0.56	0.19	0.13	0.08	0.08	0.04	0.04
Crack	0.37	0.21	0.14	0.06	0.01	0.05	0.03	0.02
Heroin	0.13	0.08	0.05	0.05	0.03	0.03	0.03	0.02
Hallucinogens	0.34	0.34	0.15	0.13	0.04	0.13	0.02	0.03
LSD	0.15	0.13	0.03	0.07	0.01	0.01	0.00	0.01
PCP	0.07	0.01	0.01	0.03	0.00	*	*	0.01
Ecstasy	0.27	0.27	0.12	0.10	0.03	0.09	0.01	0.01
Inhalants	0.13	0.19	0.12	0.06	0.03	0.05	0.02	0.03
Nonmedical Use of Psychotherapeutics ^{3,4}	0.61	0.71	0.31	0.29	0.17	0.16	0.11	0.11
Pain Relievers	0.52	0.63	0.24	0.24	0.13	0.12	0.09	0.09
OxyContin®	0.17	0.17	0.04	0.09	0.02	0.02	0.02	0.05
Tranquilizers	0.39	0.36	0.21	0.16	0.11	0.06	0.07	0.05
Stimulants ⁴	0.26	0.36	0.14	0.15	0.07	0.07	0.03	0.04
Methamphetamine ⁴	0.15	0.25	0.12	0.09	0.04	0.06	0.03	0.03
Sedatives	0.07	0.12	0.08	0.03	0.03	0.06	0.02	0.04
ILLICIT DRUGS OTHER THAN MARIJUANA²	0.77	0.83	0.37	0.33	0.19	0.21	0.11	0.12

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.11P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	0.0968	0.9645	0.7327	0.8904				
Marijuana and Hashish	0.1240	0.7166	0.4166	0.6708				
Cocaine	0.5205	0.1211	0.5355	0.2959				
Crack	0.4542	0.3317	0.0208	0.3843				
Heroin	0.2487	0.7675	0.3897	0.3810				
Hallucinogens	0.9437	0.7452	0.1658	0.3915				
LSD	0.6890	0.4569	0.9520	0.4194				
PCP	0.2314	0.5986	*	*				
Ecstasy	0.8638	0.9450	0.7739	0.9338				
Inhalants	0.6366	0.1346	0.4661	0.3099				
Nonmedical Use of Psychotherapeutics ^{3,4}	0.9769	0.5120	0.1570	0.9796				
Pain Relievers	0.9398	0.2200	0.4121	0.6851				
OxyContin®	0.8382	0.2735	0.4316	0.2585				
Tranquilizers	0.6858	0.3547	0.0319	0.3696				
Stimulants ⁴	0.3023	0.9116	0.9027	0.1287				
Methamphetamine ⁴	0.0819	0.6959	0.7279	0.1641				
Sedatives	0.8107	0.0329	0.7152	0.4196				
ILLICIT DRUGS OTHER THAN MARIJUANA²	0.9723	0.3336	0.8073	0.9906				

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.12C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	21	17	32	35	28	24	53	47
Marijuana and Hashish	19	17	30	33	26	24	40	37
Cocaine	3	6	6	3	2	1	3	6
Crack	0	0	1	0	*	1	*	*
Heroin	*	*	*	3	*	1	*	3
Hallucinogens	9	7	10	10	5	5	11	10
LSD	3	6	3	7	3	1	3	5
PCP	*	1	1	3	2	1	2	1
Ecstasy	6	4	7	6	4	1	6	5
Inhalants	9	4	8	6	7	3	14	13
Nonmedical Use of Psychotherapeutics ^{3,4}	10	8	17	14	12	11	31	27
Pain Relievers	8	7	15	13	11	8	26	24
OxyContin®	3	3	8	4	3	1	3	3
Tranquilizers	7	3	8	8	4	6	16	8
Stimulants ⁴	6	3	9	6	4	5	8	8
Methamphetamine ⁴	2	2	4	3	2	1	4	5
Sedatives	2	2	4	2	0	1	5	6
ILLICIT DRUGS OTHER THAN MARIJUANA²	14	11	20	16	14	12	35	31

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.12D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	3.64	4.26	1.72	2.08	1.60	1.61	0.23	0.20
Marijuana and Hashish	3.91	4.27	1.64	2.04	1.55	1.58	0.18	0.17
Cocaine	0.77	2.02	0.37	0.21	0.13	0.11	0.01	0.03
Crack	0.05	0.03	0.04	0.03	*	0.09	*	*
Heroin	*	*	*	0.27	*	0.09	*	0.01
Hallucinogens	2.65	2.39	0.63	0.80	0.32	0.38	0.05	0.05
LSD	0.95	1.90	0.19	0.51	0.20	0.08	0.01	0.02
PCP	*	0.21	0.04	0.23	0.16	0.05	0.01	0.01
Ecstasy	1.74	1.43	0.46	0.48	0.28	0.10	0.03	0.02
Inhalants	2.46	1.24	0.51	0.44	0.49	0.20	0.06	0.06
Nonmedical Use of Psychotherapeutics ^{3,4}	2.75	2.62	1.10	1.10	0.79	0.80	0.14	0.12
Pain Relievers	2.33	2.36	0.99	0.99	0.73	0.61	0.12	0.11
OxyContin®	0.83	0.86	0.56	0.36	0.24	0.11	0.02	0.02
Tranquilizers	2.20	1.03	0.52	0.64	0.25	0.46	0.07	0.03
Stimulants ⁴	1.78	1.15	0.62	0.44	0.30	0.36	0.04	0.04
Methamphetamine ⁴	0.68	0.80	0.28	0.23	0.14	0.07	0.02	0.02
Sedatives	0.47	0.65	0.25	0.16	0.02	0.08	0.02	0.03
ILLICIT DRUGS OTHER THAN MARIJUANA²	3.37	3.28	1.22	1.25	0.91	0.87	0.16	0.14

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.12P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	0.3020	0.0840	0.2848		0.3343			
Marijuana and Hashish	0.6883	0.0337	0.7205		0.6271			
Cocaine	0.0331	0.2741	0.8772		0.1880			
Crack	0.7937	0.7242	*		*		*	
Heroin	*	*	*		*		*	
Hallucinogens	0.2372	0.2777	0.6923		0.9380			
LSD	0.1719	0.0178	0.2381		0.3061			
PCP	*	0.2382	0.5061		0.9853			
Ecstasy	0.2139	0.8472	0.0984		0.7928			
Inhalants	0.0943	0.2610	0.0079		0.3577			
Nonmedical Use of Psychotherapeutics ^{3,4}	0.1517	0.1152	0.6097		0.1198			
Pain Relievers	0.7515	0.0790	0.2687		0.0685			
OxyContin®	0.8851	0.1764	0.6502		0.7403			
Tranquilizers	0.0300	0.9419	0.6612		0.1224			
Stimulants ⁴	0.1946	0.0460	0.8244		0.9117			
Methamphetamine ⁴	0.9228	0.2434	0.3797		0.7432			
Sedatives	0.9039	0.4627	0.2448		0.9111			
ILLICIT DRUGS OTHER THAN MARIJUANA²	0.1576	0.3377	0.2105		0.1710			

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.13C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	84	82	94	103	69	64	65	61
Marijuana and Hashish	81	77	90	99	66	60	58	51
Cocaine	24	27	19	18	7	7	11	6
Crack	7	10	4	4	2	1	4	5
Heroin	7	6	11	5	6	8	17	12
Hallucinogens	33	30	25	27	11	11	13	19
LSD	8	10	13	13	4	5	5	5
PCP	3	1	2	*	2	*	*	2
Ecstasy	31	22	18	21	9	4	8	7
Inhalants	14	9	12	10	5	4	11	12
Nonmedical Use of Psychotherapeutics ^{3,4}	46	42	39	40	26	27	34	28
Pain Relievers	36	35	33	33	23	23	33	24
OxyContin®	10	13	10	12	11	6	16	4
Tranquilizers	24	21	22	22	13	11	23	13
Stimulants ⁴	22	27	19	19	6	13	9	11
Methamphetamine ⁴	10	15	10	7	2	6	5	4
Sedatives	9	6	7	5	3	2	4	4
ILLICIT DRUGS OTHER THAN MARIJUANA²	55	50	47	48	28	30	38	35

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.13D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	1.28	1.35	0.77	0.86	0.84	0.79	0.44	0.41
Marijuana and Hashish	1.25	1.37	0.77	0.83	0.80	0.74	0.41	0.36
Cocaine	0.52	0.68	0.20	0.19	0.10	0.09	0.08	0.04
Crack	0.15	0.26	0.04	0.04	0.03	0.01	0.03	0.04
Heroin	0.16	0.14	0.11	0.05	0.09	0.11	0.12	0.08
Hallucinogens	0.74	0.73	0.26	0.29	0.15	0.14	0.09	0.13
LSD	0.18	0.25	0.14	0.14	0.06	0.06	0.04	0.03
PCP	0.07	0.03	0.02	*	0.03	*	*	0.01
Ecstasy	0.70	0.56	0.20	0.22	0.12	0.05	0.05	0.05
Inhalants	0.32	0.23	0.13	0.11	0.07	0.06	0.08	0.08
Nonmedical Use of Psychotherapeutics ^{3,4}	0.95	0.94	0.40	0.42	0.36	0.35	0.24	0.19
Pain Relievers	0.78	0.84	0.35	0.34	0.31	0.30	0.24	0.17
OxyContin®	0.23	0.32	0.11	0.13	0.15	0.08	0.11	0.03
Tranquilizers	0.54	0.51	0.23	0.23	0.18	0.14	0.17	0.09
Stimulants ⁴	0.48	0.64	0.20	0.20	0.09	0.18	0.06	0.08
Methamphetamine ⁴	0.23	0.37	0.10	0.08	0.03	0.08	0.04	0.03
Sedatives	0.20	0.16	0.07	0.06	0.05	0.03	0.03	0.03
ILLICIT DRUGS OTHER THAN MARIJUANA²	1.10	1.09	0.47	0.49	0.39	0.39	0.27	0.24

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.13P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	0.3444		0.0009		0.2745		0.2345	
Marijuana and Hashish	0.6329		0.0004		0.2488		0.4889	
Cocaine	0.1812		0.7266		0.7144		0.3431	
Crack	0.8307		0.4990		0.7599		0.9968	
Heroin	0.4671		0.0867		0.5702		0.1711	
Hallucinogens	0.9122		0.5520		0.9302		0.4291	
LSD	0.6703		0.4543		0.9673		0.7999	
PCP	0.3416		*		*		*	
Ecstasy	0.5685		0.7208		0.1424		0.9077	
Inhalants	0.7238		0.5116		0.7208		0.7841	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.6235		0.3666		0.7831		0.2936	
Pain Relievers	0.7274		0.1214		0.9475		0.0983	
OxyContin®	0.6894		0.9270		0.2882		0.0797	
Tranquilizers	0.8891		0.3595		0.0062		0.6655	
Stimulants ⁴	0.1593		0.9760		0.1686		0.3100	
Methamphetamine ⁴	0.1568		0.9083		0.9626		0.9158	
Sedatives	0.5903		0.3576		0.3562		0.8406	
ILLICIT DRUGS OTHER THAN MARIJUANA²	0.7083		0.7118		0.9598		0.4843	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.14C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	215	217	246	237	247	261	195	231
Marijuana and Hashish	192	193	216	217	213	229	153	191
Cocaine	85	93	81	54	57	58	46	43
Crack	62	33	58	26	9	38	38	27
Heroin	21	12	16	20	22	21	29	23
Hallucinogens	47	48	56	49	25	100	13	28
LSD	25	17	6	25	*	*	*	*
PCP	11	*	4	11	*	*	*	10
Ecstasy	32	39	46	35	23	66	7	10
Inhalants	15	30	51	22	18	35	7	29
Nonmedical Use of Psychotherapeutics ^{3,4}	101	118	127	124	126	117	123	138
Pain Relievers	86	102	100	99	93	86	105	114
OxyContin®	27	22	10	36	6	18	22	63
Tranquilizers	60	55	87	64	83	44	76	60
Stimulants ⁴	37	52	59	62	54	53	37	46
Methamphetamine ⁴	21	38	52	38	28	43	33	43
Sedatives	8	19	35	14	23	45	20	44
ILLICIT DRUGS OTHER THAN MARIJUANA²	131	138	153	138	138	165	129	147

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.14D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	1.47	1.49	0.71	0.67	0.36	0.37	0.22	0.25
Marijuana and Hashish	1.36	1.37	0.64	0.62	0.31	0.33	0.17	0.21
Cocaine	0.68	0.73	0.25	0.16	0.08	0.08	0.05	0.05
Crack	0.50	0.27	0.18	0.08	0.01	0.06	0.04	0.03
Heroin	0.17	0.10	0.05	0.06	0.03	0.03	0.03	0.03
Hallucinogens	0.38	0.39	0.17	0.15	0.04	0.15	0.01	0.03
LSD	0.20	0.14	0.02	0.08	*	*	*	*
PCP	0.09	*	0.01	0.03	*	*	*	0.01
Ecstasy	0.26	0.32	0.15	0.11	0.03	0.10	0.01	0.01
Inhalants	0.12	0.24	0.16	0.07	0.03	0.05	0.01	0.03
Nonmedical Use of Psychotherapeutics ^{3,4}	0.78	0.90	0.39	0.37	0.18	0.17	0.14	0.15
Pain Relievers	0.68	0.79	0.31	0.30	0.14	0.13	0.12	0.13
OxyContin®	0.22	0.18	0.03	0.11	0.01	0.03	0.02	0.07
Tranquilizers	0.48	0.44	0.27	0.19	0.12	0.06	0.09	0.07
Stimulants ⁴	0.29	0.42	0.18	0.19	0.08	0.08	0.04	0.05
Methamphetamine ⁴	0.17	0.30	0.16	0.11	0.04	0.06	0.04	0.05
Sedatives	0.06	0.15	0.11	0.04	0.03	0.07	0.02	0.05
ILLICIT DRUGS OTHER THAN MARIJUANA²	0.99	1.05	0.46	0.41	0.20	0.24	0.14	0.16

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.14P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2013 Versus 2012

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	0.0635	0.3040	0.4999		0.5086			
Marijuana and Hashish	0.0633	0.4256	0.2572		0.5953			
Cocaine	0.8467	0.1482	0.5016		0.3057			
Crack	0.4544	0.2985	0.0216		0.3720			
Heroin	0.3500	0.7942	0.5217		0.6392			
Hallucinogens	0.7486	0.9626	0.1667		0.6287			
LSD	0.9481	0.3547	*		*			
PCP	*	0.5492	*		*			
Ecstasy	0.3014	0.9637	0.5532		0.9345			
Inhalants	0.2112	0.2636	0.1770		0.0823			
Nonmedical Use of Psychotherapeutics ^{3,4}	0.6724	0.9168	0.1733		0.5924			
Pain Relievers	0.8529	0.6643	0.4568		0.6995			
OxyContin®	0.7152	0.1031	0.0801		0.1068			
Tranquilizers	0.9075	0.5250	0.0816		0.5839			
Stimulants ⁴	0.4238	0.6712	0.8655		0.1785			
Methamphetamine ⁴	0.1495	0.7467	0.7069		0.1657			
Sedatives	0.5534	0.0591	0.6941		0.4371			
ILLICIT DRUGS OTHER THAN MARIJUANA²	0.8369	0.5159	0.8672		0.6263			

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.15C Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Numbers in Thousands, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-20 (2012)	Aged 12-20 (2013)	Aged 21+ (2012)	Aged 21+ (2013)
ILLICIT DRUGS¹	460	458	122	106	444	439
Marijuana and Hashish	406	407	114	102	389	387
Cocaine (Including Crack)	135	127	22	19	134	125
Heroin	37	33	8	5	36	33
Hallucinogens	88	129	35	30	81	126
Inhalants	62	54	22	11	57	52
Pain Relievers ²	177	174	40	35	174	173
Tranquilizers ²	141	100	24	20	139	99
Stimulants ^{2,3}	93	106	21	23	91	103
Methamphetamine ³	66	73	9	9	64	72
Sedatives ²	45	50	7	6	44	51
ILLICIT DRUGS OTHER THAN MARIJUANA¹	273	269	61	50	265	267

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.15D Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Percentages, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-20 (2012)	Aged 12-20 (2013)	Aged 21+ (2012)	Aged 21+ (2013)
ILLICIT DRUGS¹	0.32	0.32	0.83	0.87	0.33	0.32
Marijuana and Hashish	0.28	0.28	0.80	0.87	0.29	0.29
Cocaine (Including Crack)	0.10	0.09	0.23	0.22	0.11	0.10
Heroin	0.03	0.02	0.09	0.05	0.03	0.03
Hallucinogens	0.07	0.09	0.37	0.34	0.06	0.10
Inhalants	0.05	0.04	0.23	0.13	0.05	0.04
Pain Relievers ²	0.13	0.13	0.43	0.39	0.14	0.13
Tranquilizers ²	0.10	0.07	0.26	0.23	0.11	0.08
Stimulants ^{2,3}	0.07	0.08	0.22	0.25	0.07	0.08
Methamphetamine ³	0.05	0.05	0.10	0.10	0.05	0.06
Sedatives ²	0.03	0.04	0.08	0.07	0.03	0.04
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.20	0.19	0.61	0.53	0.21	0.20

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.15P Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: *P* Values from Tests of Differences of Percentages, 2013 Versus 2012

Drug	Total (2012)	Total (2013)	Aged 12-20 (2012)	Aged 12-20 (2013)	Aged 21+ (2012)	Aged 21+ (2013)
ILLICIT DRUGS¹	0.4653		0.9149		0.2646	
Marijuana and Hashish	0.2169		0.5427		0.1263	
Cocaine (Including Crack)	0.7908		0.7950		0.7790	
Heroin	0.8159		0.1512		0.9530	
Hallucinogens	0.3213		0.5051		0.1587	
Inhalants	0.6184		0.0041		0.6575	
Pain Relievers ²	0.2093		0.0049		0.5433	
Tranquilizers ²	0.0350		0.1539		0.0704	
Stimulants ^{2,3}	0.5430		0.5921		0.4107	
Methamphetamine ³	0.4471		0.8456		0.4133	
Sedatives ²	0.3663		0.6021		0.4187	
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.4233		0.0358		0.7937	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.16C Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Numbers in Thousands, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-20 (2012)	Aged 12-20 (2013)	Aged 21+ (2012)	Aged 21+ (2013)
ILLICIT DRUGS¹	302	300	84	69	290	288
Marijuana and Hashish	285	284	83	69	274	271
Cocaine (Including Crack)	100	79	13	6	99	79
Heroin	24	25	4	1	24	25
Hallucinogens	27	75	8	6	26	75
Inhalants	6	9	5	2	*	9
Pain Relievers ²	73	90	11	7	72	89
Tranquilizers ²	38	31	5	7	38	30
Stimulants ²	37	17	3	9	37	15
Methamphetamine ³	58	40	6	5	57	40
Sedatives ²	12	19	1	4	11	19
ILLICIT DRUGS OTHER THAN MARIJUANA¹	132	140	21	15	130	139

*Low precision; no estimate reported.

NOTE: Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.16D Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Percentages, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-20 (2012)	Aged 12-20 (2013)	Aged 21+ (2012)	Aged 21+ (2013)
ILLICIT DRUGS¹	0.21	0.21	0.74	0.72	0.22	0.22
Marijuana and Hashish	0.20	0.20	0.73	0.72	0.21	0.21
Cocaine (Including Crack)	0.07	0.06	0.14	0.07	0.08	0.06
Heroin	0.02	0.02	0.04	0.01	0.02	0.02
Hallucinogens	0.02	0.05	0.09	0.07	0.02	0.06
Inhalants	0.00	0.01	0.06	0.02	*	0.01
Pain Relievers ²	0.05	0.07	0.12	0.09	0.06	0.07
Tranquilizers ²	0.03	0.02	0.06	0.08	0.03	0.02
Stimulants ²	0.03	0.01	0.04	0.10	0.03	0.01
Methamphetamine ³	0.04	0.03	0.06	0.05	0.04	0.03
Sedatives ²	0.01	0.01	0.01	0.05	0.01	0.01
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.10	0.10	0.23	0.18	0.10	0.11

*Low precision; no estimate reported.

NOTE: Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.16P Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Drug	Total (2012)	Total (2013)	Aged 12-20 (2012)	Aged 12-20 (2013)	Aged 21+ (2012)	Aged 21+ (2013)
ILLICIT DRUGS¹	0.4119		0.4658		0.6779	
Marijuana and Hashish	0.3508		0.7576		0.5476	
Cocaine (Including Crack)	0.8148		0.4508		0.8790	
Heroin	0.4188		0.3048		0.3525	
Hallucinogens	0.6438		0.0845		0.4817	
Inhalants	0.9399		0.1442		*	
Pain Relievers ²	0.7800		0.0305		0.9775	
Tranquilizers ²	0.4180		0.8148		0.4427	
Stimulants ²	0.3179		0.1780		0.1847	
Methamphetamine ³	0.4467		0.8626		0.4515	
Sedatives ²	0.5971		0.5028		0.6788	
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.7339		0.0331		0.9701	

*Low precision; no estimate reported.

NOTE: Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.17C Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
ILLICIT DRUGS¹	460	458	48	46	163	156	425	421
Marijuana and Hashish	406	407	45	45	156	148	371	376
Cocaine (Including Crack)	135	127	6	7	31	33	130	122
Heroin	37	33	*	4	14	11	34	32
Hallucinogens	88	129	14	13	45	44	76	122
Inhalants	62	54	13	7	19	14	56	51
Pain Relievers ²	177	174	21	17	55	56	170	165
Tranquilizers ²	141	100	11	9	36	33	136	94
Stimulants ^{2,3}	93	106	12	8	31	37	88	97
Methamphetamine ³	66	73	5	4	14	18	63	69
Sedatives ²	45	50	4	3	12	8	43	51
ILLICIT DRUGS OTHER THAN MARIJUANA¹	273	269	28	23	82	79	253	258

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.17D Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
ILLCIT DRUGS¹	0.32	0.32	1.20	1.28	0.54	0.58	0.36	0.35
Marijuana and Hashish	0.28	0.28	1.17	1.25	0.54	0.55	0.32	0.32
Cocaine (Including Crack)	0.10	0.09	0.19	0.24	0.14	0.16	0.12	0.11
Heroin	0.03	0.02	*	0.12	0.07	0.05	0.03	0.03
Hallucinogens	0.07	0.09	0.42	0.46	0.21	0.21	0.07	0.11
Inhalants	0.05	0.04	0.41	0.24	0.09	0.07	0.05	0.05
Pain Relievers ²	0.13	0.13	0.61	0.59	0.26	0.26	0.15	0.15
Tranquilizers ²	0.10	0.07	0.34	0.31	0.17	0.15	0.12	0.08
Stimulants ^{2,3}	0.07	0.08	0.37	0.28	0.14	0.17	0.08	0.09
Methamphetamine ³	0.05	0.05	0.16	0.13	0.07	0.09	0.06	0.06
Sedatives ²	0.03	0.04	0.12	0.10	0.06	0.04	0.04	0.05
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.20	0.19	0.79	0.75	0.35	0.35	0.22	0.23

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.17P Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Drug	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
ILLICIT DRUGS¹	0.4653		1.0000		0.1973		0.4935	
Marijuana and Hashish	0.2169		0.3872		0.1905		0.2660	
Cocaine (Including Crack)	0.7908		0.4387		0.7683		0.7102	
Heroin	0.8159		*		0.2456		0.9424	
Hallucinogens	0.3213		0.8351		0.9868		0.2475	
Inhalants	0.6184		0.0025		0.3680		0.6682	
Pain Relievers ²	0.2093		0.0368		0.1335		0.5957	
Tranquilizers ²	0.0350		0.2911		0.0541		0.1402	
Stimulants ^{2,3}	0.5430		0.0388		0.2913		0.5663	
Methamphetamine ³	0.4471		0.1939		0.2975		0.5152	
Sedatives ²	0.3663		0.7597		0.2140		0.5489	
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.4233		0.0372		0.5490		0.7217	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.18C Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
ILLCIT DRUGS¹	302	300	34	31	113	107	276	277
Marijuana and Hashish	285	284	33	31	110	103	260	261
Cocaine (Including Crack)	100	79	3	2	18	22	98	77
Heroin	24	25	*	*	5	8	24	23
Hallucinogens	27	75	3	3	13	12	24	74
Inhalants	6	9	3	*	4	3	*	8
Pain Relievers ²	73	90	5	5	23	16	69	88
Tranquilizers ²	38	31	2	5	12	11	36	28
Stimulants ²	37	17	3	3	7	9	36	14
Methamphetamine ³	58	40	2	2	11	7	56	40
Sedatives ²	12	19	1	*	2	6	11	18
ILLCIT DRUGS OTHER THAN MARIJUANA¹	132	140	9	9	35	34	126	135

*Low precision; no estimate reported.

NOTE: Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.18D Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
ILLCIT DRUGS¹	0.21	0.21	0.99	1.01	0.45	0.45	0.24	0.24
Marijuana and Hashish	0.20	0.20	0.97	1.03	0.45	0.43	0.23	0.23
Cocaine (Including Crack)	0.07	0.06	0.09	0.07	0.08	0.11	0.09	0.07
Heroin	0.02	0.02	*	*	0.02	0.04	0.02	0.02
Hallucinogens	0.02	0.05	0.11	0.10	0.06	0.06	0.02	0.07
Inhalants	0.00	0.01	0.11	*	0.02	0.01	*	0.01
Pain Relievers ²	0.05	0.07	0.17	0.17	0.11	0.08	0.06	0.08
Tranquilizers ²	0.03	0.02	0.08	0.19	0.06	0.05	0.03	0.02
Stimulants ²	0.03	0.01	0.09	0.11	0.04	0.04	0.03	0.01
Methamphetamine ³	0.04	0.03	0.06	0.06	0.05	0.03	0.05	0.04
Sedatives ²	0.01	0.01	0.04	*	0.01	0.03	0.01	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.10	0.10	0.29	0.30	0.16	0.16	0.11	0.12

*Low precision; no estimate reported.

NOTE: Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.18P Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Drug	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
ILLICIT DRUGS¹	0.4119	0.7255		0.8279		0.4877		
Marijuana and Hashish	0.3508	0.9590		0.9808		0.4452		
Cocaine (Including Crack)	0.8148	0.5259		0.3460		0.6622		
Heroin	0.4188	*		0.1331		0.6836		
Hallucinogens	0.6438	0.3251		0.3490		0.4527		
Inhalants	0.9399	*		0.6951		*		
Pain Relievers ²	0.7800	0.2880		0.0591		0.7889		
Tranquilizers ²	0.4180	0.7355		0.4295		0.5424		
Stimulants ²	0.3179	0.8277		0.8105		0.2814		
Methamphetamine ³	0.4467	0.7228		0.6168		0.4993		
Sedatives ²	0.5971	*		0.3029		0.7526		
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.7339	0.1369		0.5983		0.9276		

*Low precision; no estimate reported.

NOTE: Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.