

Table 7.29C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2013

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
TOTAL												
TOBACCO PRODUCTS¹	832	803	794	845	850	844	871	831	852	846	882	835
Cigarettes	795	789	768	777	807	809	810	797	780	816	839	798
Smokeless Tobacco	279	284	245	267	284	283	312	291	314	278	298	300
Cigars	350	331	323	354	352	337	366	361	363	373	388	354
Pipe Tobacco	166	131	147	146	167	170	147	151	147	144	184	151
ALCOHOL	989	919	957	975	968	1,019	983	957	990	1,006	1,016	1,076
Binge Alcohol Use ²	732	682	698	701	746	770	735	783	752	734	797	804
Heavy Alcohol Use ²	392	380	382	379	427	429	414	406	422	423	448	438
MALE												
TOBACCO PRODUCTS¹	572	573	570	576	585	598	628	602	626	581	618	612
Cigarettes	527	545	537	515	545	577	565	565	536	537	556	549
Smokeless Tobacco	264	256	234	241	273	252	296	283	302	264	281	282
Cigars	327	305	290	324	320	312	333	336	321	336	351	312
Pipe Tobacco	139	127	139	141	160	168	131	144	137	135	170	146
ALCOHOL	656	599	626	621	610	666	632	651	675	674	675	693
Binge Alcohol Use ²	540	517	513	529	559	585	586	605	577	567	625	607
Heavy Alcohol Use ²	346	321	319	321	354	374	357	349	353	355	382	367
FEMALE												
TOBACCO PRODUCTS¹	513	507	497	543	511	511	558	505	535	542	554	524
Cigarettes	501	488	491	538	503	493	553	498	526	532	539	513
Smokeless Tobacco	100	126	65	87	76	113	75	51	78	82	103	94
Cigars	114	123	122	112	131	116	114	130	153	142	146	161
Pipe Tobacco	90	35	43	42	46	30	55	37	54	46	61	35
ALCOHOL	644	600	664	637	645	676	673	687	635	684	686	736
Binge Alcohol Use ²	415	382	390	402	399	395	398	441	441	441	427	463
Heavy Alcohol Use ²	177	174	174	163	169	171	192	193	201	207	203	232

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2013.

Table 7.29D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Percentages, 2002-2013

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
TOTAL												
TOBACCO PRODUCTS¹	0.35	0.34	0.33	0.35	0.35	0.34	0.35	0.33	0.34	0.33	0.34	0.32
Cigarettes	0.34	0.33	0.32	0.32	0.33	0.33	0.32	0.32	0.31	0.32	0.32	0.30
Smokeless Tobacco	0.12	0.12	0.10	0.11	0.12	0.11	0.12	0.12	0.12	0.11	0.11	0.11
Cigars	0.15	0.14	0.13	0.15	0.14	0.14	0.15	0.14	0.14	0.14	0.15	0.14
Pipe Tobacco	0.07	0.06	0.06	0.06	0.07	0.07	0.06	0.06	0.06	0.06	0.07	0.06
ALCOHOL	0.42	0.39	0.40	0.40	0.39	0.41	0.39	0.38	0.39	0.39	0.39	0.41
Binge Alcohol Use ²	0.31	0.29	0.29	0.29	0.30	0.31	0.29	0.31	0.30	0.29	0.31	0.31
Heavy Alcohol Use ²	0.17	0.16	0.16	0.16	0.17	0.17	0.17	0.16	0.17	0.16	0.17	0.17
MALE												
TOBACCO PRODUCTS¹	0.50	0.50	0.49	0.49	0.49	0.50	0.52	0.49	0.51	0.47	0.49	0.48
Cigarettes	0.46	0.47	0.46	0.44	0.46	0.48	0.47	0.46	0.43	0.43	0.44	0.43
Smokeless Tobacco	0.23	0.22	0.20	0.20	0.23	0.21	0.24	0.23	0.25	0.21	0.22	0.22
Cigars	0.29	0.27	0.25	0.27	0.27	0.26	0.27	0.27	0.26	0.27	0.28	0.25
Pipe Tobacco	0.12	0.11	0.12	0.12	0.13	0.14	0.11	0.12	0.11	0.11	0.14	0.12
ALCOHOL	0.58	0.52	0.54	0.53	0.51	0.55	0.52	0.53	0.55	0.54	0.54	0.55
Binge Alcohol Use ²	0.48	0.45	0.44	0.45	0.47	0.49	0.48	0.49	0.47	0.45	0.50	0.48
Heavy Alcohol Use ²	0.30	0.28	0.27	0.27	0.30	0.31	0.29	0.29	0.29	0.28	0.30	0.29
FEMALE												
TOBACCO PRODUCTS¹	0.42	0.41	0.40	0.43	0.40	0.40	0.43	0.39	0.41	0.41	0.41	0.39
Cigarettes	0.41	0.40	0.40	0.43	0.40	0.39	0.43	0.38	0.40	0.40	0.40	0.38
Smokeless Tobacco	0.08	0.10	0.05	0.07	0.06	0.09	0.06	0.04	0.06	0.06	0.08	0.07
Cigars	0.09	0.10	0.10	0.09	0.10	0.09	0.09	0.10	0.12	0.11	0.11	0.12
Pipe Tobacco	0.07	0.03	0.03	0.03	0.04	0.02	0.04	0.03	0.04	0.03	0.05	0.03
ALCOHOL	0.53	0.49	0.54	0.51	0.51	0.53	0.52	0.53	0.49	0.51	0.51	0.54
Binge Alcohol Use ²	0.34	0.31	0.31	0.32	0.32	0.31	0.31	0.34	0.34	0.33	0.32	0.34
Heavy Alcohol Use ²	0.15	0.14	0.14	0.13	0.13	0.13	0.15	0.15	0.15	0.16	0.15	0.17

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2013.

Table 7.29P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: P Values from Tests of Differences of Percentages, 2013 Versus 2002-2012

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
TOTAL												
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0268	0.0050	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0654	0.0411	
Smokeless Tobacco	0.7737	0.5108	0.0122	0.2125	0.9240	0.5020	0.4694	0.7038	0.3970	0.3200	0.4649	
Cigars	0.0007	0.0007	0.0000	0.0000	0.0000	0.0007	0.0037	0.0037	0.0120	0.1914	0.0351	
Pipe Tobacco	0.3067	0.0208	0.2228	0.6750	0.3890	0.6546	0.1798	0.7173	0.8419	0.6559	0.2396	
ALCOHOL	0.0404	0.0002	0.0010	0.5512	0.0328	0.0865	0.3210	0.6166	0.5136	0.5032	0.9167	
Binge Alcohol Use ²	0.9693	0.5230	0.7443	0.5567	0.7249	0.3069	0.2678	0.0629	0.6502	0.5707	0.8875	
Heavy Alcohol Use ²	0.0617	0.0347	0.0060	0.1892	0.0149	0.0213	0.0051	0.0274	0.1360	0.5372	0.3092	
MALE												
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.0002	0.0777	0.0043	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0083	0.0041	0.2462	0.0911	
Smokeless Tobacco	0.7938	0.3605	0.0231	0.2243	0.7358	0.6361	0.3087	0.4446	0.3354	0.3352	0.3536	
Cigars	0.0000	0.0002	0.0000	0.0000	0.0000	0.0000	0.0001	0.0021	0.0106	0.1749	0.0163	
Pipe Tobacco	0.3140	0.1039	0.4622	0.4980	0.2015	0.9378	0.1137	0.9190	0.6581	0.5658	0.4243	
ALCOHOL	0.7186	0.7711	0.8118	0.2010	0.8874	0.5510	0.4895	0.5778	0.7668	0.6480	0.4621	
Binge Alcohol Use ²	0.1309	0.2603	0.1749	0.5729	0.0777	0.0287	0.0358	0.0387	0.2837	0.7655	0.7559	
Heavy Alcohol Use ²	0.0032	0.0275	0.0049	0.0410	0.0046	0.0098	0.0050	0.0463	0.1547	0.7058	0.3506	
FEMALE												
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0001	0.0199	0.1298	0.2473	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0029	0.1041	0.1788	
Smokeless Tobacco	0.9811	0.5908	0.2339	0.6812	0.1337	0.4539	0.3250	0.1124	0.5658	0.8443	0.6792	
Cigars	0.0752	0.8689	0.5735	0.2799	0.6572	0.2776	0.1101	0.8599	0.6097	0.8188	1.0000	
Pipe Tobacco	0.8123	0.0041	0.0536	0.3338	0.1848	0.0063	0.7094	0.2446	0.5476	0.6797	0.1490	
ALCOHOL	0.0007	0.0000	0.0000	0.0322	0.0022	0.0543	0.0310	0.2018	0.1905	0.5979	0.5396	
Binge Alcohol Use ²	0.0501	0.0101	0.0191	0.0864	0.1003	0.2189	0.2994	0.6962	0.4783	0.5824	0.8719	
Heavy Alcohol Use ²	0.1999	0.5772	0.4180	0.3261	0.9630	0.8905	0.4660	0.3310	0.6040	0.5843	0.6413	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2013.

Table 7.30C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Numbers in Thousands, 2002-2013

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
TOTAL												
TOBACCO PRODUCTS¹	81	80	80	78	72	76	69	71	69	68	61	59
Cigarettes	74	73	76	71	66	67	61	64	63	60	55	51
Smokeless Tobacco	27	30	32	29	31	36	30	32	30	30	31	31
Cigars	47	44	44	46	41	45	41	39	36	41	32	32
Pipe Tobacco	14	18	19	17	18	19	17	21	17	18	18	17
ALCOHOL	80	82	81	81	82	87	79	79	80	77	78	73
Binge Alcohol Use ²	67	67	73	65	69	69	57	62	62	55	56	54
Heavy Alcohol Use ²	31	33	36	32	33	37	30	30	28	25	25	23
MALE												
TOBACCO PRODUCTS¹	59	56	55	59	54	57	50	52	52	53	47	45
Cigarettes	53	47	52	53	46	48	42	44	45	46	41	36
Smokeless Tobacco	26	29	30	28	29	34	28	30	28	28	29	28
Cigars	37	36	37	38	34	37	34	33	31	31	26	28
Pipe Tobacco	13	17	15	14	15	16	12	18	14	14	12	12
ALCOHOL	55	57	59	57	57	61	53	58	57	53	55	52
Binge Alcohol Use ²	46	46	52	48	47	51	41	48	44	42	41	40
Heavy Alcohol Use ²	24	23	27	27	26	28	23	23	23	22	19	18
FEMALE												
TOBACCO PRODUCTS¹	51	52	53	50	47	48	46	46	43	43	37	40
Cigarettes	49	50	52	48	46	46	44	44	40	40	34	37
Smokeless Tobacco	9	6	11	8	10	8	10	9	12	10	8	9
Cigars	26	23	25	24	24	22	21	25	20	24	19	16
Pipe Tobacco	9	7	13	11	10	11	12	11	10	12	12	12
ALCOHOL	58	60	59	58	59	61	56	52	54	53	54	52
Binge Alcohol Use ²	46	46	50	43	49	45	42	40	43	37	38	37
Heavy Alcohol Use ²	19	23	23	20	21	23	20	19	17	15	16	16

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2013.

Table 7.30D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Percentages, 2002-2013

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
TOTAL												
TOBACCO PRODUCTS¹	0.33	0.32	0.32	0.31	0.29	0.30	0.28	0.29	0.28	0.27	0.25	0.24
Cigarettes	0.30	0.29	0.30	0.28	0.26	0.27	0.25	0.26	0.26	0.24	0.22	0.20
Smokeless Tobacco	0.11	0.12	0.13	0.12	0.12	0.14	0.12	0.13	0.12	0.12	0.12	0.12
Cigars	0.19	0.17	0.18	0.18	0.16	0.18	0.16	0.16	0.15	0.16	0.13	0.13
Pipe Tobacco	0.06	0.07	0.08	0.07	0.07	0.08	0.07	0.09	0.07	0.07	0.07	0.07
ALCOHOL	0.32	0.33	0.32	0.32	0.32	0.34	0.32	0.32	0.33	0.31	0.31	0.29
Binge Alcohol Use ²	0.27	0.27	0.29	0.26	0.27	0.27	0.23	0.25	0.25	0.22	0.22	0.22
Heavy Alcohol Use ²	0.12	0.13	0.14	0.13	0.13	0.15	0.12	0.12	0.11	0.10	0.10	0.09
MALE												
TOBACCO PRODUCTS¹	0.47	0.44	0.43	0.46	0.41	0.45	0.39	0.42	0.42	0.41	0.37	0.35
Cigarettes	0.42	0.37	0.40	0.41	0.36	0.37	0.33	0.35	0.36	0.36	0.32	0.28
Smokeless Tobacco	0.21	0.23	0.23	0.22	0.22	0.26	0.22	0.24	0.23	0.22	0.23	0.22
Cigars	0.30	0.28	0.28	0.29	0.26	0.29	0.27	0.27	0.25	0.25	0.20	0.22
Pipe Tobacco	0.10	0.13	0.11	0.11	0.12	0.12	0.10	0.14	0.11	0.11	0.09	0.10
ALCOHOL	0.43	0.45	0.45	0.44	0.44	0.47	0.41	0.46	0.46	0.42	0.43	0.41
Binge Alcohol Use ²	0.37	0.36	0.40	0.37	0.36	0.40	0.32	0.38	0.35	0.33	0.32	0.32
Heavy Alcohol Use ²	0.19	0.18	0.21	0.21	0.20	0.22	0.18	0.18	0.18	0.17	0.15	0.14
FEMALE												
TOBACCO PRODUCTS¹	0.42	0.42	0.43	0.40	0.38	0.39	0.38	0.38	0.36	0.35	0.30	0.33
Cigarettes	0.40	0.41	0.42	0.39	0.37	0.37	0.36	0.37	0.34	0.33	0.28	0.30
Smokeless Tobacco	0.08	0.05	0.09	0.07	0.08	0.07	0.08	0.08	0.11	0.08	0.06	0.07
Cigars	0.21	0.19	0.20	0.19	0.19	0.18	0.17	0.21	0.17	0.20	0.16	0.13
Pipe Tobacco	0.07	0.06	0.10	0.09	0.08	0.09	0.10	0.09	0.09	0.10	0.10	0.10
ALCOHOL	0.48	0.49	0.48	0.47	0.47	0.49	0.46	0.43	0.45	0.43	0.44	0.43
Binge Alcohol Use ²	0.38	0.38	0.41	0.35	0.39	0.37	0.35	0.33	0.36	0.30	0.31	0.30
Heavy Alcohol Use ²	0.16	0.19	0.19	0.16	0.17	0.18	0.16	0.15	0.14	0.12	0.13	0.13

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2013.

Table 7.30P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: P Values from Tests of Differences of Percentages, 2013 Versus 2002-2012

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
TOTAL												
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0188	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0013	
Smokeless Tobacco	0.9464	0.6181	0.0822	0.5362	0.0158	0.0097	0.1191	0.0215	0.0290	0.4981	0.5338	
Cigars	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.1311	
Pipe Tobacco	0.3328	1.0000	0.5306	0.8570	0.8337	0.4497	0.4705	0.0082	0.9605	0.3743	0.8496	
ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0021	
Binge Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0026	
Heavy Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0011	0.0098	0.3321	
MALE												
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0937	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0076	
Smokeless Tobacco	0.9910	0.3426	0.0616	0.4658	0.0149	0.0061	0.1283	0.0296	0.0796	0.5987	0.4548	
Cigars	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0007	0.0002	0.3286	
Pipe Tobacco	0.9463	0.1144	0.2628	0.2868	0.2220	0.1595	0.5218	0.0006	0.5167	0.4866	0.9383	
ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0005	0.0193	
Binge Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.0067	0.0892	
Heavy Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0020	0.0238	0.8916	
FEMALE												
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.1022	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0596	
Smokeless Tobacco	0.8763	0.0956	1.0000	0.8101	0.7442	0.9932	0.6215	0.3638	0.1189	0.5998	0.7262	
Cigars	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0014	0.0000	0.2230	
Pipe Tobacco	0.1325	0.0219	0.7791	0.1528	0.2484	0.6276	0.6879	0.8681	0.3906	0.5734	0.7272	
ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0088	0.0192	0.0328	
Binge Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0023	0.0116	0.0066	
Heavy Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0016	0.0000	0.1487	0.2388	0.1826	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2013.

Table 7.31C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Numbers in Thousands, 2002-2013

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
TOTAL												
TOBACCO PRODUCTS¹	150	151	161	155	160	162	153	169	166	167	164	170
Cigarettes	150	151	159	152	156	159	149	162	160	162	161	162
Smokeless Tobacco	59	56	60	64	65	63	68	78	81	70	70	71
Cigars	83	84	97	92	94	92	96	99	101	100	93	100
Pipe Tobacco	24	25	30	37	33	32	34	40	41	48	39	49
ALCOHOL	164	160	164	166	168	170	163	174	169	184	170	183
Binge Alcohol Use ²	161	155	166	171	169	169	167	182	168	189	175	182
Heavy Alcohol Use ²	112	114	120	120	125	114	124	128	127	125	121	107
MALE												
TOBACCO PRODUCTS¹	100	106	106	110	109	116	109	121	121	113	118	121
Cigarettes	99	109	105	107	108	116	106	121	113	111	113	117
Smokeless Tobacco	56	55	58	60	62	61	63	73	75	65	65	67
Cigars	73	72	83	79	78	81	80	81	84	89	80	86
Pipe Tobacco	22	21	27	33	30	30	29	35	33	39	34	41
ALCOHOL	111	100	110	101	108	110	109	117	112	119	118	121
Binge Alcohol Use ²	110	105	113	111	115	116	118	130	121	126	124	121
Heavy Alcohol Use ²	87	91	88	96	95	94	95	99	101	87	97	85
FEMALE												
TOBACCO PRODUCTS¹	101	93	102	101	102	97	101	98	96	105	104	101
Cigarettes	102	92	101	101	98	95	100	96	97	102	102	97
Smokeless Tobacco	8	11	10	12	13	13	14	18	20	14	14	18
Cigars	38	43	44	41	44	43	47	45	49	50	46	46
Pipe Tobacco	12	12	12	16	11	14	17	15	23	26	19	22
ALCOHOL	102	114	104	112	112	114	107	110	109	123	118	117
Binge Alcohol Use ²	99	100	96	104	103	110	102	105	101	116	110	112
Heavy Alcohol Use ²	55	59	60	56	63	60	65	61	59	72	63	65

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2013.

Table 7.31D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Percentages, 2002-2013

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
TOTAL												
TOBACCO PRODUCTS¹	0.48	0.48	0.50	0.48	0.49	0.50	0.47	0.50	0.49	0.49	0.47	0.49
Cigarettes	0.48	0.47	0.49	0.47	0.48	0.49	0.45	0.48	0.47	0.47	0.47	0.46
Smokeless Tobacco	0.19	0.18	0.19	0.20	0.20	0.19	0.20	0.23	0.24	0.20	0.20	0.20
Cigars	0.27	0.26	0.30	0.28	0.29	0.28	0.29	0.29	0.30	0.29	0.27	0.29
Pipe Tobacco	0.08	0.08	0.09	0.11	0.10	0.10	0.10	0.12	0.12	0.14	0.11	0.14
ALCOHOL	0.53	0.50	0.51	0.51	0.51	0.52	0.49	0.52	0.50	0.54	0.49	0.53
Binge Alcohol Use ²	0.52	0.49	0.52	0.53	0.52	0.52	0.51	0.54	0.49	0.55	0.51	0.52
Heavy Alcohol Use ²	0.36	0.36	0.37	0.37	0.38	0.35	0.38	0.38	0.37	0.36	0.35	0.31
MALE												
TOBACCO PRODUCTS¹	0.64	0.67	0.66	0.67	0.66	0.70	0.66	0.72	0.70	0.66	0.68	0.70
Cigarettes	0.64	0.69	0.65	0.66	0.66	0.70	0.64	0.71	0.65	0.65	0.65	0.67
Smokeless Tobacco	0.36	0.34	0.36	0.37	0.38	0.37	0.38	0.43	0.43	0.38	0.38	0.38
Cigars	0.47	0.45	0.51	0.48	0.48	0.49	0.48	0.48	0.49	0.52	0.46	0.50
Pipe Tobacco	0.14	0.13	0.17	0.20	0.18	0.18	0.17	0.21	0.19	0.23	0.20	0.24
ALCOHOL	0.71	0.63	0.68	0.62	0.66	0.67	0.66	0.69	0.65	0.69	0.68	0.69
Binge Alcohol Use ²	0.71	0.66	0.70	0.68	0.70	0.70	0.71	0.77	0.70	0.73	0.71	0.69
Heavy Alcohol Use ²	0.56	0.57	0.54	0.59	0.58	0.57	0.57	0.59	0.59	0.51	0.56	0.49
FEMALE												
TOBACCO PRODUCTS¹	0.65	0.59	0.64	0.63	0.63	0.59	0.62	0.59	0.57	0.61	0.60	0.58
Cigarettes	0.66	0.58	0.63	0.63	0.60	0.58	0.61	0.58	0.58	0.60	0.59	0.56
Smokeless Tobacco	0.05	0.07	0.06	0.08	0.08	0.08	0.08	0.11	0.12	0.08	0.08	0.11
Cigars	0.25	0.27	0.27	0.26	0.27	0.26	0.29	0.27	0.29	0.29	0.27	0.27
Pipe Tobacco	0.08	0.08	0.08	0.10	0.07	0.08	0.10	0.09	0.14	0.15	0.11	0.13
ALCOHOL	0.66	0.72	0.65	0.69	0.69	0.70	0.65	0.66	0.65	0.72	0.68	0.68
Binge Alcohol Use ²	0.64	0.63	0.60	0.65	0.64	0.68	0.62	0.63	0.60	0.68	0.64	0.64
Heavy Alcohol Use ²	0.35	0.37	0.37	0.35	0.39	0.37	0.39	0.36	0.35	0.42	0.37	0.38

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2013.

Table 7.31P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: P Values from Tests of Differences of Percentages, 2013 Versus 2002-2012

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
TOTAL												
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	0.1211	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0600	
Smokeless Tobacco	0.0006	0.0000	0.0017	0.0166	0.0440	0.0475	0.1247	0.2938	0.0521	0.2066	0.3403	
Cigars	0.0089	0.0002	0.0000	0.0000	0.0000	0.0000	0.0004	0.0002	0.0012	0.0290	0.0531	
Pipe Tobacco	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0266	0.0479	0.1060	0.0400	
ALCOHOL	0.2553	0.0176	0.2340	0.0879	0.0007	0.0302	0.0384	0.0049	0.0142	0.1541	0.3966	
Binge Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0111	0.0211	
Heavy Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0669	0.0025	
MALE												
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.0118	0.0003	0.0072	0.2026	0.7024	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0010	0.0000	0.0535	0.0324	0.8090	
Smokeless Tobacco	0.0063	0.0002	0.0110	0.0425	0.1097	0.0868	0.2484	0.2989	0.0500	0.3324	0.5568	
Cigars	0.0063	0.0003	0.0000	0.0000	0.0000	0.0000	0.0002	0.0001	0.0053	0.1360	0.0319	
Pipe Tobacco	0.0000	0.0000	0.0001	0.0036	0.0010	0.0000	0.0001	0.1905	0.0132	0.0913	0.1025	
ALCOHOL	0.0030	0.0000	0.0063	0.0000	0.0001	0.0020	0.0478	0.0003	0.0003	0.3324	0.5380	
Binge Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0001	0.0547	0.1574	
Heavy Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0937	0.0032	
FEMALE												
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0407	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0070	
Smokeless Tobacco	0.0001	0.0109	0.0011	0.0292	0.0141	0.1045	0.0322	0.8365	0.7861	0.1901	0.0823	
Cigars	0.5431	0.2391	0.0523	0.1484	0.2023	0.7503	0.3919	0.3070	0.1102	0.0677	0.8325	
Pipe Tobacco	0.0000	0.0000	0.0000	0.0007	0.0000	0.0000	0.0108	0.0109	0.8727	0.6747	0.1657	
ALCOHOL	0.1877	0.2338	0.3230	0.1404	0.2374	0.8006	0.2637	0.4501	0.9946	0.2255	0.5228	
Binge Alcohol Use ²	0.7318	0.6522	0.3153	0.5679	0.0016	0.0101	0.0032	0.0077	0.1483	0.0426	0.0374	
Heavy Alcohol Use ²	0.0134	0.0030	0.0068	0.0059	0.0000	0.0000	0.0009	0.0510	0.0167	0.3501	0.2780	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2013.

Table 7.32C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2013

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
TOTAL												
TOBACCO PRODUCTS¹	797	748	752	798	799	801	849	784	813	805	841	803
Cigarettes	754	738	718	726	760	763	785	753	747	771	796	770
Smokeless Tobacco	270	274	236	256	267	268	303	274	302	266	286	288
Cigars	331	332	311	335	332	312	352	341	340	352	372	335
Pipe Tobacco	160	126	142	141	162	166	141	141	141	135	176	143
ALCOHOL	953	887	933	936	914	986	958	931	946	978	972	1,021
Binge Alcohol Use ²	702	639	654	650	697	714	703	751	715	699	749	766
Heavy Alcohol Use ²	355	349	356	347	392	402	391	377	398	396	420	420
MALE												
TOBACCO PRODUCTS¹	553	546	552	554	566	575	613	574	599	560	605	590
Cigarettes	506	521	512	493	527	551	546	538	514	514	545	530
Smokeless Tobacco	252	247	226	232	258	238	286	267	290	252	267	270
Cigars	311	302	280	309	305	290	325	318	306	321	334	293
Pipe Tobacco	132	124	135	136	156	164	128	136	134	129	165	140
ALCOHOL	630	577	610	598	583	640	616	631	645	665	652	669
Binge Alcohol Use ²	527	492	492	507	541	562	565	585	552	552	594	583
Heavy Alcohol Use ²	323	306	307	302	333	355	340	326	337	343	363	350
FEMALE												
TOBACCO PRODUCTS¹	488	477	469	515	477	486	543	485	516	520	525	511
Cigarettes	475	460	466	508	472	469	539	481	509	510	512	500
Smokeless Tobacco	99	125	63	85	74	111	73	49	72	80	102	91
Cigars	99	117	107	98	116	104	101	116	140	130	135	152
Pipe Tobacco	89	32	39	37	43	26	50	32	47	36	57	26
ALCOHOL	628	582	654	615	614	658	656	668	620	663	662	701
Binge Alcohol Use ²	393	347	361	370	356	364	382	422	427	406	406	435
Heavy Alcohol Use ²	158	155	160	146	147	157	181	181	190	184	192	219

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2013.

Table 7.32D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Percentages, 2002-2013

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
TOTAL												
TOBACCO PRODUCTS¹	0.44	0.41	0.41	0.43	0.43	0.42	0.44	0.40	0.42	0.41	0.42	0.40
Cigarettes	0.42	0.41	0.39	0.39	0.40	0.40	0.41	0.39	0.38	0.39	0.40	0.38
Smokeless Tobacco	0.15	0.15	0.13	0.14	0.14	0.14	0.16	0.14	0.15	0.13	0.14	0.14
Cigars	0.18	0.18	0.17	0.18	0.18	0.16	0.18	0.18	0.17	0.18	0.19	0.17
Pipe Tobacco	0.09	0.07	0.08	0.08	0.09	0.09	0.07	0.07	0.07	0.07	0.09	0.07
ALCOHOL	0.53	0.49	0.51	0.51	0.49	0.52	0.50	0.48	0.48	0.49	0.48	0.50
Binge Alcohol Use ²	0.39	0.35	0.36	0.35	0.37	0.38	0.37	0.39	0.37	0.35	0.37	0.38
Heavy Alcohol Use ²	0.20	0.19	0.19	0.19	0.21	0.21	0.20	0.19	0.20	0.20	0.21	0.21
MALE												
TOBACCO PRODUCTS¹	0.65	0.63	0.63	0.62	0.63	0.63	0.67	0.62	0.64	0.59	0.63	0.61
Cigarettes	0.59	0.60	0.59	0.56	0.59	0.61	0.59	0.58	0.55	0.54	0.57	0.55
Smokeless Tobacco	0.29	0.29	0.26	0.26	0.29	0.26	0.31	0.29	0.31	0.27	0.28	0.28
Cigars	0.36	0.35	0.32	0.35	0.34	0.32	0.35	0.34	0.33	0.34	0.35	0.30
Pipe Tobacco	0.15	0.14	0.15	0.15	0.17	0.18	0.14	0.15	0.14	0.14	0.17	0.14
ALCOHOL	0.74	0.67	0.70	0.67	0.65	0.70	0.67	0.68	0.69	0.70	0.68	0.69
Binge Alcohol Use ²	0.62	0.57	0.56	0.57	0.60	0.62	0.61	0.63	0.59	0.58	0.62	0.60
Heavy Alcohol Use ²	0.38	0.35	0.35	0.34	0.37	0.39	0.37	0.35	0.36	0.36	0.38	0.36
FEMALE												
TOBACCO PRODUCTS¹	0.52	0.50	0.49	0.53	0.49	0.49	0.54	0.48	0.51	0.50	0.50	0.48
Cigarettes	0.51	0.49	0.49	0.53	0.48	0.47	0.54	0.48	0.50	0.49	0.49	0.47
Smokeless Tobacco	0.11	0.13	0.07	0.09	0.08	0.11	0.07	0.05	0.07	0.08	0.10	0.09
Cigars	0.11	0.12	0.11	0.10	0.12	0.11	0.10	0.12	0.14	0.13	0.13	0.14
Pipe Tobacco	0.10	0.03	0.04	0.04	0.04	0.03	0.05	0.03	0.05	0.03	0.05	0.02
ALCOHOL	0.67	0.61	0.68	0.64	0.63	0.66	0.66	0.66	0.61	0.64	0.63	0.66
Binge Alcohol Use ²	0.42	0.37	0.38	0.38	0.36	0.37	0.38	0.42	0.42	0.39	0.39	0.41
Heavy Alcohol Use ²	0.17	0.16	0.17	0.15	0.15	0.16	0.18	0.18	0.19	0.18	0.18	0.21

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2013.

Table 7.32P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: P Values from Tests of Differences of Percentages, 2013 Versus 2002-2012

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
TOTAL												
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0025	0.0075	0.2438	0.0148	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0082	0.0206	0.5622	0.1229	
Smokeless Tobacco	0.5607	0.7952	0.0458	0.4598	0.8644	0.6231	0.3244	0.9005	0.8962	0.4270	0.3483	
Cigars	0.0744	0.1720	0.0516	0.0250	0.0445	0.2547	0.2198	0.3045	0.2438	0.8739	0.1258	
Pipe Tobacco	0.4611	0.8153	0.8403	0.0971	0.0260	0.3557	0.8568	0.9500	0.6868	0.9841	0.0770	
ALCOHOL	0.0076	0.0000	0.0001	0.2581	0.0016	0.0139	0.1135	0.1504	0.1511	0.2690	0.6696	
Binge Alcohol Use ²	0.0902	0.0069	0.0169	0.0080	0.0660	0.4624	0.7402	0.8743	0.4201	0.1529	0.5597	
Heavy Alcohol Use ²	0.5779	0.5781	0.9217	0.1068	0.7738	0.9103	0.4540	0.6216	0.9725	0.2116	0.8245	
MALE												
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0268	0.0038	0.2115	0.0065	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0007	0.2024	0.0399	0.7160	0.1308	
Smokeless Tobacco	0.6332	0.9111	0.0580	0.4337	0.5986	0.7250	0.2231	0.7520	0.7586	0.4132	0.2951	
Cigars	0.0010	0.0399	0.0011	0.0005	0.0101	0.0355	0.0144	0.1173	0.1352	0.5074	0.0639	
Pipe Tobacco	0.7942	0.7841	0.9174	0.1609	0.0361	0.3337	0.5080	0.8943	0.8736	0.8799	0.2162	
ALCOHOL	0.7686	0.4820	0.3632	0.5766	0.3127	0.1779	0.7944	0.6467	0.5058	0.4488	0.3306	
Binge Alcohol Use ²	0.9820	0.4597	0.7161	0.1689	0.9204	0.4253	0.2277	0.5026	0.9750	0.4011	0.9635	
Heavy Alcohol Use ²	0.3967	0.9434	0.6567	0.6088	0.4531	0.3542	0.2516	0.5979	0.8675	0.3814	0.8077	
FEMALE												
TOBACCO PRODUCTS¹	0.0002	0.0003	0.0376	0.0033	0.0056	0.0269	0.0203	0.0254	0.4013	0.6749	0.4898	
Cigarettes	0.0000	0.0003	0.0134	0.0010	0.0015	0.0090	0.0081	0.0056	0.1784	0.6139	0.4521	
Smokeless Tobacco	0.5781	0.3145	0.5327	0.9890	0.2490	0.6164	0.5163	0.0692	0.4523	0.9926	0.9151	
Cigars	0.0021	0.2722	0.0196	0.0101	0.4117	0.0464	0.0097	0.3295	0.6531	0.2988	0.8123	
Pipe Tobacco	0.2450	0.8141	0.6407	0.1954	0.3471	0.8388	0.1303	0.7951	0.4045	0.5445	0.0294	
ALCOHOL	0.0003	0.0000	0.0000	0.0244	0.0002	0.0238	0.0087	0.1001	0.1286	0.3880	0.6963	
Binge Alcohol Use ²	0.0042	0.0002	0.0002	0.0068	0.0008	0.0121	0.0190	0.4618	0.1624	0.1779	0.3436	
Heavy Alcohol Use ²	0.0106	0.3426	0.5948	0.0172	0.0428	0.1246	0.6029	0.9737	0.7722	0.3245	0.9573	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2013.

Table 7.33C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2013

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
TOTAL												
TOBACCO PRODUCTS¹	829	784	793	839	838	838	872	822	840	845	874	834
Cigarettes	790	774	765	772	799	803	809	790	772	814	832	796
Smokeless Tobacco	276	283	243	264	281	280	311	288	310	275	298	299
Cigars	346	333	323	351	348	331	363	358	358	369	387	354
Pipe Tobacco	163	130	145	145	165	169	146	149	147	143	183	150
ALCOHOL	972	909	950	969	954	1,011	977	945	974	1,006	1,004	1,052
Binge Alcohol Use ²	730	675	699	701	742	766	737	784	748	736	792	799
Heavy Alcohol Use ²	389	376	383	378	425	426	414	404	421	423	448	438
MALE												
TOBACCO PRODUCTS¹	565	561	568	576	581	594	630	597	616	580	613	615
Cigarettes	521	537	532	514	542	573	565	564	529	535	552	549
Smokeless Tobacco	260	255	233	239	271	249	295	281	299	261	280	279
Cigars	322	306	291	324	316	306	332	332	318	337	349	312
Pipe Tobacco	136	127	138	141	159	168	131	143	136	135	169	146
ALCOHOL	643	590	616	616	599	658	626	644	666	678	666	687
Binge Alcohol Use ²	540	517	511	529	561	585	588	606	573	568	621	606
Heavy Alcohol Use ²	346	320	318	322	352	373	358	349	352	355	382	367
FEMALE												
TOBACCO PRODUCTS¹	509	498	494	541	506	505	553	501	532	539	552	522
Cigarettes	497	481	488	537	497	487	548	494	524	529	538	511
Smokeless Tobacco	99	126	64	87	75	112	74	51	76	82	103	94
Cigars	110	121	119	109	128	112	113	127	150	140	145	161
Pipe Tobacco	90	35	41	40	45	28	54	35	53	45	60	33
ALCOHOL	640	593	662	635	641	673	667	679	634	677	682	717
Binge Alcohol Use ²	411	375	389	401	395	392	399	442	444	443	426	461
Heavy Alcohol Use ²	175	170	173	162	168	169	192	192	201	207	202	231

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2013.

Table 7.33D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Percentages, 2002-2013

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
TOTAL												
TOBACCO PRODUCTS¹	0.39	0.37	0.37	0.39	0.38	0.38	0.39	0.36	0.37	0.36	0.37	0.35
Cigarettes	0.38	0.36	0.36	0.35	0.36	0.36	0.36	0.35	0.34	0.35	0.35	0.34
Smokeless Tobacco	0.13	0.13	0.11	0.12	0.13	0.13	0.14	0.13	0.14	0.12	0.13	0.13
Cigars	0.16	0.16	0.15	0.16	0.16	0.15	0.16	0.16	0.16	0.16	0.16	0.15
Pipe Tobacco	0.08	0.06	0.07	0.07	0.07	0.08	0.06	0.07	0.06	0.06	0.08	0.06
ALCOHOL	0.46	0.43	0.44	0.44	0.43	0.45	0.43	0.42	0.42	0.43	0.43	0.44
Binge Alcohol Use ²	0.35	0.32	0.32	0.32	0.34	0.34	0.33	0.34	0.33	0.32	0.34	0.34
Heavy Alcohol Use ²	0.19	0.18	0.18	0.17	0.19	0.19	0.18	0.18	0.18	0.18	0.19	0.18
MALE												
TOBACCO PRODUCTS¹	0.56	0.55	0.55	0.55	0.55	0.55	0.58	0.54	0.56	0.52	0.54	0.54
Cigarettes	0.52	0.53	0.51	0.49	0.51	0.53	0.52	0.51	0.48	0.48	0.49	0.48
Smokeless Tobacco	0.26	0.25	0.22	0.23	0.25	0.23	0.27	0.26	0.27	0.23	0.25	0.24
Cigars	0.32	0.30	0.28	0.31	0.30	0.29	0.31	0.30	0.29	0.30	0.31	0.27
Pipe Tobacco	0.13	0.12	0.13	0.13	0.15	0.16	0.12	0.13	0.12	0.12	0.15	0.13
ALCOHOL	0.64	0.58	0.59	0.59	0.56	0.61	0.58	0.59	0.60	0.61	0.59	0.60
Binge Alcohol Use ²	0.53	0.51	0.49	0.50	0.53	0.55	0.54	0.55	0.52	0.51	0.55	0.53
Heavy Alcohol Use ²	0.34	0.31	0.31	0.31	0.33	0.35	0.33	0.32	0.32	0.32	0.34	0.32
FEMALE												
TOBACCO PRODUCTS¹	0.47	0.45	0.44	0.48	0.44	0.44	0.47	0.43	0.45	0.45	0.45	0.42
Cigarettes	0.45	0.44	0.44	0.48	0.44	0.42	0.47	0.42	0.44	0.44	0.44	0.42
Smokeless Tobacco	0.09	0.11	0.06	0.08	0.07	0.10	0.06	0.04	0.06	0.07	0.08	0.08
Cigars	0.10	0.11	0.11	0.10	0.11	0.10	0.10	0.11	0.13	0.12	0.12	0.13
Pipe Tobacco	0.08	0.03	0.04	0.04	0.04	0.02	0.05	0.03	0.04	0.04	0.05	0.03
ALCOHOL	0.59	0.54	0.59	0.56	0.56	0.58	0.57	0.58	0.54	0.56	0.56	0.58
Binge Alcohol Use ²	0.38	0.34	0.35	0.36	0.35	0.34	0.34	0.38	0.37	0.37	0.35	0.37
Heavy Alcohol Use ²	0.16	0.15	0.16	0.14	0.15	0.15	0.16	0.16	0.17	0.17	0.17	0.19

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2013.

Table 7.33P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: P Values from Tests of Differences of Percentages, 2013 Versus 2002-2012

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
TOTAL												
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0622	0.0073	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0007	0.1452	0.0604	
Smokeless Tobacco	0.8419	0.5345	0.0081	0.2118	0.7774	0.3551	0.5382	0.8705	0.5199	0.2909	0.4974	
Cigars	0.0161	0.0169	0.0001	0.0004	0.0006	0.0134	0.0279	0.0369	0.0361	0.4319	0.0460	
Pipe Tobacco	0.3646	0.0207	0.1971	0.6355	0.3848	0.6028	0.1554	0.4774	0.8491	0.5826	0.2429	
ALCOHOL	0.0163	0.0000	0.0002	0.4365	0.0139	0.0451	0.2316	0.3912	0.3465	0.4148	0.7921	
Binge Alcohol Use ²	0.5012	0.1922	0.2710	0.2685	0.7731	0.6054	0.4795	0.1647	0.8931	0.4582	0.9728	
Heavy Alcohol Use ²	0.1280	0.0835	0.0197	0.3287	0.0330	0.0469	0.0101	0.0584	0.1864	0.4792	0.3236	
MALE												
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0040	0.0007	0.1377	0.0058	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0322	0.0139	0.4133	0.1224	
Smokeless Tobacco	0.8831	0.3608	0.0159	0.2258	0.8660	0.4554	0.3538	0.5668	0.4155	0.3134	0.3852	
Cigars	0.0001	0.0030	0.0000	0.0000	0.0001	0.0006	0.0007	0.0110	0.0241	0.2911	0.0189	
Pipe Tobacco	0.3315	0.0778	0.4242	0.5251	0.2249	0.9913	0.1056	0.6411	0.6164	0.5310	0.4145	
ALCOHOL	0.8167	0.8566	0.6876	0.1999	0.7705	0.4347	0.5538	0.8378	0.9819	0.5718	0.3969	
Binge Alcohol Use ²	0.2390	0.4144	0.3437	0.7419	0.1543	0.0639	0.0524	0.0857	0.3919	0.6966	0.8027	
Heavy Alcohol Use ²	0.0061	0.0418	0.0101	0.0667	0.0079	0.0176	0.0078	0.0708	0.2025	0.6589	0.3404	
FEMALE												
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.0003	0.0003	0.0504	0.2040	0.2869	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0092	0.1658	0.2138	
Smokeless Tobacco	0.9902	0.5096	0.2287	0.6964	0.1158	0.4522	0.2919	0.0881	0.4335	0.8029	0.7004	
Cigars	0.0081	0.5329	0.1398	0.0650	0.7361	0.0805	0.0343	0.5667	0.9069	0.7149	0.8851	
Pipe Tobacco	0.9683	0.0171	0.0427	0.5433	0.2627	0.0052	0.8074	0.2296	0.4063	0.8055	0.1659	
ALCOHOL	0.0002	0.0000	0.0000	0.0149	0.0007	0.0317	0.0176	0.1435	0.1389	0.5280	0.6187	
Binge Alcohol Use ²	0.0091	0.0009	0.0017	0.0228	0.0207	0.0916	0.1264	0.9860	0.3342	0.4718	0.7027	
Heavy Alcohol Use ²	0.1043	0.9298	0.6883	0.1934	0.7876	0.8744	0.6106	0.5187	0.6760	0.5425	0.7047	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2013.