

Table 2.47A (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	174,105	174,409	83,514	83,384	69,825	69,663
GEOGRAPHIC DIVISION						
Northeast	31,783	32,738	14,890	14,881	12,131	12,139
New England	8,694	8,938	3,815	3,581	3,114	2,853
Middle Atlantic	23,089	23,800	11,076	11,300	9,017	9,285
Midwest	39,989	40,067	19,985	19,629	16,678	16,456
East North Central	27,455	27,888	13,719	13,625	11,496	11,301
West North Central	12,534	12,179	6,266	6,004	5,182	5,154
South	62,947	63,259	31,379	31,701	26,955	27,178
South Atlantic	33,409	32,885	15,785	15,702	13,474	13,484
East South Central	10,854	10,852	5,749	6,112	5,129	5,432
West South Central	18,684 ^a	19,522	9,846	9,887	8,351	8,262
West	39,385	38,344	17,260	17,173	14,061	13,891
Mountain	12,744	12,226	5,562	5,726	4,528	4,772
Pacific	26,641	26,118	11,698	11,447	9,534	9,119
COUNTY TYPE						
Large Metro	89,862	91,150	41,973	42,446	34,238	34,832
Small Metro	53,949	53,677	25,931	25,934	21,860	21,746
250K – 1 Mil. Pop.	35,031	35,166	16,749	16,805	13,947	14,098
< 250K Pop.	18,918	18,511	9,182	9,129	7,912	7,648
Nonmetro	30,293	29,582	15,610	15,003	13,728	13,085
Urbanized	13,025	12,700	6,562	6,020	5,680	5,229
Less Urbanized	14,036	14,200	7,562	7,664	6,747	6,683
Completely Rural	3,232	2,683	1,486	1,319	1,300	1,172

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.47B (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	69.1	68.8	33.2	32.9	27.7	27.5
GEOGRAPHIC DIVISION						
Northeast	68.5	70.4	32.1	32.0	26.2	26.1
New England	71.4	73.0	31.3	29.2	25.6	23.3
Middle Atlantic	67.5	69.4	32.4	33.0	26.4	27.1
Midwest	72.5	72.4	36.2	35.5	30.2	29.7
East North Central	71.4	72.4	35.7	35.4	29.9	29.3
West North Central	75.1	72.5	37.5	35.7	31.0	30.7
South	68.4	68.0	34.1	34.1	29.3	29.2
South Atlantic	68.8	67.1	32.5	32.0	27.7	27.5
East South Central	72.4	72.0	38.4	40.5	34.2	36.0
West South Central	65.6	67.6	34.6	34.2	29.3	28.6
West	67.7 ^a	65.2	29.6	29.2	24.2	23.6
Mountain	71.5 ^a	67.8	31.2	31.7	25.4	26.5
Pacific	66.0	64.1	29.0	28.1	23.6	22.4
COUNTY TYPE						
Large Metro	67.2	67.2	31.4	31.3	25.6	25.7
Small Metro	70.3	69.3	33.8	33.5	28.5	28.1
250K – 1 Mil. Pop.	69.2	68.3	33.1	32.6	27.5	27.4
< 250K Pop.	72.5	71.3	35.2	35.2	30.3	29.5
Nonmetro	73.4	73.1	37.8	37.1	33.3	32.3
Urbanized	72.2	74.4	36.4	35.3	31.5	30.6
Less Urbanized	73.8	72.3	39.8	39.0	35.5	34.0
Completely Rural	77.2	72.0	35.5	35.4	31.1	31.5

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.48A (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	6,668 ^b	6,112	4,799 ^b	4,419	2,894 ^b	2,612
GEOGRAPHIC DIVISION						
Northeast	1,068	1,017	810	765	486	433
New England	286	281	220	223	126	118
Middle Atlantic	782	737	590	542	360	315
Midwest	1,515 ^b	1,338	1,133 ^b	997	720 ^b	615
East North Central	1,054 ^b	908	795 ^b	669	502 ^b	394
West North Central	462	431	338	328	218	221
South	2,591 ^a	2,393	1,773	1,711	1,057	990
South Atlantic	1,262	1,194	875	856	525	497
East South Central	482	443	326	294	206	182
West South Central	847	757	573	561	327	310
West	1,494	1,363	1,083	946	631	575
Mountain	481	461	334	325	204	218
Pacific	1,013	902	749 ^a	621	427	357
COUNTY TYPE						
Large Metro	3,258 ^b	2,913	2,317	2,137	1,364	1,226
Small Metro	2,107	2,030	1,559	1,433	953 ^a	841
250K – 1 Mil. Pop.	1,368	1,363	1,009	974	622	558
< 250K Pop.	739	666	550	459	331	283
Nonmetro	1,302 ^a	1,169	923	849	576	545
Urbanized	544 ^a	428	399 ^b	303	246 ^b	175
Less Urbanized	613	647	424	472	265	316
Completely Rural	146 ^a	94	99	74	65	54

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.48B (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	27.1 ^b	25.1	19.5 ^b	18.1	11.8 ^b	10.7
GEOGRAPHIC DIVISION						
Northeast	24.8	24.0	18.8	18.1	11.3	10.2
New England	25.7	25.7	19.7	20.4	11.3	10.8
Middle Atlantic	24.5	23.4	18.5	17.3	11.3	10.0
Midwest	28.0 ^b	25.1	20.9 ^b	18.7	13.3 ^a	11.5
East North Central	27.8 ^b	24.3	21.0 ^b	17.9	13.2 ^b	10.5
West North Central	28.5	27.1	20.9	20.6	13.5	13.9
South	28.8 ^a	26.7	19.7	19.1	11.7	11.0
South Atlantic	27.8	26.6	19.3	19.1	11.6	11.1
East South Central	33.1	30.6	22.4	20.3	14.1	12.6
West South Central	28.1	25.0	19.0	18.6	10.8	10.3
West	25.4	23.4	18.4	16.3	10.7	9.9
Mountain	26.7	25.8	18.5	18.2	11.3	12.2
Pacific	24.8	22.4	18.4	15.4	10.5	8.9
COUNTY TYPE						
Large Metro	25.0 ^b	22.3	17.8	16.4	10.5	9.4
Small Metro	27.5	27.0	20.3	19.0	12.4 ^a	11.2
250K – 1 Mil. Pop.	26.7	27.1	19.7	19.3	12.2	11.1
< 250K Pop.	28.9	26.8	21.5 ^a	18.5	13.0	11.4
Nonmetro	33.4	30.9	23.6	22.4	14.8	14.4
Urbanized	31.7	28.1	23.3	19.9	14.3 ^a	11.4
Less Urbanized	34.1	33.3	23.6	24.3	14.7	16.3
Completely Rural	37.6 ^a	29.6	25.5	23.3	16.9	17.1

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.49A (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	23,383	23,437	17,601	17,484	13,986	13,934
GEOGRAPHIC DIVISION						
Northeast	4,365	4,198	3,271	3,100	2,572	2,450
New England	1,206	1,171	909	866	710	699
Middle Atlantic	3,159	3,027	2,363	2,233	1,862	1,751
Midwest	5,356	5,456	4,152	4,217	3,362	3,465
East North Central	3,732	3,701	2,911	2,862	2,353	2,368
West North Central	1,624 ^b	1,756	1,241 ^a	1,355	1,008	1,098
South	8,219	8,445	6,214	6,374	5,032	5,130
South Atlantic	4,144	4,251	3,105	3,151	2,464	2,520
East South Central	1,320	1,392	1,032	1,092	920	925
West South Central	2,755	2,802	2,077	2,131	1,648	1,685
West	5,443	5,338	3,964	3,793	3,020	2,889
Mountain	1,643	1,708	1,209	1,202	927	911
Pacific	3,800	3,630	2,755	2,591	2,092	1,978
COUNTY TYPE						
Large Metro	11,951	12,043	8,865	8,806	6,902	6,870
Small Metro	7,508	7,800	5,656	5,867	4,479	4,663
250K – 1 Mil. Pop.	4,677	4,868	3,490	3,598	2,717	2,868
< 250K Pop.	2,831	2,932	2,166	2,269	1,762	1,795
Nonmetro	3,924	3,594	3,080	2,811	2,605	2,401
Urbanized	1,912	1,627	1,478	1,282	1,252	1,088
Less Urbanized	1,762	1,700	1,400	1,333	1,181	1,136
Completely Rural	251	267	202	196	171	177

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.49B (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	69.6	68.8	52.4	51.3	41.6	40.9
GEOGRAPHIC DIVISION						
Northeast	71.3	69.4	53.4	51.3	42.0	40.5
New England	75.5	72.1	56.9	53.3	44.5	43.0
Middle Atlantic	69.8	68.5	52.2	50.5	41.2	39.6
Midwest	73.0	72.8	56.6	56.3	45.8	46.2
East North Central	73.5	72.0	57.3	55.7	46.4	46.1
West North Central	71.8	74.5	54.9	57.5	44.6	46.6
South	68.4	68.0	51.7	51.3	41.9	41.3
South Atlantic	67.8	67.3	50.8	49.9	40.3	39.9
East South Central	69.4	70.0	54.3	54.9	48.4	46.5
West South Central	68.7	68.1	51.8	51.8	41.1	40.9
West	67.2	65.8	48.9	46.8	37.3	35.6
Mountain	67.9	68.4	50.0	48.1	38.3	36.5
Pacific	66.9	64.7	48.5	46.1	36.8	35.2
COUNTY TYPE						
Large Metro	67.0	66.3	49.7	48.5	38.7	37.8
Small Metro	71.6	70.1	53.9	52.8	42.7	41.9
250K – 1 Mil. Pop.	69.9	69.3	52.2	51.2	40.6	40.8
< 250K Pop.	74.4	71.6	56.9	55.4	46.3	43.8
Nonmetro	74.9	75.0	58.8	58.7	49.7	50.1
Urbanized	76.0	77.1	58.8	60.7	49.8	51.5
Less Urbanized	73.3	72.9	58.3	57.2	49.2	48.7
Completely Rural	77.2	76.6	62.1	56.3	52.8	50.7

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.50A (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	144,054	144,860	61,115	61,482	52,946	53,117
GEOGRAPHIC DIVISION						
Northeast	26,350 ^a	27,523	10,809	11,016	9,073	9,256
New England	7,202	7,486	2,686	2,492	2,278	2,036
Middle Atlantic	19,148 ^a	20,036	8,123	8,524	6,795	7,220
Midwest	33,118	33,272	14,700	14,415	12,596	12,375
East North Central	22,670	23,280	10,013	10,094	8,640	8,540
West North Central	10,448	9,993	4,687	4,322	3,956	3,836
South	52,137	52,421	23,392	23,616	20,866	21,058
South Atlantic	28,002	27,441	11,805	11,695	10,485	10,466
East South Central	9,052	9,017	4,391	4,726	4,004	4,324
West South Central	15,083 ^a	15,963	7,196	7,195	6,377	6,268
West	32,448	31,643	12,213	12,434	10,411	10,428
Mountain	10,620 ^a	10,058	4,019	4,199	3,397	3,644
Pacific	21,828	21,585	8,194	8,235	7,014	6,784
COUNTY TYPE						
Large Metro	74,653	76,194	30,790	31,504	25,972	26,736
Small Metro	44,334	43,847	18,716	18,635	16,428	16,242
250K – 1 Mil. Pop.	28,986	28,934	12,250	12,233	10,609	10,671
< 250K Pop.	15,348	14,913	6,466	6,401	5,819	5,571
Nonmetro	25,066	24,819	11,608	11,343	10,547	10,138
Urbanized	10,570	10,644	4,685	4,436	4,182	3,966
Less Urbanized	11,662	11,854	5,737	5,859	5,302	5,231
Completely Rural	2,835	2,321	1,186	1,049	1,064	941

^aLow precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.50B (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	74.4	74.2	31.6	31.5	27.3	27.2
GEOGRAPHIC DIVISION						
Northeast	73.3	75.9	30.1	30.4	25.2	25.5
New England	76.1	78.6	28.4	26.2	24.1	21.4
Middle Atlantic	72.3	75.0	30.7	31.9	25.7	27.0
Midwest	78.1	78.3	34.7	33.9	29.7	29.1
East North Central	76.6	78.5	33.8	34.0	29.2	28.8
West North Central	81.5 ^a	77.7	36.6	33.6	30.9	29.8
South	73.4	73.2	32.9	33.0	29.4	29.4
South Atlantic	73.9	71.8	31.1	30.6	27.7	27.4
East South Central	77.8	77.4	37.8	40.6	34.4	37.1
West South Central	70.3	73.4	33.5	33.1	29.7	28.8
West	73.4 ^a	70.6	27.6	27.7	23.5	23.3
Mountain	78.1 ^a	73.1	29.6	30.5	25.0	26.5
Pacific	71.3	69.4	26.7	26.5	22.9	21.8
COUNTY TYPE						
Large Metro	72.5	72.9	29.9	30.1	25.2	25.6
Small Metro	75.7	74.6	31.9	31.7	28.0	27.6
250K – 1 Mil. Pop.	74.6	73.4	31.5	31.0	27.3	27.1
< 250K Pop.	77.7	76.9	32.7	33.0	29.5	28.7
Nonmetro	78.1	77.9	36.2	35.6	32.8	31.8
Urbanized	76.5	79.2	33.9	33.0	30.3	29.5
Less Urbanized	78.7	77.1	38.7	38.1	35.8	34.0
Completely Rural	81.6	75.9	34.1	34.3	30.6	30.8

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.51A (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	167,437	168,297	78,716	78,965	66,932	67,051
GEOGRAPHIC DIVISION						
Northeast	30,716 ^a	31,721	14,080	14,116	11,646	11,706
New England	8,409	8,658	3,595	3,358	2,988	2,735
Middle Atlantic	22,307	23,063	10,485	10,757	8,657	8,971
Midwest	38,474	38,729	18,852	18,633	15,958	15,841
East North Central	26,402	26,981	12,924	12,956	10,994	10,907
West North Central	12,072	11,748	5,928	5,677	4,964	4,934
South	60,356	60,866	29,606	29,990	25,898	26,188
South Atlantic	32,146	31,692	14,910	14,845	12,949	12,986
East South Central	10,372	10,409	5,423	5,819	4,924	5,249
West South Central	17,838 ^a	18,766	9,273	9,326	8,025	7,952
West	37,891	36,981	16,177	16,227	13,431	13,316
Mountain	12,263	11,766	5,228	5,401	4,324	4,555
Pacific	25,628	25,216	10,949	10,826	9,106	8,762
COUNTY TYPE						
Large Metro	86,604	88,236	39,656	40,310	32,873	33,606
Small Metro	51,842	51,647	24,372	24,502	20,907	20,905
250K – 1 Mil. Pop.	33,663	33,802	15,740	15,831	13,325	13,539
< 250K Pop.	18,179	17,845	8,632	8,670	7,581	7,366
Nonmetro	28,991	28,413	14,688	14,154	13,152	12,539
Urbanized	12,482	12,271	6,163	5,717	5,434	5,055
Less Urbanized	13,423	13,553	7,137	7,192	6,482	6,367
Completely Rural	3,086	2,588	1,387	1,245	1,235	1,118

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.51B (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	73.7	73.4	34.6	34.4	29.5	29.2
GEOGRAPHIC DIVISION						
Northeast	73.0	75.0	33.5	33.4	27.7	27.7
New England	76.0	77.6	32.5	30.1	27.0	24.5
Middle Atlantic	71.9	74.1	33.8	34.5	27.9	28.8
Midwest	77.3	77.4	37.9	37.3	32.1	31.7
East North Central	76.1	77.5	37.3	37.2	31.7	31.3
West North Central	80.1	77.2	39.3	37.3	32.9	32.4
South	72.7	72.5	35.7	35.7	31.2	31.2
South Atlantic	73.0	71.2	33.9	33.4	29.4	29.2
East South Central	76.7	76.3	40.1	42.7	36.4	38.5
West South Central	70.0	72.6	36.4	36.1	31.5	30.8
West	72.4 ^a	69.8	30.9	30.6	25.7	25.1
Mountain	76.6 ^a	72.4	32.6	33.2	27.0	28.0
Pacific	70.6	68.7	30.2	29.5	25.1	23.9
COUNTY TYPE						
Large Metro	71.7	71.9	32.8	32.9	27.2	27.4
Small Metro	75.1	73.9	35.3	35.0	30.3	29.9
250K – 1 Mil. Pop.	74.0	72.8	34.6	34.1	29.3	29.1
< 250K Pop.	77.2	76.0	36.6	36.9	32.2	31.4
Nonmetro	77.6	77.5	39.3	38.6	35.2	34.2
Urbanized	76.4	78.9	37.7	36.8	33.3	32.5
Less Urbanized	77.9	76.5	41.4	40.6	37.6	36.0
Completely Rural	81.2	76.0	36.5	36.5	32.5	32.8

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.