

Table 2.47C (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	888	874	865	920	831	852
GEOGRAPHIC DIVISION						
Northeast	373	337	357	332	315	309
New England	172	169	159	153	151	139
Middle Atlantic	331	288	322	292	277	275
Midwest	308	317	350	381	345	369
East North Central	260	266	291	303	286	287
West North Central	162	169	202	217	196	218
South	573	563	584	623	562	589
South Atlantic	464	441	460	461	444	439
East South Central	186	215	210	227	210	217
West South Central	277	282	304	291	288	287
West	476	471	421	453	404	409
Mountain	236	196	221	222	199	208
Pacific	412	427	359	396	353	355
COUNTY TYPE						
Large Metro	1,649	1,618	865	892	777	787
Small Metro	1,130	1,235	681	697	625	618
250K – 1 Mil. Pop.	963	995	571	532	512	471
< 250K Pop.	749	747	409	426	377	378
Nonmetro	879	898	525	550	486	496
Urbanized	605	716	370	356	336	324
Less Urbanized	606	650	376	411	346	368
Completely Rural	410	329	185	170	173	153

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.47D (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.35	0.34	0.34	0.36	0.33	0.34
GEOGRAPHIC DIVISION						
Northeast	0.80	0.73	0.77	0.71	0.68	0.66
New England	1.41	1.38	1.30	1.25	1.24	1.13
Middle Atlantic	0.97	0.84	0.94	0.85	0.81	0.80
Midwest	0.56	0.57	0.64	0.69	0.63	0.67
East North Central	0.68	0.69	0.76	0.79	0.74	0.75
West North Central	0.97	1.00	1.21	1.29	1.18	1.30
South	0.62	0.61	0.63	0.67	0.61	0.63
South Atlantic	0.96	0.90	0.95	0.94	0.91	0.90
East South Central	1.24	1.42	1.40	1.51	1.40	1.44
West South Central	0.97	0.98	1.07	1.01	1.01	0.99
West	0.82	0.80	0.72	0.77	0.69	0.70
Mountain	1.33	1.09	1.24	1.23	1.11	1.16
Pacific	1.02	1.05	0.89	0.97	0.87	0.87
COUNTY TYPE						
Large Metro	0.53	0.50	0.47	0.50	0.45	0.47
Small Metro	0.58	0.61	0.65	0.65	0.63	0.61
250K – 1 Mil. Pop.	0.72	0.74	0.78	0.72	0.76	0.70
< 250K Pop.	1.01	1.07	1.13	1.21	1.12	1.11
Nonmetro	0.74	0.76	0.84	0.92	0.80	0.87
Urbanized	1.13	1.20	1.26	1.42	1.23	1.36
Less Urbanized	0.99	1.11	1.27	1.29	1.19	1.23
Completely Rural	2.48	2.39	2.13	2.50	2.26	2.33

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.47P (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.4403		0.5490		0.5631	
GEOGRAPHIC DIVISION						
Northeast	0.0957		0.9023		0.9391	
New England	0.4225		0.1800		0.1001	
Middle Atlantic	0.1432		0.6457		0.5170	
Midwest	0.9072		0.4185		0.5939	
East North Central	0.3011		0.7736		0.5874	
West North Central	0.0767		0.3218		0.8447	
South	0.6792		0.9899		0.9540	
South Atlantic	0.1750		0.7208		0.8589	
East South Central	0.7997		0.3007		0.3717	
West South Central	0.1368		0.8028		0.5715	
West	0.0286		0.6853		0.5957	
Mountain	0.0364		0.7668		0.5152	
Pacific	0.1834		0.5110		0.3225	
COUNTY TYPE						
Large Metro	0.9880		0.8936		0.8971	
Small Metro	0.2127		0.7316		0.6342	
250K – 1 Mil. Pop.	0.3734		0.6731		0.8693	
< 250K Pop.	0.3882		0.9949		0.5870	
Nonmetro	0.7701		0.5308		0.4192	
Urbanized	0.1800		0.5172		0.6198	
Less Urbanized	0.2653		0.6749		0.3885	
Completely Rural	0.1640		0.9785		0.8989	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.48C (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	102	101	93	89	71	69
GEOGRAPHIC DIVISION						
Northeast	38	36	36	32	27	25
New England	17	15	16	14	13	12
Middle Atlantic	34	32	32	29	24	22
Midwest	38	36	35	33	29	27
East North Central	31	30	28	27	24	21
West North Central	21	21	19	19	17	17
South	64	68	55	62	44	48
South Atlantic	47	52	42	48	34	39
East South Central	23	21	20	19	16	14
West South Central	35	35	30	32	22	23
West	54	53	51	48	36	34
Mountain	25	26	23	23	16	18
Pacific	48	46	46	42	32	29
COUNTY TYPE						
Large Metro	93	94	80	80	56	57
Small Metro	74	70	63	53	44	38
250K – 1 Mil. Pop.	59	58	48	45	36	32
< 250K Pop.	47	40	41	31	27	22
Nonmetro	55	55	43	46	33	36
Urbanized	39	35	31	27	23	21
Less Urbanized	36	45	27	37	21	29
Completely Rural	21	12	16	11	11	9

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.48D (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.42	0.42	0.38	0.37	0.29	0.28
GEOGRAPHIC DIVISION						
Northeast	0.89	0.84	0.84	0.77	0.64	0.60
New England	1.55	1.39	1.43	1.29	1.19	1.06
Middle Atlantic	1.07	1.02	1.01	0.92	0.75	0.71
Midwest	0.70	0.68	0.64	0.61	0.54	0.50
East North Central	0.81	0.79	0.75	0.72	0.63	0.56
West North Central	1.32	1.34	1.17	1.21	1.02	1.08
South	0.71	0.76	0.61	0.69	0.49	0.53
South Atlantic	1.03	1.17	0.92	1.08	0.74	0.88
East South Central	1.59	1.43	1.39	1.31	1.10	0.97
West South Central	1.18	1.16	0.98	1.05	0.73	0.75
West	0.91	0.90	0.87	0.82	0.61	0.59
Mountain	1.38	1.47	1.26	1.27	0.91	1.00
Pacific	1.18	1.13	1.12	1.05	0.79	0.72
COUNTY TYPE						
Large Metro	0.59	0.59	0.54	0.53	0.40	0.40
Small Metro	0.68	0.67	0.63	0.57	0.50	0.45
250K – 1 Mil. Pop.	0.79	0.82	0.75	0.72	0.59	0.57
< 250K Pop.	1.30	1.17	1.15	1.01	0.89	0.81
Nonmetro	0.95	1.02	0.83	0.96	0.69	0.81
Urbanized	1.48	1.48	1.28	1.32	1.06	1.03
Less Urbanized	1.31	1.54	1.18	1.52	0.98	1.32
Completely Rural	2.81	3.07	2.45	2.86	1.90	2.57

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.48P (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.0005		0.0088		0.0069	
GEOGRAPHIC DIVISION						
Northeast	0.5158		0.4980		0.2031	
New England	0.9905		0.7415		0.7435	
Middle Atlantic	0.4587		0.3536		0.2025	
Midwest	0.0017		0.0084		0.0110	
East North Central	0.0010		0.0017		0.0007	
West North Central	0.4064		0.8712		0.8007	
South	0.0412		0.5167		0.3106	
South Atlantic	0.4164		0.8810		0.6461	
East South Central	0.2650		0.2979		0.2989	
West South Central	0.0552		0.7400		0.5661	
West	0.1455		0.0795		0.3191	
Mountain	0.6750		0.8613		0.5274	
Pacific	0.1528		0.0597		0.1312	
COUNTY TYPE						
Large Metro	0.0016		0.0648		0.0562	
Small Metro	0.6057		0.1187		0.0492	
250K – 1 Mil. Pop.	0.7593		0.7066		0.1789	
< 250K Pop.	0.1946		0.0345		0.1435	
Nonmetro	0.0563		0.3300		0.7218	
Urbanized	0.0725		0.0534		0.0362	
Less Urbanized	0.6835		0.6948		0.3042	
Completely Rural	0.0487		0.5513		0.9486	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.49C (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	153	159	168	171	169	166
GEOGRAPHIC DIVISION						
Northeast	57	65	68	70	65	66
New England	25	30	33	35	31	32
Middle Atlantic	51	57	59	60	57	58
Midwest	58	58	62	63	66	65
East North Central	45	48	49	51	53	53
West North Central	34	31	36	35	38	38
South	91	105	95	108	105	106
South Atlantic	75	84	76	85	86	83
East South Central	33	37	35	36	36	37
West South Central	44	50	49	57	52	55
West	90	86	96	96	87	90
Mountain	44	38	40	41	38	40
Pacific	78	76	88	87	79	81
COUNTY TYPE						
Large Metro	245	282	198	227	162	192
Small Metro	212	242	176	194	146	154
250K – 1 Mil. Pop.	161	173	129	137	105	108
< 250K Pop.	144	170	122	138	101	111
Nonmetro	215	137	171	118	157	104
Urbanized	193	98	148	82	137	73
Less Urbanized	103	100	91	88	79	76
Completely Rural	33	37	28	29	24	27

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.49D (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.45	0.47	0.50	0.50	0.50	0.49
GEOGRAPHIC DIVISION						
Northeast	0.93	1.07	1.11	1.16	1.07	1.10
New England	1.55	1.86	2.07	2.15	1.93	1.97
Middle Atlantic	1.13	1.29	1.31	1.37	1.27	1.31
Midwest	0.78	0.78	0.84	0.84	0.90	0.87
East North Central	0.90	0.94	0.96	1.00	1.05	1.03
West North Central	1.52	1.33	1.60	1.49	1.69	1.60
South	0.76	0.85	0.79	0.87	0.87	0.85
South Atlantic	1.22	1.33	1.24	1.35	1.40	1.31
East South Central	1.76	1.87	1.87	1.81	1.89	1.88
West South Central	1.11	1.22	1.22	1.38	1.29	1.33
West	1.11	1.06	1.19	1.18	1.08	1.11
Mountain	1.80	1.51	1.67	1.62	1.55	1.61
Pacific	1.38	1.36	1.54	1.55	1.39	1.45
COUNTY TYPE						
Large Metro	0.66	0.69	0.70	0.72	0.67	0.69
Small Metro	0.73	0.82	0.86	0.90	0.89	0.88
250K – 1 Mil. Pop.	0.88	0.93	1.02	1.11	1.04	1.11
< 250K Pop.	1.28	1.46	1.42	1.50	1.44	1.46
Nonmetro	0.88	0.89	0.98	1.05	1.14	1.10
Urbanized	1.23	1.36	1.30	1.51	1.57	1.66
Less Urbanized	1.40	1.29	1.62	1.61	1.80	1.59
Completely Rural	3.13	3.90	3.68	3.29	3.87	3.48

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.49P (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.1733		0.0991		0.2635	
GEOGRAPHIC DIVISION						
Northeast	0.1861		0.1646		0.3123	
New England	0.1471		0.1729		0.5576	
Middle Atlantic	0.4272		0.3667		0.3878	
Midwest	0.8404		0.7714		0.7316	
East North Central	0.2358		0.2203		0.8388	
West North Central	0.1394		0.2032		0.3623	
South	0.7391		0.7462		0.6331	
South Atlantic	0.7830		0.6089		0.8251	
East South Central	0.8245		0.7954		0.4964	
West South Central	0.6746		0.9807		0.9113	
West	0.3335		0.1598		0.2724	
Mountain	0.8213		0.3604		0.3718	
Pacific	0.2171		0.2542		0.4194	
COUNTY TYPE						
Large Metro	0.4834		0.2189		0.3826	
Small Metro	0.1664		0.3190		0.5073	
250K – 1 Mil. Pop.	0.6095		0.4962		0.8850	
< 250K Pop.	0.1055		0.4085		0.1691	
Nonmetro	0.9192		0.9432		0.7916	
Urbanized	0.5753		0.3648		0.4439	
Less Urbanized	0.8233		0.6156		0.8445	
Completely Rural	0.9026		0.2253		0.6895	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.50C (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	833	818	804	870	784	813
GEOGRAPHIC DIVISION						
Northeast	361	313	324	316	292	297
New England	167	148	140	138	137	126
Middle Atlantic	320	272	293	281	258	267
Midwest	287	298	326	362	324	350
East North Central	246	247	266	288	264	274
West North Central	145	162	193	204	190	205
South	554	539	560	594	540	566
South Atlantic	450	424	432	438	423	419
East South Central	181	205	207	218	201	209
West South Central	267	271	296	275	279	272
West	441	431	402	425	388	386
Mountain	199	182	204	204	184	196
Pacific	390	392	345	376	341	336
COUNTY TYPE						
Large Metro	1,569	1,522	792	813	717	731
Small Metro	1,033	1,132	596	607	558	550
250K – 1 Mil. Pop.	880	922	505	472	466	425
< 250K Pop.	678	675	347	367	327	332
Nonmetro	796	827	454	492	430	448
Urbanized	523	652	306	313	284	289
Less Urbanized	548	591	326	357	310	327
Completely Rural	379	304	165	151	157	134

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.50D (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.43	0.42	0.42	0.45	0.40	0.42
GEOGRAPHIC DIVISION						
Northeast	1.00	0.86	0.90	0.87	0.81	0.82
New England	1.77	1.56	1.48	1.45	1.45	1.32
Middle Atlantic	1.21	1.02	1.11	1.05	0.97	1.00
Midwest	0.68	0.70	0.77	0.85	0.76	0.82
East North Central	0.83	0.83	0.90	0.97	0.89	0.92
West North Central	1.13	1.26	1.51	1.59	1.48	1.59
South	0.78	0.75	0.79	0.83	0.76	0.79
South Atlantic	1.19	1.11	1.14	1.15	1.11	1.10
East South Central	1.55	1.76	1.78	1.87	1.73	1.79
West South Central	1.24	1.25	1.38	1.26	1.30	1.25
West	1.00	0.96	0.91	0.95	0.88	0.86
Mountain	1.46	1.32	1.50	1.48	1.35	1.43
Pacific	1.27	1.26	1.13	1.21	1.11	1.08
COUNTY TYPE						
Large Metro	0.64	0.60	0.57	0.62	0.55	0.58
Small Metro	0.71	0.73	0.78	0.78	0.76	0.74
250K – 1 Mil. Pop.	0.87	0.90	0.96	0.88	0.94	0.85
< 250K Pop.	1.25	1.32	1.39	1.49	1.37	1.37
Nonmetro	0.94	0.96	1.01	1.12	0.99	1.07
Urbanized	1.47	1.47	1.54	1.72	1.50	1.66
Less Urbanized	1.23	1.41	1.53	1.58	1.47	1.54
Completely Rural	2.90	2.97	2.57	2.96	2.66	2.72

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.50P (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.7515		0.9113		0.8142	
GEOGRAPHIC DIVISION						
Northeast	0.0507		0.7867		0.7847	
New England	0.2910		0.2197		0.1039	
Middle Atlantic	0.0962		0.4159		0.3186	
Midwest	0.8490		0.5220		0.6155	
East North Central	0.1035		0.8725		0.7568	
West North Central	0.0363		0.1959		0.6634	
South	0.8553		0.9626		0.9733	
South Atlantic	0.1998		0.7411		0.8708	
East South Central	0.8525		0.2917		0.2849	
West South Central	0.0650		0.7990		0.5817	
West	0.0374		0.9309		0.8174	
Mountain	0.0163		0.6541		0.4408	
Pacific	0.2823		0.8737		0.4884	
COUNTY TYPE						
Large Metro	0.6569		0.7806		0.6520	
Small Metro	0.2538		0.8082		0.6863	
250K – 1 Mil. Pop.	0.2968		0.6828		0.8349	
< 250K Pop.	0.6509		0.8866		0.7129	
Nonmetro	0.8743		0.6979		0.4607	
Urbanized	0.1878		0.6635		0.7175	
Less Urbanized	0.3415		0.7795		0.3997	
Completely Rural	0.2056		0.9682		0.9663	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.51C (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	856	844	851	906	822	840
GEOGRAPHIC DIVISION						
Northeast	370	324	354	329	312	309
New England	170	157	156	153	148	140
Middle Atlantic	329	279	319	289	274	275
Midwest	297	306	345	379	340	367
East North Central	253	254	284	301	280	283
West North Central	152	168	200	216	194	220
South	567	551	576	618	558	583
South Atlantic	460	432	454	454	441	432
East South Central	184	210	210	225	208	216
West South Central	274	276	302	288	285	284
West	450	451	418	446	404	402
Mountain	210	185	214	219	194	207
Pacific	397	411	360	390	354	348
COUNTY TYPE						
Large Metro	1,631	1,595	851	874	767	775
Small Metro	1,107	1,212	664	682	614	609
250K – 1 Mil. Pop.	943	975	559	520	506	464
< 250K Pop.	732	731	394	416	369	372
Nonmetro	859	873	513	533	479	486
Urbanized	588	696	360	345	331	318
Less Urbanized	591	630	366	395	341	360
Completely Rural	398	323	177	164	167	148

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.51D (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.38	0.37	0.37	0.40	0.36	0.37
GEOGRAPHIC DIVISION						
Northeast	0.88	0.77	0.84	0.78	0.74	0.73
New England	1.54	1.41	1.41	1.38	1.34	1.25
Middle Atlantic	1.06	0.90	1.03	0.93	0.88	0.88
Midwest	0.60	0.61	0.69	0.76	0.68	0.73
East North Central	0.73	0.73	0.82	0.87	0.81	0.81
West North Central	1.01	1.11	1.33	1.42	1.29	1.44
South	0.68	0.66	0.69	0.74	0.67	0.69
South Atlantic	1.04	0.97	1.03	1.02	1.00	0.97
East South Central	1.36	1.54	1.55	1.65	1.54	1.58
West South Central	1.08	1.07	1.18	1.12	1.12	1.10
West	0.86	0.85	0.80	0.84	0.77	0.76
Mountain	1.31	1.14	1.33	1.35	1.21	1.27
Pacific	1.09	1.12	0.99	1.06	0.98	0.95
COUNTY TYPE						
Large Metro	0.56	0.53	0.51	0.54	0.49	0.51
Small Metro	0.62	0.65	0.71	0.71	0.69	0.67
250K – 1 Mil. Pop.	0.76	0.79	0.86	0.80	0.83	0.77
< 250K Pop.	1.07	1.14	1.26	1.32	1.24	1.21
Nonmetro	0.82	0.84	0.92	1.00	0.89	0.96
Urbanized	1.26	1.30	1.38	1.54	1.34	1.49
Less Urbanized	1.08	1.22	1.39	1.42	1.32	1.37
Completely Rural	2.65	2.67	2.32	2.69	2.43	2.50

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.51P (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.5751		0.6980		0.6664	
GEOGRAPHIC DIVISION						
Northeast	0.0942		0.9345		1.0000	
New England	0.4320		0.1528		0.0949	
Middle Atlantic	0.1378		0.5987		0.4712	
Midwest	0.8925		0.5422		0.7012	
East North Central	0.1732		0.9774		0.7524	
West North Central	0.0706		0.3188		0.8125	
South	0.7973		0.9601		0.9888	
South Atlantic	0.1838		0.7137		0.8653	
East South Central	0.8706		0.2664		0.3428	
West South Central	0.0847		0.8204		0.5925	
West	0.0273		0.8154		0.6285	
Mountain	0.0225		0.7567		0.5523	
Pacific	0.2019		0.6472		0.3706	
COUNTY TYPE						
Large Metro	0.7627		0.9697		0.7929	
Small Metro	0.1577		0.7979		0.6894	
250K – 1 Mil. Pop.	0.2596		0.6614		0.9067	
< 250K Pop.	0.4138		0.8725		0.6356	
Nonmetro	0.9080		0.5798		0.4255	
Urbanized	0.1647		0.5983		0.6818	
Less Urbanized	0.3388		0.6725		0.3712	
Completely Rural	0.1993		0.9958		0.9300	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.