

MIDWEST REGION

Table 5 Selected Drug Use, Perceptions of Great Risk, Average Annual Marijuana Initiates, Past Year Substance Dependence or Abuse, Needing But Not Receiving Treatment, and Past Year Mental Health Measures in the Midwest Region, by Age Group: Estimated Numbers (in Thousands), Annual Averages Based on 2009-2010 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLICIT DRUGS					
Past Month Illicit Drug Use ¹	4,477	524	1,509	2,445	3,953
Past Year Marijuana Use	5,963	700	2,200	3,062	5,262
Past Month Marijuana Use	3,505	383	1,302	1,820	3,122
Past Month Use of Illicit Drugs Other Than Marijuana ¹	1,832	241	593	998	1,591
Past Year Cocaine Use	853	50	317	486	803
Past Year Nonmedical Pain Reliever Use	2,656	353	889	1,413	2,303
Perception of Great Risk of Smoking Marijuana Once a Month	17,656	1,670	1,297	14,689	15,985
Average Annual Number of Marijuana Initiates ²	591	273	285	32	317
ALCOHOL					
Past Month Alcohol Use	30,202	763	4,818	24,622	29,439
Past Month Binge Alcohol Use ³	14,112	468	3,329	10,316	13,644
Perception of Great Risk of Drinking Five or More Drinks Once or Twice a Week	21,534	2,045	2,189	17,299	19,488
Past Month Alcohol Use (Persons Aged 12 to 20)	2,286 ⁴	--	--	--	--
Past Month Binge Alcohol Use (Persons Aged 12 to 20) ³	1,580 ⁴	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁵	16,632	675	3,396	12,561	15,957
Past Month Cigarette Use	13,879	530	2,880	10,468	13,348
Perception of Great Risk of Smoking One or More Packs of Cigarettes per Day	37,252	3,457	4,529	29,266	33,795
PAST YEAR DEPENDENCE, ABUSE, AND TREATMENT⁶					
Illicit Drug Dependence ¹	1,021	127	381	512	893
Illicit Drug Dependence or Abuse ¹	1,480	237	558	685	1,243
Alcohol Dependence	1,867	94	520	1,253	1,772
Alcohol Dependence or Abuse	4,121	246	1,247	2,628	3,874
Alcohol or Illicit Drug Dependence or Abuse ¹	4,964	384	1,526	3,054	4,579
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,7}	1,322	220	512	590	1,102
Needing But Not Receiving Treatment for Alcohol Use ⁷	3,856	235	1,180	2,442	3,622
PAST YEAR MENTAL HEALTH					
Had at Least One Major Depressive Episode ⁸	--	441	632	2,855	3,486
Serious Mental Illness ⁹	--	--	284	1,817	2,100
Any Mental Illness ¹⁰	--	--	1,392	7,880	9,272
Had Serious Thoughts of Suicide	--	--	499	1,520	2,019

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

NOTE: Estimates for any mental illness (AMI) and serious mental illness (SMI) presented in this table may differ from previously published AMI and SMI estimates because of revised estimation procedures. For details, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates include data from original methamphetamine questions but do not include new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Average annual number of marijuana initiates = $X_1 \div 2$, where X_1 is the number of marijuana initiates in the past 24 months.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

⁴ Underage drinking is defined for persons aged 12 to 20; therefore, the "Total" estimate reflects that age group and not persons aged 12 or older.

⁵ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁶ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁷ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs (or alcohol), but not receiving treatment for an illicit drug (or alcohol) problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

⁸ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from persons aged 18 or older to produce an estimate for those aged 12 or older. For more details, see Section A.11 in Appendix A of the report on *State Estimates of Substance Use and Mental Disorders from the 2009-2010 NSDUHs*.

⁹ Serious mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in serious functional impairment.

¹⁰ Any mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012; SMI and AMI Estimates Revised October 2013).

MIDWEST REGION

Table 6 Selected Drug Use, Perceptions of Great Risk, Average Annual Rates of First Use of Marijuana, Past Year Substance Dependence or Abuse, Needing But Not Receiving Treatment, and Past Year Mental Health Measures in the Midwest Region, by Age Group: Percentages, Annual Averages Based on 2009-2010 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLICIT DRUGS					
Past Month Illicit Drug Use ¹	8.10	9.76	20.34	5.75	7.92
Past Year Marijuana Use	10.79	13.04	29.66	7.21	10.55
Past Month Marijuana Use	6.34	7.13	17.55	4.28	6.26
Past Month Use of Illicit Drugs Other Than Marijuana ¹	3.31	4.49	8.00	2.35	3.19
Past Year Cocaine Use	1.54	0.93	4.27	1.14	1.61
Past Year Nonmedical Pain Reliever Use	4.80	6.58	11.99	3.33	4.61
Perception of Great Risk of Smoking Marijuana Once a Month	31.95	31.09	17.48	34.59	32.05
Average Annual Rate of First Use of Marijuana ²	1.83	5.71	7.46	0.14	1.16
ALCOHOL					
Past Month Alcohol Use	54.66	14.20	64.95	57.98	59.02
Past Month Binge Alcohol Use ³	25.54	8.71	44.88	24.29	27.35
Perception of Great Risk of Drinking Five or More Drinks Once or Twice a Week	38.97	38.07	29.51	40.73	39.06
Past Month Alcohol Use (Persons Aged 12 to 20)	27.16 ⁴	--	--	--	--
Past Month Binge Alcohol Use (Persons Aged 12 to 20) ³	18.78 ⁴	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁵	30.10	12.56	45.79	29.58	31.99
Past Month Cigarette Use	25.11	9.87	38.84	24.65	26.76
Perception of Great Risk of Smoking One or More Packs of Cigarettes per Day	67.42	64.34	61.07	68.91	67.75
PAST YEAR DEPENDENCE, ABUSE, AND TREATMENT⁶					
Illicit Drug Dependence ¹	1.85	2.37	5.14	1.21	1.79
Illicit Drug Dependence or Abuse ¹	2.68	4.41	7.53	1.61	2.49
Alcohol Dependence	3.38	1.76	7.01	2.95	3.55
Alcohol Dependence or Abuse	7.46	4.58	16.81	6.19	7.77
Alcohol or Illicit Drug Dependence or Abuse ¹	8.98	7.15	20.57	7.19	9.18
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,7}	2.39	4.09	6.91	1.39	2.21
Needing But Not Receiving Treatment for Alcohol Use ⁷	6.98	4.37	15.91	5.75	7.26
PAST YEAR MENTAL HEALTH					
Had at Least One Major Depressive Episode ⁸	--	8.20	8.52	6.72	6.99
Serious Mental Illness ⁹	--	--	3.83	4.28	4.21
Any Mental Illness ¹⁰	--	--	18.77	18.56	18.59
Had Serious Thoughts of Suicide	--	--	6.73	3.58	4.05

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

NOTE: Estimates for any mental illness (AMI) and serious mental illness (SMI) presented in this table may differ from previously published AMI and SMI estimates because of revised estimation procedures. For details, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates include data from original methamphetamine questions but do not include new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Average annual marijuana initiation rate = $100 * \{ [X_1 + (0.5 * X_1 + X_2)] \div 2 \}$, where X_1 is the number of marijuana initiates in the past 24 months and X_2 is the number of persons who never used marijuana. Both of the computation components, X_1 and X_2 , are based on a survey-weighted hierarchical Bayes estimation approach. Note that the age group is based on a respondent's age at the time of the interview, not his or her age at first use.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

⁴ Underage drinking is defined for persons aged 12 to 20; therefore, the "Total" estimate reflects that age group and not persons aged 12 or older.

⁵ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁶ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁷ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs (or alcohol), but not receiving treatment for an illicit drug (or alcohol) problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

⁸ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from persons aged 18 or older to produce an estimate for those aged 12 or older. For more details, see Section A.11 in Appendix A of the report on *State Estimates of Substance Use and Mental Disorders from the 2009-2010 NSDUHs*.

⁹ Serious mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in serious functional impairment.

¹⁰ Any mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012; SMI and AMI Estimates Revised October 2013).