

Table 2.72A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	172,843	171,827	83,415	83,066	70,757	70,257
GEOGRAPHIC DIVISION						
Northeast	32,290	31,951	15,406	15,180	13,019	12,642
New England	8,801	8,826	3,974	3,793	3,164	3,206
Middle Atlantic	23,488	23,125	11,432	11,387	9,856	9,437
Midwest	40,826	40,761	20,799	20,423	17,514	17,549
East North Central	28,536	28,399	14,406	14,271	12,121	12,273
West North Central	12,291	12,362	6,393	6,152	5,393	5,275
South	61,829	61,896	30,989	31,501	26,946	27,359
South Atlantic	32,948	32,981	15,442	16,439	13,301	14,219
East South Central	10,578	10,384	5,783	5,471	5,046	4,873
West South Central	18,303	18,531	9,764	9,591	8,599	8,267
West	37,897	37,218	16,221	15,962	13,277	12,707
Mountain	11,287	11,547	5,609	5,308	4,759	4,366
Pacific	26,610	25,672	10,611	10,654	8,518	8,341
COUNTY TYPE						
Large Metro	88,928	88,589	41,457	41,730	34,681	34,379
Small Metro	53,120	52,580	26,369	25,815	22,441	22,211
250K – 1 Mil. Pop.	35,023	33,853	17,146	16,377	14,486	13,895
< 250K Pop.	18,097	18,726	9,223	9,438	7,955	8,316
Nonmetro	30,795	30,658	15,589	15,521	13,635	13,666
Urbanized	13,720	12,518	6,990	6,544	6,161	5,652
Less Urbanized	13,916	15,418	6,997	7,812	6,039 ^a	6,980
Completely Rural	3,159	2,723	1,602	1,165	1,435	1,033

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.72B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	72.7 ^b	71.4	35.1	34.5	29.8	29.2
GEOGRAPHIC DIVISION						
Northeast	71.3	70.2	34.0	33.4	28.7	27.8
New England	74.0	73.9	33.4	31.7	26.6	26.8
Middle Atlantic	70.3	68.9	34.2	33.9	29.5	28.1
Midwest	75.9	75.2	38.7	37.7	32.6	32.4
East North Central	75.8	74.9	38.2	37.6	32.2	32.4
West North Central	76.3	76.0	39.7	37.8	33.5	32.4
South	72.9	71.9	36.6	36.6	31.8	31.8
South Atlantic	74.2	73.0	34.8	36.4	29.9	31.5
East South Central	74.4	72.1	40.7	38.0	35.5	33.9
West South Central	70.1	69.8	37.4	36.1	32.9	31.1
West	70.4 ^a	68.1	30.1	29.2	24.7	23.2
Mountain	72.3	72.2	35.9 ^a	33.2	30.5 ^b	27.3
Pacific	69.6 ^a	66.4	27.8	27.5	22.3	21.6
COUNTY TYPE						
Large Metro	70.5 ^a	69.0	32.9	32.5	27.5	26.8
Small Metro	74.4	73.4	36.9	36.0	31.4	31.0
250K – 1 Mil. Pop.	74.6 ^b	71.8	36.5	34.7	30.9	29.5
< 250K Pop.	74.0	76.5	37.7	38.6	32.5	34.0
Nonmetro	76.6	75.6	38.8	38.3	33.9	33.7
Urbanized	76.8	75.0	39.1	39.2	34.5	33.9
Less Urbanized	76.4	76.0	38.4	38.5	33.1	34.4
Completely Rural	77.1	76.3	39.1	32.7	35.0 ^a	29.0

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.73A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	8,631	8,249	5,634	5,562	3,610	3,635
GEOGRAPHIC DIVISION						
Northeast	1,502	1,419	997	1,052	635	660
New England	384	380	265	294	172	183
Middle Atlantic	1,118	1,039	732	758	463	477
Midwest	2,016	1,959	1,373	1,311	923	882
East North Central	1,377	1,345	939	901	613	592
West North Central	639	614	434	410	310	290
South	3,271	3,046	2,112	2,009	1,366	1,314
South Atlantic	1,594	1,513	1,007	974	645	645
East South Central	590	557	386	399	267	284
West South Central	1,087	976	719	636	455	385
West	1,841	1,825	1,152	1,190	686	779
Mountain	621	567	400	372	252	247
Pacific	1,220	1,258	752	818	433	532
COUNTY TYPE						
Large Metro	4,051	3,869	2,590	2,560	1,588	1,600
Small Metro	2,752	2,668	1,815	1,849	1,187	1,220
250K – 1 Mil. Pop.	1,804	1,735	1,168	1,214	746	783
< 250K Pop.	948	932	647	635	441	437
Nonmetro	1,827	1,712	1,229	1,153	835	815
Urbanized	749 ^a	628	503	429	345	305
Less Urbanized	908	938	608	637	401	450
Completely Rural	170	145	118	86	89	60

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.73B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	34.5 ^b	32.7	22.5	22.1	14.4	14.4
GEOGRAPHIC DIVISION						
Northeast	33.5	31.3	22.2	23.2	14.1	14.6
New England	32.8	32.4	22.6	25.1	14.7	15.6
Middle Atlantic	33.7	30.9	22.1	22.5	13.9	14.2
Midwest	35.4	34.5	24.1	23.1	16.2	15.5
East North Central	34.5	33.6	23.5	22.5	15.4	14.8
West North Central	37.6	36.7	25.5	24.5	18.2	17.3
South	36.6 ^b	33.8	23.7	22.3	15.3	14.6
South Atlantic	35.4	33.0	22.4	21.3	14.3	14.1
East South Central	40.6	38.5	26.6	27.5	18.4	19.6
West South Central	36.6 ^a	32.8	24.2	21.4	15.3	12.9
West	31.3	30.4	19.6	19.8	11.7	13.0
Mountain	35.9	32.5	23.2	21.3	14.6	14.2
Pacific	29.4	29.6	18.1	19.2	10.4	12.5
COUNTY TYPE						
Large Metro	30.4	29.0	19.5	19.2	11.9	12.0
Small Metro	37.2 ^a	34.9	24.5	24.2	16.0	16.0
250K – 1 Mil. Pop.	36.2	34.0	23.5	23.8	15.0	15.3
< 250K Pop.	39.2	36.7	26.7	25.0	18.2	17.2
Nonmetro	42.6	40.5	28.7	27.3	19.5	19.3
Urbanized	40.3	36.9	27.1	25.3	18.6	17.9
Less Urbanized	45.7	43.7	30.6	29.7	20.2	21.0
Completely Rural	38.4	38.5	26.7	22.9	20.1	16.0

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.74A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	23,723	23,658	17,073	17,481	14,201	14,359
GEOGRAPHIC DIVISION						
Northeast	4,215	4,206	3,117	3,208	2,523	2,645
New England	1,138	1,126	816	871	649	729
Middle Atlantic	3,077	3,080	2,301	2,337	1,874	1,916
Midwest	5,819	5,910	4,318	4,506	3,573	3,779
East North Central	3,976	4,007	2,938	3,068	2,431	2,582
West North Central	1,843	1,903	1,380	1,438	1,141	1,197
South	8,367	8,337	6,083	6,137	5,212	5,132
South Atlantic	4,054	4,164	2,942	3,065	2,479	2,550
East South Central	1,448	1,439	1,071	1,109	949	946
West South Central	2,865	2,734	2,069	1,963	1,784	1,635
West	5,322	5,204	3,556	3,631	2,894	2,803
Mountain	1,640	1,664	1,100	1,187	915	963
Pacific	3,683	3,540	2,456	2,444	1,978	1,840
COUNTY TYPE						
Large Metro	11,979	11,826	8,433	8,565	6,921	6,906
Small Metro	7,792	7,931	5,684	5,907	4,755	4,856
250K – 1 Mil. Pop.	5,045	5,024	3,670	3,722	3,057	3,015
< 250K Pop.	2,747	2,907	2,014	2,186	1,697	1,841
Nonmetro	3,951	3,901	2,956	3,009	2,526	2,597
Urbanized	1,890	1,798	1,410	1,415	1,194	1,214
Less Urbanized	1,755	1,827	1,318	1,376	1,124	1,193
Completely Rural	305	276	227	217	208	190

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.74B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	74.8 ^a	73.5	53.8	54.3	44.8	44.6
GEOGRAPHIC DIVISION						
Northeast	74.8	74.0	55.3	56.4	44.8	46.5
New England	77.4	75.7	55.5	58.5	44.1 ^a	49.0
Middle Atlantic	73.9	73.4	55.3	55.7	45.0	45.7
Midwest	79.1	79.4	58.7	60.6	48.6	50.8
East North Central	78.2	78.2	57.8	59.8	47.8	50.4
West North Central	81.0	82.2	60.7	62.1	50.2	51.7
South	73.9	72.3	53.7	53.2	46.0	44.5
South Atlantic	72.6	73.0	52.7	53.7	44.4	44.7
East South Central	74.8	73.8	55.4	56.8	49.0	48.5
West South Central	75.3 ^a	70.4	54.4 ^a	50.6	46.9 ^b	42.1
West	71.8	69.1	48.0	48.2	39.0	37.2
Mountain	73.3	72.7	49.2	51.9	40.9	42.1
Pacific	71.2	67.5	47.5	46.6	38.2	35.1
COUNTY TYPE						
Large Metro	72.1	71.0	50.8	51.4	41.7	41.5
Small Metro	76.3	74.8	55.7	55.7	46.6	45.8
250K – 1 Mil. Pop.	75.9 ^a	72.9	55.2	54.0	46.0	43.8
< 250K Pop.	77.3	78.2	56.7	58.8	47.7	49.5
Nonmetro	80.3	79.0	60.1	60.9	51.4	52.6
Urbanized	80.0	78.1	59.7	61.5	50.5	52.7
Less Urbanized	80.1	79.0	60.1	59.5	51.3	51.6
Completely Rural	84.0	85.0	62.6	67.0	57.2	58.7

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.75A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	140,489	139,921	60,708	60,023	52,946	52,263
GEOGRAPHIC DIVISION						
Northeast	26,573	26,326	11,292	10,921	9,862	9,337
New England	7,280	7,319	2,893	2,627	2,343	2,293
Middle Atlantic	19,293	19,006	8,399	8,293	7,519	7,044
Midwest	32,992	32,892	15,108	14,606	13,018	12,888
East North Central	23,182	23,047	10,529	10,302	9,076	9,099
West North Central	9,809	9,845	4,579	4,304	3,942	3,789
South	50,191	50,514	22,795	23,355	20,368	20,913
South Atlantic	27,300	27,305	11,493	12,400	10,177	11,023
East South Central	8,540	8,387	4,326	3,963	3,831	3,643
West South Central	14,351	14,821	6,976	6,992	6,360	6,247
West	30,734	30,189	11,513	11,142	9,698	9,124
Mountain	9,026	9,316	4,109	3,749	3,591	3,155
Pacific	21,707	20,874	7,403	7,392	6,107	5,969
COUNTY TYPE						
Large Metro	72,897	72,894	30,434	30,606	26,172	25,874
Small Metro	42,575	41,981	18,870	18,058	16,500	16,135
250K – 1 Mil. Pop.	28,173	27,094	12,308	11,441	10,683	10,097
< 250K Pop.	14,402	14,887	6,562	6,617	5,817	6,038
Nonmetro	25,017	25,046	11,404	11,359	10,274	10,254
Urbanized	11,080	10,092	5,077	4,700	4,622	4,134
Less Urbanized	11,252	12,653	5,070	5,799	4,514 ^a	5,338
Completely Rural	2,684	2,302	1,256	861	1,139	783

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.75B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	77.6 ^a	76.4	33.5	32.8	29.3	28.5
GEOGRAPHIC DIVISION						
Northeast	75.5	74.6	32.1	31.0	28.0	26.5
New England	78.8	78.8	31.3 ^a	28.3	25.3	24.7
Middle Atlantic	74.4	73.1	32.4	31.9	29.0	27.1
Midwest	81.0	80.1	37.1	35.5	32.0	31.4
East North Central	81.1	80.0	36.8	35.8	31.7	31.6
West North Central	80.9	80.1	37.8	35.0	32.5	30.8
South	77.8	77.0	35.3	35.6	31.6	31.9
South Atlantic	79.5	78.2	33.5	35.5	29.6	31.6
East South Central	78.9	76.3	39.9	36.1	35.4	33.1
West South Central	74.2	75.2	36.1	35.5	32.9	31.7
West	75.8	73.4	28.4	27.1	23.9	22.2
Mountain	77.5	77.9	35.3 ^a	31.4	30.8 ^b	26.4
Pacific	75.1 ^a	71.5	25.6	25.3	21.1	20.5
COUNTY TYPE						
Large Metro	75.8	74.1	31.6	31.1	27.2	26.3
Small Metro	79.2	78.6	35.1	33.8	30.7	30.2
250K – 1 Mil. Pop.	79.8 ^a	77.0	34.9	32.5	30.3	28.7
< 250K Pop.	77.9 ^a	81.7	35.5	36.3	31.5	33.2
Nonmetro	80.7	79.8	36.8	36.2	33.2	32.7
Urbanized	81.2	79.5	37.2	37.0	33.9	32.6
Less Urbanized	80.1	79.9	36.1	36.6	32.1	33.7
Completely Rural	81.5	80.3	38.1	30.0	34.6	27.3

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.76A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	164,212	163,579	77,781	77,504	67,147	66,622
GEOGRAPHIC DIVISION						
Northeast	30,787	30,532	14,409	14,128	12,385	11,982
New England	8,417	8,446	3,709	3,498	2,992	3,022
Middle Atlantic	22,370	22,087	10,700	10,630	9,393	8,960
Midwest	38,810	38,803	19,426	19,112	16,591	16,667
East North Central	27,159	27,054	13,467	13,370	11,508	11,681
West North Central	11,652	11,748	5,959	5,742	5,083	4,986
South	58,558	58,850	28,878	29,492	25,580	26,045
South Atlantic	31,354	31,469	14,435	15,465	12,656	13,574
East South Central	9,988	9,827	5,397	5,072	4,780	4,589
West South Central	17,216	17,555	9,045	8,954	8,144	7,882
West	36,056	35,394	15,068	14,773	12,592	11,928
Mountain	10,666	10,980	5,209	4,936	4,507	4,118
Pacific	25,390	24,414	9,859	9,837	8,085	7,809
COUNTY TYPE						
Large Metro	84,876	84,720	38,867	39,170	33,093	32,780
Small Metro	50,368	49,912	24,554	23,966	21,254	20,991
250K – 1 Mil. Pop.	33,219	32,118	15,978	15,162	13,740	13,112
< 250K Pop.	17,149	17,794	8,576	8,803	7,514	7,879
Nonmetro	28,968	28,947	14,359	14,368	12,800	12,851
Urbanized	12,971	11,890	6,487	6,115	5,816	5,348
Less Urbanized	13,008	14,480	6,388	7,175	5,638 ^a	6,530
Completely Rural	2,989	2,577	1,484	1,078	1,347	973

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.76B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	77.2 ^a	76.0	36.6	36.0	31.6	30.9
GEOGRAPHIC DIVISION						
Northeast	75.4	74.5	35.3	34.5	30.3	29.3
New England	78.6	78.4	34.6	32.5	27.9	28.1
Middle Atlantic	74.3	73.2	35.5	35.2	31.2	29.7
Midwest	80.7	80.0	40.4	39.4	34.5	34.3
East North Central	80.7	79.7	40.0	39.4	34.2	34.4
West North Central	80.9	80.5	41.4	39.3	35.3	34.1
South	77.2	76.3	38.1	38.2	33.7	33.8
South Atlantic	78.5	77.5	36.2	38.1	31.7	33.4
East South Central	78.3	75.9	42.3	39.2	37.4	35.5
West South Central	74.4	74.4	39.1	38.0	35.2	33.4
West	75.2 ^a	72.7	31.4	30.4	26.3	24.5
Mountain	76.8	77.1	37.5	34.7	32.5 ^b	28.9
Pacific	74.5 ^a	70.9	28.9	28.6	23.7	22.7
COUNTY TYPE						
Large Metro	75.3 ^a	73.7	34.5	34.1	29.3	28.5
Small Metro	78.7	78.0	38.4	37.5	33.2	32.8
250K – 1 Mil. Pop.	79.2 ^b	76.4	38.1	36.0	32.8	31.2
< 250K Pop.	77.8 ^a	81.1	38.9	40.1	34.1	35.9
Nonmetro	80.7	79.7	40.0	39.5	35.7	35.4
Urbanized	81.0	79.3	40.5	40.8	36.3	35.7
Less Urbanized	80.1	79.8	39.3	39.5	34.7	36.0
Completely Rural	81.8	80.8	40.6	33.8	36.8	30.5

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.