

Table 1.98A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	7,949	7,840	1,406	1,304	604	467
GEOGRAPHIC DIVISION						
Northeast	1,373	1,316	220	195	69	88
New England	408	352	41	48	14	14
Middle Atlantic	964	964	178	147	55	74
Midwest	1,643	1,688	336	425	115	155
East North Central	1,071	1,196	219	329	97	129
West North Central	572	492	117	96	19	27
South	2,779	2,939	639	490	318	206
South Atlantic	1,583	1,723	322	318	184	144
East South Central	346 ^a	510	132	105	59	32
West South Central	850	707	184 ^b	67	75	30
West	2,153	1,896	211	194	102 ^b	17
Mountain	707	608	102	50	47	11
Pacific	1,446	1,288	110	144	55 ^a	6
COUNTY TYPE						
Large Metro	4,445	4,153	769	657	352	245
Small Metro	2,328	2,438	408	425	158	147
250K – 1 Mil. Pop.	1,495	1,649	232	268	90	86
< 250K Pop.	833	789	176	157	67	61
Nonmetro	1,175	1,249	229	222	94	75
Urbanized	552	655	85	99	21	36
Less Urbanized	536	535	127	109	65	35
Completely Rural	87	59	17	13	8	4

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.98B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	3.3	3.3	0.6	0.5	0.3	0.2
GEOGRAPHIC DIVISION						
Northeast	3.0	2.9	0.5	0.4	0.2	0.2
New England	3.4	2.9	0.3	0.4	0.1	0.1
Middle Atlantic	2.9	2.9	0.5	0.4	0.2	0.2
Midwest	3.1	3.1	0.6	0.8	0.2	0.3
East North Central	2.8	3.2	0.6	0.9	0.3	0.3
West North Central	3.6	3.0	0.7	0.6	0.1	0.2
South	3.3	3.4	0.8	0.6	0.4	0.2
South Atlantic	3.6	3.8	0.7	0.7	0.4	0.3
East South Central	2.4 ^a	3.5	0.9	0.7	0.4	0.2
West South Central	3.3	2.7	0.7 ^b	0.3	0.3	0.1
West	4.0	3.5	0.4	0.4	0.2 ^b	0.0
Mountain	4.5	3.8	0.7	0.3	0.3	0.1
Pacific	3.8	3.3	0.3	0.4	0.1 ^a	0.0
COUNTY TYPE						
Large Metro	3.5	3.2	0.6	0.5	0.3	0.2
Small Metro	3.3	3.4	0.6	0.6	0.2	0.2
250K – 1 Mil. Pop.	3.2	3.5	0.5	0.6	0.2	0.2
< 250K Pop.	3.4	3.2	0.7	0.6	0.3	0.2
Nonmetro	2.9	3.1	0.6	0.5	0.2	0.2
Urbanized	3.1	3.9	0.5	0.6	0.1	0.2
Less Urbanized	2.9	2.6	0.7	0.5	0.4	0.2
Completely Rural	2.1	1.7	0.4	0.4	0.2	0.1

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.99A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	157	125	95	63	32	16
GEOGRAPHIC DIVISION						
Northeast	16	22	10	13	1	4
New England	3	10	1	8	1	0
Middle Atlantic	14	13	9	6	1	3
Midwest	42 ^a	23	27	15	14 ^b	3
East North Central	23	19	14	13	5	2
West North Central	19 ^b	4	13 ^a	2	9 ^a	1
South	53	40	34	21	12	6
South Atlantic	22	14	18 ^a	4	6	2
East South Central	3	9	2	2	1	*
West South Central	28	17	14	14	5	4
West	45	40	25	14	5	4
Mountain	19	17	15	7	5	2
Pacific	26	23	10	6	*	2
COUNTY TYPE						
Large Metro	83 ^a	48	51 ^a	23	21 ^a	4
Small Metro	36	54	21	26	7	7
250K – 1 Mil. Pop.	19	36	11	17	4	4
< 250K Pop.	17	18	11	9	3	3
Nonmetro	37	23	23	14	5	5
Urbanized	12	9	7	4	2	*
Less Urbanized	18	13	12	9	2	5
Completely Rural	8	2	3	1	0	*

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.99B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.6	0.5	0.4 ^a	0.3	0.1	0.1
GEOGRAPHIC DIVISION						
Northeast	0.4	0.5	0.2	0.3	0.0	0.1
New England	0.2	0.8	0.1	0.6	0.0	0.0
Middle Atlantic	0.4	0.4	0.3	0.2	0.0	0.1
Midwest	0.7 ^a	0.4	0.5	0.3	0.2 ^b	0.0
East North Central	0.6	0.5	0.3	0.3	0.1	0.1
West North Central	1.1 ^b	0.2	0.8 ^a	0.1	0.5 ^a	0.0
South	0.6	0.4	0.4	0.2	0.1	0.1
South Atlantic	0.5	0.3	0.4 ^a	0.1	0.1	0.1
East South Central	0.2	0.6	0.1	0.2	0.1	*
West South Central	0.9	0.6	0.5	0.5	0.2	0.1
West	0.8	0.7	0.4	0.2	0.1	0.1
Mountain	1.1	1.0	0.8	0.4	0.3	0.1
Pacific	0.6	0.5	0.2	0.2	*	0.0
COUNTY TYPE						
Large Metro	0.6 ^a	0.4	0.4 ^a	0.2	0.2 ^a	0.0
Small Metro	0.5	0.7	0.3	0.3	0.1	0.1
250K – 1 Mil. Pop.	0.4	0.7	0.2	0.3	0.1	0.1
< 250K Pop.	0.7	0.7	0.4	0.4	0.1	0.1
Nonmetro	0.9	0.6	0.5	0.3	0.1	0.1
Urbanized	0.6	0.5	0.4	0.3	0.1	*
Less Urbanized	0.9	0.6	0.6	0.4	0.1	0.2
Completely Rural	1.9	0.5	0.7	0.3	0.0	*

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.100A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	1,193	1,135	275	263	55	80
GEOGRAPHIC DIVISION						
Northeast	157	188	42	55	6 ^a	21
New England	44	60	14	29	2	13
Middle Atlantic	114	128	28	25	5	8
Midwest	282	275	78	73	13	26
East North Central	175	158	49	48	8	14
West North Central	106	117	29	25	5	11
South	348	360	81	85	10	26
South Atlantic	153	175	36	45	7	13
East South Central	70	88	16	21	2	5
West South Central	126	97	29	18	1	7
West	406	313	73	50	25	8
Mountain	124	134	28	21	9	5
Pacific	282 ^a	179	46	29	16	4
COUNTY TYPE						
Large Metro	565	465	118	85	29	24
Small Metro	424	457	105	116	19	39
250K – 1 Mil. Pop.	278	268	68	68	11	24
< 250K Pop.	146	189	37	49	8	15
Nonmetro	204	213	51	62	6	18
Urbanized	97	86	26	27	4	8
Less Urbanized	93	107	20	32	2	10
Completely Rural	14	20	5	*	*	0

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.100B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	3.8	3.5	0.9	0.8	0.2	0.3
GEOGRAPHIC DIVISION						
Northeast	2.8	3.3	0.7	1.0	0.1 ^a	0.4
New England	3.0	4.0	1.0	2.0	0.1 ^a	0.9
Middle Atlantic	2.7	3.0	0.7	0.6	0.1	0.2
Midwest	3.8	3.7	1.1	1.0	0.2	0.3
East North Central	3.4	3.1	1.0	0.9	0.2	0.3
West North Central	4.7	5.0	1.3	1.1	0.2	0.5
South	3.1	3.1	0.7	0.7	0.1	0.2
South Atlantic	2.7	3.1	0.6	0.8	0.1	0.2
East South Central	3.6	4.5	0.8	1.1	0.1	0.3
West South Central	3.3	2.5	0.8	0.5	0.0	0.2
West	5.5 ^a	4.2	1.0	0.7	0.3	0.1
Mountain	5.5	5.9	1.2	0.9	0.4	0.2
Pacific	5.5 ^a	3.4	0.9	0.6	0.3	0.1
COUNTY TYPE						
Large Metro	3.4	2.8	0.7	0.5	0.2	0.1
Small Metro	4.2	4.3	1.0	1.1	0.2	0.4
250K – 1 Mil. Pop.	4.2	3.9	1.0	1.0	0.2	0.3
< 250K Pop.	4.1	5.1	1.0	1.3	0.2	0.4
Nonmetro	4.1	4.3	1.0	1.2	0.1	0.4
Urbanized	4.1	3.7	1.1	1.2	0.2	0.3
Less Urbanized	4.3	4.6	0.9	1.4	0.1	0.4
Completely Rural	3.9	6.1	1.5	*	*	0.0

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.101A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	6,599	6,580	1,036	978	517	370
GEOGRAPHIC DIVISION						
Northeast	1,199	1,107	168	127	61	63
New England	362	283	26	11	12 ^a	1
Middle Atlantic	837	824	142	116	49	63
Midwest	1,320	1,391	231	337	88	127
East North Central	873	1,020	156	268	83	113
West North Central	447	371	75	68	5	15
South	2,378	2,540	524	385	296	175
South Atlantic	1,408	1,533	268	269	171	129
East South Central	273	413	114	81	55	27
West South Central	697	593	142 ^b	34	70	19
West	1,702	1,543	113	130	72 ^a	5
Mountain	564	457	60	22	33	5
Pacific	1,139	1,086	54	109	39	*
COUNTY TYPE						
Large Metro	3,797	3,640	599	549	302	217
Small Metro	1,868	1,927	281	283	132	101
250K – 1 Mil. Pop.	1,198	1,345	153	183	76	58
< 250K Pop.	670	582	128	100	56	44
Nonmetro	934	1,013	155	146	83	53
Urbanized	444	560	52	67	14	29
Less Urbanized	426	415	95	68	61	20
Completely Rural	64	38	8	10	8	4

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.101B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	3.6	3.6	0.6	0.5	0.3	0.2
GEOGRAPHIC DIVISION						
Northeast	3.4	3.1	0.5	0.4	0.2	0.2
New England	3.9	3.0	0.3	0.1	0.1 ^a	0.0
Middle Atlantic	3.2	3.2	0.5	0.4	0.2	0.2
Midwest	3.2	3.4	0.6	0.8	0.2	0.3
East North Central	3.1	3.5	0.5	0.9	0.3	0.4
West North Central	3.7	3.0	0.6	0.6	0.0	0.1
South	3.7	3.9	0.8	0.6	0.5	0.3
South Atlantic	4.1	4.4	0.8	0.8	0.5	0.4
East South Central	2.5	3.8	1.1	0.7	0.5	0.2
West South Central	3.6	3.0	0.7 ^b	0.2	0.4	0.1
West	4.2	3.8	0.3	0.3	0.2 ^a	0.0
Mountain	4.8	3.8	0.5	0.2	0.3	0.0
Pacific	3.9	3.7	0.2	0.4	0.1	*
COUNTY TYPE						
Large Metro	3.9	3.7	0.6	0.6	0.3	0.2
Small Metro	3.5	3.6	0.5	0.5	0.2	0.2
250K – 1 Mil. Pop.	3.4	3.8	0.4	0.5	0.2	0.2
< 250K Pop.	3.6	3.2	0.7	0.5	0.3	0.2
Nonmetro	3.0	3.2	0.5	0.5	0.3	0.2
Urbanized	3.3	4.4	0.4	0.5	0.1	0.2
Less Urbanized	3.0	2.6	0.7	0.4	0.4	0.1
Completely Rural	1.9	1.3	0.3	0.4	0.3	0.1

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.102A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	7,792	7,715	1,310	1,241	572	451
GEOGRAPHIC DIVISION						
Northeast	1,356	1,294	210	182	68	84
New England	406	342	40	40	14	13
Middle Atlantic	950	952	170	141	54	71
Midwest	1,601	1,665	309	410	101	153
East North Central	1,048	1,177	205	316	91	127
West North Central	553	488	104	94	10	26
South	2,726	2,899	605	469	306	200
South Atlantic	1,561	1,708	304	314	178	142
East South Central	343 ^a	501	130	103	58	32
West South Central	822	690	170 ^b	52	70	26
West	2,108	1,856	187	181	97 ^a	13
Mountain	688	591	87	43	42	9
Pacific	1,421	1,265	100	138	55 ^a	4
COUNTY TYPE						
Large Metro	4,362	4,105	718	634	331	241
Small Metro	2,292	2,384	386	399	151	140
250K – 1 Mil. Pop.	1,476	1,613	221	251	87	81
< 250K Pop.	816	771	165	148	64	58
Nonmetro	1,138	1,225	206	208	90	70
Urbanized	541	646	78	95	19	36
Less Urbanized	519	522	115	101	63	30
Completely Rural	78	57	14	12	8	4

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.102B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	3.7	3.6	0.6	0.6	0.3	0.2
GEOGRAPHIC DIVISION						
Northeast	3.3	3.2	0.5	0.4	0.2	0.2
New England	3.8	3.2	0.4	0.4	0.1	0.1
Middle Atlantic	3.2	3.2	0.6	0.5	0.2	0.2
Midwest	3.3	3.4	0.6	0.8	0.2	0.3
East North Central	3.1	3.5	0.6	0.9	0.3	0.4
West North Central	3.8	3.3	0.7	0.6	0.1	0.2
South	3.6	3.8	0.8	0.6	0.4	0.3
South Atlantic	3.9	4.2	0.8	0.8	0.4	0.3
East South Central	2.7 ^a	3.9	1.0	0.8	0.5	0.3
West South Central	3.6	2.9	0.7 ^b	0.2	0.3	0.1
West	4.4	3.8	0.4	0.4	0.2 ^b	0.0
Mountain	5.0	4.1	0.6	0.3	0.3	0.1
Pacific	4.2	3.7	0.3	0.4	0.2 ^a	0.0
COUNTY TYPE						
Large Metro	3.9	3.6	0.6	0.6	0.3	0.2
Small Metro	3.6	3.7	0.6	0.6	0.2	0.2
250K – 1 Mil. Pop.	3.5	3.8	0.5	0.6	0.2	0.2
< 250K Pop.	3.7	3.5	0.7	0.7	0.3	0.3
Nonmetro	3.2	3.4	0.6	0.6	0.2	0.2
Urbanized	3.4	4.3	0.5	0.6	0.1	0.2
Less Urbanized	3.2	2.9	0.7	0.6	0.4	0.2
Completely Rural	2.1	1.8	0.4	0.4	0.2	0.1

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.