

**Table 1.123A Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	71,128	70,657	20,305	19,658	8,849	8,247
<b>GEOGRAPHIC DIVISION</b>						
Northeast	13,002	13,242	3,417	3,381	1,424	1,487
New England	3,962	3,787	1,154	931	410	415
Middle Atlantic	9,040	9,455	2,263	2,449	1,014	1,072
Midwest	15,446	15,081	4,246	4,230	1,804	1,737
East North Central	10,903	10,900	2,966	3,125	1,283	1,295
West North Central	4,543	4,181	1,280	1,105	521	443
South	23,081	24,504	7,063	6,843	3,166	3,119
South Atlantic	12,177 <sup>a</sup>	13,345	3,660	3,537	1,601	1,608
East South Central	3,686	3,769	1,074	1,203	462	570
West South Central	7,218	7,391	2,329	2,103	1,102	941
West	19,599 <sup>a</sup>	17,830	5,579	5,205	2,455 <sup>a</sup>	1,904
Mountain	5,794	5,388	1,748	1,637	751	650
Pacific	13,804 <sup>a</sup>	12,442	3,831	3,568	1,704 <sup>a</sup>	1,254
<b>COUNTY TYPE</b>						
Large Metro	39,323	38,942	10,863	10,379	4,763	4,189
Small Metro	21,387	21,451	6,430	6,277	2,758	2,828
250K – 1 Mil. Pop.	14,096	14,308	4,135	4,215	1,711	1,880
< 250K Pop.	7,291	7,143	2,295	2,062	1,047	947
Nonmetro	10,418	10,263	3,012	3,002	1,328	1,231
Urbanized	4,990	4,650	1,480	1,357	645	568
Less Urbanized	4,559	4,772	1,364	1,434	624	581
Completely Rural	869	841	168	212	59	82

\*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.123B Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2003 and 2004**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	29.9	29.4	8.5	8.2	3.7	3.4
<b>GEOGRAPHIC DIVISION</b>						
Northeast	28.7	29.1	7.5	7.4	3.1	3.3
New England	33.3	31.7	9.7	7.8	3.4	3.5
Middle Atlantic	27.0	28.2	6.8	7.3	3.0	3.2
Midwest	28.7	27.8	7.9	7.8	3.4	3.2
East North Central	28.9	28.7	7.9	8.2	3.4	3.4
West North Central	28.2	25.7	7.9	6.8	3.2	2.7
South	27.2	28.4	8.3	7.9	3.7	3.6
South Atlantic	27.4	29.5	8.2	7.8	3.6	3.6
East South Central	25.9	26.2	7.6	8.4	3.3	4.0
West South Central	27.6	27.8	8.9	7.9	4.2	3.5
West	36.4 <sup>b</sup>	32.6	10.4	9.5	4.6 <sup>a</sup>	3.5
Mountain	37.1 <sup>a</sup>	33.7	11.2	10.2	4.8	4.1
Pacific	36.1 <sup>b</sup>	32.2	10.0	9.2	4.5 <sup>a</sup>	3.2
<b>COUNTY TYPE</b>						
Large Metro	31.2	30.3	8.6	8.1	3.8 <sup>a</sup>	3.3
Small Metro	30.0	29.9	9.0	8.8	3.9	3.9
250K – 1 Mil. Pop.	30.0	30.3	8.8	8.9	3.6	4.0
< 250K Pop.	29.8	29.2	9.4	8.4	4.3	3.9
Nonmetro	25.9	25.3	7.5	7.4	3.3	3.0
Urbanized	27.9	27.9	8.3	8.1	3.6	3.4
Less Urbanized	25.0	23.5	7.5	7.1	3.4	2.9
Completely Rural	21.2	23.6	4.1	5.9	1.4	2.3

\*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.124A Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	5,319	5,355	3,351	3,269	1,417	1,328
<b>GEOGRAPHIC DIVISION</b>						
Northeast	847	891	542	561	235	242
New England	241	248	161	169	59	71
Middle Atlantic	606	643	381	392	175	171
Midwest	1,160	1,176	723	722	291	284
East North Central	841	849	527	523	215	203
West North Central	319	327	195	198	76	81
South	1,912	1,874	1,213	1,135	514	457
South Atlantic	914	945	579	568	222	213
East South Central	337	320	210	200	88	75
West South Central	661	609	425	367	204	169
West	1,399	1,414	873	852	377	344
Mountain	435	394	283	240	125	90
Pacific	964	1,021	590	612	253	254
<b>COUNTY TYPE</b>						
Large Metro	2,667	2,672	1,656	1,608	669	647
Small Metro	1,692	1,699	1,071	1,050	475	435
250K – 1 Mil. Pop.	1,130	1,111	696	676	319	279
< 250K Pop.	563	588	375	373	156	156
Nonmetro	959	984	624	612	273	245
Urbanized	430	384	290	248	131	99
Less Urbanized	434	518	276	317	116	131
Completely Rural	95	81	57	47	26	16

\*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.124B Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2003 and 2004**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	21.3	21.2	13.4	13.0	5.7	5.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	18.9	19.6	12.1	12.4	5.2	5.3
New England	20.5	21.1	13.7	14.4	5.1	6.0
Middle Atlantic	18.3	19.1	11.5	11.7	5.3	5.1
Midwest	20.4	20.7	12.7	12.7	5.1	5.0
East North Central	21.1	21.2	13.2	13.1	5.4	5.1
West North Central	18.8	19.5	11.5	11.9	4.5	4.9
South	21.4	20.8	13.6	12.6	5.8	5.1
South Atlantic	20.3	20.6	12.9	12.4	4.9	4.7
East South Central	23.2	22.1	14.4	13.8	6.0	5.2
West South Central	22.3	20.5	14.3	12.4	6.9	5.7
West	23.8	23.6	14.8	14.2	6.4	5.7
Mountain	25.2	22.6	16.4	13.7	7.2 <sup>a</sup>	5.2
Pacific	23.2	24.0	14.2	14.4	6.1	6.0
<b>COUNTY TYPE</b>						
Large Metro	20.0	20.0	12.4	12.0	5.0	4.8
Small Metro	22.9	22.2	14.5	13.7	6.4	5.7
250K – 1 Mil. Pop.	22.7	21.8	14.0	13.3	6.4	5.5
< 250K Pop.	23.2	23.1	15.5	14.7	6.5	6.2
Nonmetro	22.4	23.3	14.6	14.5	6.4	5.8
Urbanized	23.1	22.6	15.6	14.6	7.0	5.8
Less Urbanized	21.8	24.1	13.9	14.7	5.8	6.1
Completely Rural	21.6	21.5	13.0	12.5	5.9	4.1

\*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.125A Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	12,770	12,630	6,266	6,201	2,665	2,619
<b>GEOGRAPHIC DIVISION</b>						
Northeast	2,256	2,248	1,180	1,214	493	534
New England	705	637	374	347	152	168
Middle Atlantic	1,551	1,611	807	867	341	366
Midwest	2,938	2,846	1,378	1,407	583	564
East North Central	2,054	2,009	954	1,011	415	410
West North Central	885	837	425	396	169	154
South	4,291	4,364	2,101	2,132	955	970
South Atlantic	2,091	2,135	1,023	1,000	481	462
East South Central	703	757	335	363	137	169
West South Central	1,497	1,472	743	769	337	340
West	3,284	3,172	1,606	1,447	633	550
Mountain	1,068	1,034	503	501	191	173
Pacific	2,216	2,138	1,103	946	442	377
<b>COUNTY TYPE</b>						
Large Metro	6,591	6,349	3,224	3,062	1,382	1,226
Small Metro	4,218	4,358	2,055	2,188	863	961
250K – 1 Mil. Pop.	2,774	2,778	1,328	1,411	553	609
< 250K Pop.	1,443	1,580	727	777	309	352
Nonmetro	1,962	1,922	987	951	420	432
Urbanized	983	935	514	463	239	202
Less Urbanized	867	861	429	419	166	197
Completely Rural	111	126	44	70	15	33

\*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.125B Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2003 and 2004**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	40.2	39.2	19.7	19.3	8.4	8.1
<b>GEOGRAPHIC DIVISION</b>						
Northeast	40.0	39.5	21.0	21.4	8.8	9.4
New England	47.9 <sup>a</sup>	42.8	25.4	23.3	10.3	11.3
Middle Atlantic	37.3	38.4	19.4	20.7	8.2	8.7
Midwest	39.9	38.3	18.7	18.9	7.9	7.6
East North Central	40.4	39.2	18.8	19.7	8.2	8.0
West North Central	38.9	36.2	18.7	17.1	7.4	6.7
South	37.9	37.8	18.6	18.5	8.4	8.4
South Atlantic	37.4	37.4	18.3	17.5	8.6	8.1
East South Central	36.4	38.8	17.3	18.6	7.1	8.7
West South Central	39.3	37.9	19.5	19.8	8.9	8.8
West	44.3	42.1	21.7	19.2	8.5	7.3
Mountain	47.7	45.2	22.5	21.9	8.5	7.6
Pacific	42.8	40.8	21.3	18.0	8.5	7.2
<b>COUNTY TYPE</b>						
Large Metro	39.7	38.1	19.4	18.4	8.3	7.4
Small Metro	41.3	41.1	20.1	20.6	8.5	9.1
250K – 1 Mil. Pop.	41.7	40.3	20.0	20.5	8.3	8.8
< 250K Pop.	40.6	42.5	20.4	20.9	8.7	9.5
Nonmetro	39.9	38.9	20.1	19.2	8.5	8.7
Urbanized	41.6	40.6	21.7	20.1	10.1	8.8
Less Urbanized	39.6	37.2	19.6	18.1	7.6	8.5
Completely Rural	30.5	38.8	12.2 <sup>a</sup>	21.4	4.0	10.3

\*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.126A Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	53,040	52,672	10,689	10,189	4,767	4,301
<b>GEOGRAPHIC DIVISION</b>						
Northeast	9,899	10,103	1,694	1,605	696	711
New England	3,016	2,902	620 <sup>a</sup>	415	198	176
Middle Atlantic	6,883	7,201	1,074	1,190	498	535
Midwest	11,348	11,059	2,145	2,102	929	888
East North Central	8,008	8,042	1,485	1,590	653	681
West North Central	3,340	3,017	660	511	276	207
South	16,877 <sup>a</sup>	18,267	3,749	3,576	1,697	1,692
South Atlantic	9,172 <sup>a</sup>	10,264	2,058	1,969	898	933
East South Central	2,646	2,692	530	640	238	327
West South Central	5,059	5,310	1,161	967	561	432
West	14,916 <sup>a</sup>	13,243	3,100	2,905	1,445 <sup>a</sup>	1,010
Mountain	4,292	3,960	962	896	436	387
Pacific	10,624 <sup>a</sup>	9,283	2,138	2,010	1,009 <sup>a</sup>	622
<b>COUNTY TYPE</b>						
Large Metro	30,065	29,920	5,984	5,710	2,712	2,316
Small Metro	15,477	15,394	3,304	3,039	1,420	1,432
250K – 1 Mil. Pop.	10,191	10,419	2,111	2,128	839	992
< 250K Pop.	5,286	4,975	1,193 <sup>a</sup>	911	581	439
Nonmetro	7,497	7,358	1,401	1,440	635	553
Urbanized	3,576	3,332	677	646	275	267
Less Urbanized	3,258	3,392	658	699	341	253
Completely Rural	663	634	67	95	19	33

\*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.126B Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2003 and 2004**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	29.3	28.8	5.9	5.6	2.6	2.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	28.1	28.6	4.8	4.6	2.0	2.0
New England	32.6	31.3	6.7 <sup>a</sup>	4.5	2.1	1.9
Middle Atlantic	26.5	27.7	4.1	4.6	1.9	2.1
Midwest	27.9	26.9	5.3	5.1	2.3	2.2
East North Central	28.0	27.9	5.2	5.5	2.3	2.4
West North Central	27.5	24.6	5.4	4.2	2.3	1.7
South	26.2	27.8	5.8	5.5	2.6	2.6
South Atlantic	26.7	29.4	6.0	5.6	2.6	2.7
East South Central	24.4	24.5	4.9	5.8	2.2	3.0
West South Central	26.1	27.0	6.0	4.9	2.9	2.2
West	36.8 <sup>b</sup>	32.2	7.6	7.1	3.6 <sup>a</sup>	2.5
Mountain	36.8	33.1	8.3	7.5	3.7	3.2
Pacific	36.8 <sup>b</sup>	31.8	7.4	6.9	3.5 <sup>a</sup>	2.1
<b>COUNTY TYPE</b>						
Large Metro	31.3	30.4	6.2	5.8	2.8	2.4
Small Metro	28.8	28.8	6.1	5.7	2.6	2.7
250K – 1 Mil. Pop.	28.9	29.6	6.0	6.0	2.4	2.8
< 250K Pop.	28.6	27.3	6.4	5.0	3.1	2.4
Nonmetro	24.2	23.4	4.5	4.6	2.0	1.8
Urbanized	26.2	26.2	5.0	5.1	2.0	2.1
Less Urbanized	23.2	21.4	4.7	4.4	2.4	1.6
Completely Rural	20.1	22.1	2.0	3.3	0.6	1.2

\*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.



**Table 1.127A Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	65,809	65,302	16,954	16,389	7,432	6,920
<b>GEOGRAPHIC DIVISION</b>						
Northeast	12,155	12,351	2,875	2,820	1,189	1,245
New England	3,721	3,539	993 <sup>a</sup>	763	350	344
Middle Atlantic	8,434	8,812	1,881	2,057	839	900
Midwest	14,286	13,905	3,524	3,508	1,513	1,453
East North Central	10,062	10,051	2,439	2,601	1,068	1,091
West North Central	4,224	3,854	1,085	907	445	361
South	21,169 <sup>a</sup>	22,630	5,850	5,708	2,652	2,662
South Atlantic	11,263 <sup>a</sup>	12,399	3,081	2,969	1,379	1,394
East South Central	3,349	3,449	865	1,003	375	495
West South Central	6,556	6,783	1,904	1,736	898	772
West	18,200 <sup>a</sup>	16,415	4,706	4,353	2,078 <sup>a</sup>	1,560
Mountain	5,359	4,994	1,465	1,397	627	560
Pacific	12,840 <sup>a</sup>	11,421	3,241	2,956	1,451 <sup>a</sup>	1,000
<b>COUNTY TYPE</b>						
Large Metro	36,656	36,270	9,207	8,771	4,095	3,542
Small Metro	19,695	19,752	5,359	5,227	2,283	2,392
250K – 1 Mil. Pop.	12,966	13,197	3,439	3,539	1,392	1,601
< 250K Pop.	6,729	6,555	1,919	1,688	890	791
Nonmetro	9,459	9,280	2,389	2,391	1,055	985
Urbanized	4,560	4,266	1,190	1,109	514	468
Less Urbanized	4,125	4,254	1,087	1,117	508	450
Completely Rural	774	760	111	164	33	67

\*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.127B Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2003 and 2004**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	30.9	30.3	8.0	7.6	3.5	3.2
<b>GEOGRAPHIC DIVISION</b>						
Northeast	29.8	30.2	7.0	6.9	2.9	3.0
New England	34.7	32.9	9.3	7.1	3.3	3.2
Middle Atlantic	28.0	29.2	6.2	6.8	2.8	3.0
Midwest	29.7	28.7	7.3	7.2	3.1	3.0
East North Central	29.9	29.6	7.2	7.7	3.2	3.2
West North Central	29.3	26.4	7.5	6.2	3.1	2.5
South	27.9	29.3	7.7	7.4	3.5	3.5
South Atlantic	28.2	30.5	7.7	7.3	3.5	3.4
East South Central	26.2	26.6	6.8	7.7	2.9	3.8
West South Central	28.3	28.8	8.2	7.4	3.9	3.3
West	38.0 <sup>b</sup>	33.7	9.8	8.9	4.3 <sup>a</sup>	3.2
Mountain	38.6	35.1	10.6	9.8	4.5	3.9
Pacific	37.7 <sup>b</sup>	33.2	9.5	8.6	4.3 <sup>a</sup>	2.9
<b>COUNTY TYPE</b>						
Large Metro	32.5	31.5	8.2	7.6	3.6 <sup>a</sup>	3.1
Small Metro	30.8	30.9	8.4	8.2	3.6	3.7
250K – 1 Mil. Pop.	30.9	31.4	8.2	8.4	3.3	3.8
< 250K Pop.	30.5	29.9	8.7	7.7	4.0	3.6
Nonmetro	26.3	25.5	6.7	6.6	2.9	2.7
Urbanized	28.5	28.4	7.4	7.4	3.2	3.1
Less Urbanized	25.4	23.4	6.7	6.2	3.1	2.5
Completely Rural	21.2	23.8	3.0	5.2	0.9	2.1

\*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.