

Table 3.1C Perceived Risk and Availability of Substances, by Age Group: Standard Errors of Numbers in Thousands, 2003 and 2004

Risk/Availability ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2003	2004	2003	2004	2003	2004	2003	2004
PERCEPTIONS OF GREAT RISK								
Cigarettes								
Smoke One or More Packs Per Day	829	720	104	104	138	134	795	694
Marijuana								
Smoke Once a Month	897	897	101	109	124	137	861	863
Smoke Once or Twice a Week	860	905	109	112	150	150	817	858
Cocaine								
Use Once a Month	759	735	112	106	133	138	721	702
Use Once or Twice a Week	539	488	88	92	98	107	507	474
Heroin								
Try Once or Twice	595	619	108	103	118	117	565	589
Use Once or Twice a Week	378	385	85	86	74	75	353	368
LSD								
Try Once or Twice	732	765	111	105	136	140	700	726
Use Once or Twice a Week	497	483	99	88	97	101	465	472
Alcohol								
Four or Five Drinks Nearly Every Day	770	792	112	112	144	143	753	773
Five or More Drinks Once or Twice a Week	831	855	103	101	134	149	818	834
PERCEIVED AVAILABILITY								
Fairly or Very Easy to Obtain								
Marijuana	890	880	106	108	119	139	861	855
Cocaine	768	738	90	102	144	147	736	718
Crack	750	717	91	100	142	134	733	703
Heroin	651	614	79	80	109	112	639	591
LSD	612	604	84	84	126	113	598	586
Approached by Someone Selling Drugs in the Past Month	376	380	83	85	116	122	328	341

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.1D Perceived Risk and Availability of Substances, by Age Group: Standard Errors of Percentages, 2003 and 2004

Risk/Availability ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2003	2004	2003	2004	2003	2004	2003	2004
PERCEPTIONS OF GREAT RISK								
Cigarettes								
Smoke One or More Packs Per Day	0.35	0.30	0.42	0.42	0.44	0.42	0.44	0.38
Marijuana								
Smoke Once a Month	0.38	0.38	0.41	0.44	0.39	0.43	0.48	0.48
Smoke Once or Twice a Week	0.37	0.38	0.44	0.45	0.48	0.47	0.46	0.48
Cocaine								
Use Once a Month	0.32	0.31	0.46	0.43	0.42	0.43	0.40	0.39
Use Once or Twice a Week	0.23	0.21	0.36	0.37	0.31	0.33	0.28	0.26
Heroin								
Try Once or Twice	0.25	0.26	0.44	0.42	0.38	0.37	0.32	0.33
Use Once or Twice a Week	0.16	0.16	0.35	0.35	0.24	0.24	0.20	0.20
LSD								
Try Once or Twice	0.31	0.33	0.46	0.44	0.44	0.44	0.39	0.40
Use Once or Twice a Week	0.21	0.21	0.41	0.36	0.31	0.32	0.26	0.26
Alcohol								
Four or Five Drinks Nearly Every Day	0.33	0.33	0.45	0.45	0.45	0.45	0.42	0.43
Five or More Drinks Once or Twice a Week	0.35	0.36	0.42	0.40	0.42	0.46	0.45	0.46
PERCEIVED AVAILABILITY								
Fairly or Very Easy to Obtain								
Marijuana	0.39	0.38	0.44	0.44	0.38	0.44	0.50	0.49
Cocaine	0.34	0.33	0.38	0.42	0.47	0.47	0.43	0.42
Crack	0.33	0.32	0.38	0.41	0.47	0.43	0.43	0.41
Heroin	0.29	0.27	0.33	0.33	0.36	0.36	0.38	0.35
LSD	0.27	0.27	0.36	0.35	0.41	0.37	0.35	0.34
Approached by Someone Selling Drugs in the Past Month	0.16	0.16	0.33	0.34	0.37	0.38	0.18	0.19

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.1P Perceived Risk and Availability of Substances, by Age Group: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Risk/Availability ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2003	2004	2003	2004	2003	2004	2003	2004
PERCEPTIONS OF GREAT RISK								
Cigarettes								
Smoke One or More Packs Per Day	0.0000		0.0000		0.0000		0.0000	
Marijuana								
Smoke Once a Month	0.9284		0.9528		0.1531		0.9093	
Smoke Once or Twice a Week	0.5046		0.6404		0.0430		0.2664	
Cocaine								
Use Once a Month	0.6110		0.0039		0.1095		0.8206	
Use Once or Twice a Week	0.9680		0.0763		0.0865		0.5294	
Heroin								
Try Once or Twice	0.0095		0.0041		0.3308		0.0429	
Use Once or Twice a Week	0.5306		0.0183		0.4535		0.9759	
LSD								
Try Once or Twice	0.0040		0.2164		0.5441		0.0055	
Use Once or Twice a Week	0.8303		0.3385		0.4256		0.8266	
Alcohol								
Four or Five Drinks Nearly Every Day	0.3614		0.7462		0.3020		0.2409	
Five or More Drinks Once or Twice a Week	0.0727		0.4992		0.4945		0.1018	
PERCEIVED AVAILABILITY								
Fairly or Very Easy to Obtain								
Marijuana	0.0221		0.0273		0.0009		0.1097	
Cocaine	0.3345		0.2674		0.6997		0.4353	
Crack	0.1263		0.0457		0.5581		0.2508	
Heroin	0.2377		0.0063		0.0481		0.6341	
LSD	0.0030		0.1927		0.0000		0.0786	
Approached by Someone Selling Drugs in the Past Month	0.4374		0.6600		0.1542		0.6599	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.2C Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Smoke One or More Packs of Cigarettes Per Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	104	104	104	104
AGE				
12-13	65	58	65	58
14-15	59	65	59	65
16-17	58	58	58	58
GENDER				
Male	72	77	72	77
Female	69	68	69	68
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	95	90	95	90
White	79	79	79	79
Black or African American	40	41	40	41
American Indian or Alaska Native	8	8	8	8
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	27	21	27	21
Two or More Races	11	12	11	12
Hispanic or Latino	46	53	46	53
COUNTY TYPE				
Large Metro	78	82	78	82
Small Metro	51	50	51	50
Nonmetro	39	39	39	39
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	51	52	43	44
Marijuana	46	43	38	37
Illicit Drug Other Than Marijuana ²	35	38	32	32
Cigarettes	51	51	46	49
Alcohol	63	66	51	49
Binge Alcohol ³	52	56	40	44
PAST YEAR				
Illicit Drug ²	69	72	57	52
Marijuana	60	61	49	47
Illicit Drug Other Than Marijuana ²	56	56	48	43
Cigarettes	64	65	53	55
Alcohol	79	86	63	60
LIFETIME				
Illicit Drug ²	81	78	61	59
Marijuana	69	68	55	51
Illicit Drug Other Than Marijuana ²	71	67	56	53
Cigarettes	82	80	64	62
Alcohol	88	89	64	64

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.2D Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Smoke One or More Packs of Cigarettes Per Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.42	0.42	0.42	0.42
AGE				
12-13	0.78	0.71	0.78	0.71
14-15	0.71	0.75	0.71	0.75
16-17	0.71	0.72	0.71	0.72
GENDER				
Male	0.57	0.60	0.57	0.60
Female	0.57	0.56	0.57	0.56
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.46	0.43	0.46	0.43
White	0.51	0.51	0.51	0.51
Black or African American	1.08	1.09	1.08	1.09
American Indian or Alaska Native	4.71	4.76	4.71	4.76
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.78	2.15	2.78	2.15
Two or More Races	2.77	2.76	2.77	2.76
Hispanic or Latino	1.15	1.28	1.15	1.28
COUNTY TYPE				
Large Metro	0.59	0.62	0.59	0.62
Small Metro	0.70	0.66	0.70	0.66
Nonmetro	0.92	0.92	0.92	0.92
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.32	0.31	0.49	0.54
Marijuana	0.29	0.25	0.43	0.46
Illicit Drug Other Than Marijuana ²	0.22	0.22	0.36	0.40
Cigarettes	0.32	0.30	0.52	0.61
Alcohol	0.39	0.39	0.57	0.60
Binge Alcohol ³	0.32	0.33	0.46	0.55
PAST YEAR				
Illicit Drug ²	0.43	0.43	0.64	0.64
Marijuana	0.38	0.36	0.55	0.57
Illicit Drug Other Than Marijuana ²	0.35	0.33	0.54	0.53
Cigarettes	0.40	0.39	0.60	0.67
Alcohol	0.49	0.51	0.71	0.74
LIFETIME				
Illicit Drug ²	0.51	0.46	0.69	0.73
Marijuana	0.43	0.40	0.62	0.63
Illicit Drug Other Than Marijuana ²	0.45	0.40	0.63	0.65
Cigarettes	0.51	0.47	0.72	0.77
Alcohol	0.55	0.53	0.72	0.79

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.2P Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day (II), among Persons Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

(I) Demographic Characteristic and (II) Substance Use	Smoke One or More Packs of Cigarettes Per Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.0000		0.0000	
AGE				
12-13	0.0002		0.0002	
14-15	0.0043		0.0043	
16-17	0.0013		0.0013	
GENDER				
Male	0.0020		0.0020	
Female	0.0000		0.0000	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.0000		0.0000	
White	0.0000		0.0000	
Black or African American	0.0469		0.0469	
American Indian or Alaska Native	0.0931		0.0931	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.0063		0.0063	
Two or More Races	0.8842		0.8842	
Hispanic or Latino	0.3258		0.3258	
COUNTY TYPE				
Large Metro	0.0002		0.0002	
Small Metro	0.0046		0.0046	
Nonmetro	0.0001		0.0001	
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.1479		0.8650	
Marijuana	0.6057		0.8745	
Illicit Drug Other Than Marijuana ²	0.1380		0.9434	
Cigarettes	0.3374		0.2691	
Alcohol	0.3959		0.3087	
Binge Alcohol ³	0.2010		0.4361	
PAST YEAR				
Illicit Drug ²	0.5145		0.3696	
Marijuana	0.7920		0.5810	
Illicit Drug Other Than Marijuana ²	0.6021		0.6950	
Cigarettes	0.5374		0.9107	
Alcohol	0.5316		0.1581	
LIFETIME				
Illicit Drug ²	0.7880		0.2942	
Marijuana	0.8757		0.3569	
Illicit Drug Other Than Marijuana ²	0.3770		0.6336	
Cigarettes	0.0552		0.0987	
Alcohol	0.9835		0.0853	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.3C Risk Perceptions of Smoking Marijuana Once a Month, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking Marijuana Once a Month (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Smoke Marijuana Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	101	109	101	109
AGE				
12-13	58	60	58	60
14-15	57	62	57	62
16-17	55	56	55	56
GENDER				
Male	70	73	70	73
Female	75	73	75	73
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	93	92	93	92
White	77	77	77	77
Black or African American	39	40	39	40
American Indian or Alaska Native	6	7	6	7
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	23	26	23	26
Two or More Races	12	12	12	12
Hispanic or Latino	43	56	43	56
COUNTY TYPE				
Large Metro	77	84	77	84
Small Metro	50	56	50	56
Nonmetro	40	37	40	37
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	27	25	59	63
Marijuana	16	16	55	54
Illicit Drug Other Than Marijuana ²	22	21	43	46
Cigarettes	26	29	63	68
Alcohol	36	38	69	73
Binge Alcohol ³	26	29	60	66
PAST YEAR				
Illicit Drug ²	39	39	77	76
Marijuana	26	27	70	69
Illicit Drug Other Than Marijuana ²	34	32	67	66
Cigarettes	37	39	75	75
Alcohol	50	57	81	83
LIFETIME				
Illicit Drug ²	51	47	85	82
Marijuana	32	32	78	75
Illicit Drug Other Than Marijuana ²	49	43	78	76
Cigarettes	53	50	87	87
Alcohol	58	60	87	89

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.3D Risk Perceptions of Smoking Marijuana Once a Month, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking Marijuana Once a Month (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Smoke Marijuana Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.41	0.44	0.41	0.44
AGE				
12-13	0.71	0.74	0.71	0.74
14-15	0.70	0.72	0.70	0.72
16-17	0.67	0.69	0.67	0.69
GENDER				
Male	0.56	0.58	0.56	0.58
Female	0.62	0.60	0.62	0.60
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.45	0.44	0.45	0.44
White	0.50	0.50	0.50	0.50
Black or African American	1.06	1.06	1.06	1.06
American Indian or Alaska Native	3.77	4.50	3.77	4.50
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.46	2.73	2.46	2.73
Two or More Races	3.18	2.72	3.18	2.72
Hispanic or Latino	1.08	1.34	1.08	1.34
COUNTY TYPE				
Large Metro	0.59	0.64	0.59	0.64
Small Metro	0.69	0.74	0.69	0.74
Nonmetro	0.95	0.89	0.95	0.89
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.32	0.29	0.37	0.39
Marijuana	0.19	0.18	0.34	0.33
Illicit Drug Other Than Marijuana ²	0.26	0.24	0.27	0.28
Cigarettes	0.30	0.33	0.39	0.42
Alcohol	0.41	0.44	0.43	0.45
Binge Alcohol ³	0.31	0.33	0.37	0.41
PAST YEAR				
Illicit Drug ²	0.45	0.44	0.48	0.47
Marijuana	0.30	0.31	0.43	0.42
Illicit Drug Other Than Marijuana ²	0.40	0.37	0.42	0.40
Cigarettes	0.44	0.45	0.47	0.46
Alcohol	0.58	0.65	0.51	0.51
LIFETIME				
Illicit Drug ²	0.59	0.54	0.53	0.50
Marijuana	0.37	0.37	0.48	0.47
Illicit Drug Other Than Marijuana ²	0.57	0.49	0.49	0.47
Cigarettes	0.61	0.58	0.54	0.53
Alcohol	0.67	0.69	0.54	0.55

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.3P Risk Perceptions of Smoking Marijuana Once a Month, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking Marijuana Once a Month (II), among Persons Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

(I) Demographic Characteristic and (II) Substance Use	Smoke Marijuana Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.9528		0.9528	
AGE				
12-13	0.5682		0.5682	
14-15	0.9302		0.9302	
16-17	0.5580		0.5580	
GENDER				
Male	0.4521		0.4521	
Female	0.4267		0.4267	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.7214		0.7214	
White	0.5399		0.5399	
Black or African American	0.7048		0.7048	
American Indian or Alaska Native	0.7316		0.7316	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.9639		0.9639	
Two or More Races	0.9780		0.9780	
Hispanic or Latino	0.4414		0.4414	
COUNTY TYPE				
Large Metro	0.9790		0.9790	
Small Metro	0.1440		0.1440	
Nonmetro	0.0193		0.0193	
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.0206		0.4279	
Marijuana	0.1089		0.6296	
Illicit Drug Other Than Marijuana ²	0.0858		0.4511	
Cigarettes	0.6950		0.2972	
Alcohol	0.6569		0.9702	
Binge Alcohol ³	0.6544		0.3314	
PAST YEAR				
Illicit Drug ²	0.0130		0.6189	
Marijuana	0.2787		0.4648	
Illicit Drug Other Than Marijuana ²	0.0121		0.9323	
Cigarettes	0.7469		0.2301	
Alcohol	0.5247		0.7030	
LIFETIME				
Illicit Drug ²	0.3326		0.7460	
Marijuana	0.2640		0.4134	
Illicit Drug Other Than Marijuana ²	0.5714		0.7144	
Cigarettes	0.0511		0.0200	
Alcohol	0.0723		0.6731	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.4C Risk Perceptions of Smoking Marijuana Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking Marijuana Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Smoke Marijuana Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	109	112	109	112
AGE				
12-13	58	62	58	62
14-15	59	65	59	65
16-17	64	62	64	62
GENDER				
Male	74	79	74	79
Female	79	73	79	73
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	99	99	99	99
White	87	80	87	80
Black or African American	38	47	38	47
American Indian or Alaska Native	9	9	9	9
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	29	28	29	28
Two or More Races	12	13	12	13
Hispanic or Latino	49	58	49	58
COUNTY TYPE				
Large Metro	86	84	86	84
Small Metro	55	61	55	61
Nonmetro	40	44	40	44
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	28	31	56	54
Marijuana	19	19	53	51
Illicit Drug Other Than Marijuana ²	24	25	40	41
Cigarettes	32	36	58	64
Alcohol	45	48	66	66
Binge Alcohol ³	32	36	55	61
PAST YEAR				
Illicit Drug ²	48	50	67	67
Marijuana	34	35	64	63
Illicit Drug Other Than Marijuana ²	42	41	62	60
Cigarettes	46	47	67	71
Alcohol	64	70	74	74
LIFETIME				
Illicit Drug ²	61	59	73	69
Marijuana	42	44	70	65
Illicit Drug Other Than Marijuana ²	56	53	69	69
Cigarettes	64	62	75	74
Alcohol	73	77	73	71

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.4D Risk Perceptions of Smoking Marijuana Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking Marijuana Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Smoke Marijuana Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.44	0.45	0.44	0.45
AGE				
12-13	0.70	0.75	0.70	0.75
14-15	0.72	0.75	0.72	0.75
16-17	0.77	0.77	0.77	0.77
GENDER				
Male	0.58	0.63	0.58	0.63
Female	0.65	0.59	0.65	0.59
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.48	0.48	0.48	0.48
White	0.56	0.52	0.56	0.52
Black or African American	1.02	1.26	1.02	1.26
American Indian or Alaska Native	4.89	5.17	4.89	5.17
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.99	2.91	2.99	2.91
Two or More Races	2.97	3.07	2.97	3.07
Hispanic or Latino	1.24	1.40	1.24	1.40
COUNTY TYPE				
Large Metro	0.65	0.64	0.65	0.64
Small Metro	0.76	0.81	0.76	0.81
Nonmetro	0.94	1.06	0.94	1.06
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.21	0.23	0.50	0.48
Marijuana	0.14	0.14	0.47	0.45
Illicit Drug Other Than Marijuana ²	0.18	0.18	0.36	0.37
Cigarettes	0.24	0.26	0.51	0.57
Alcohol	0.34	0.35	0.59	0.58
Binge Alcohol ³	0.24	0.26	0.49	0.54
PAST YEAR				
Illicit Drug ²	0.36	0.37	0.60	0.59
Marijuana	0.25	0.25	0.57	0.56
Illicit Drug Other Than Marijuana ²	0.31	0.30	0.55	0.53
Cigarettes	0.34	0.35	0.60	0.63
Alcohol	0.48	0.52	0.65	0.65
LIFETIME				
Illicit Drug ²	0.46	0.43	0.64	0.61
Marijuana	0.31	0.33	0.62	0.58
Illicit Drug Other Than Marijuana ²	0.42	0.39	0.61	0.61
Cigarettes	0.47	0.46	0.66	0.66
Alcohol	0.54	0.57	0.65	0.63

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.4P Risk Perceptions of Smoking Marijuana Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking Marijuana Once or Twice a Week (II), among Persons Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

(I) Demographic Characteristic and (II) Substance Use	Smoke Marijuana Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.6404		0.6404	
AGE				
12-13	0.9148		0.9148	
14-15	0.7647		0.7647	
16-17	0.6781		0.6781	
GENDER				
Male	0.2476		0.2476	
Female	0.0654		0.0654	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.6477		0.6477	
White	0.9830		0.9830	
Black or African American	0.2228		0.2228	
American Indian or Alaska Native	0.2695		0.2695	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.9171		0.9171	
Two or More Races	0.5819		0.5819	
Hispanic or Latino	0.8634		0.8634	
COUNTY TYPE				
Large Metro	0.3161		0.3161	
Small Metro	0.1969		0.1969	
Nonmetro	0.2031		0.2031	
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.3080		0.1506	
Marijuana	0.3503		0.5410	
Illicit Drug Other Than Marijuana ²	0.6430		0.1567	
Cigarettes	0.0847		0.0806	
Alcohol	0.7282		0.6631	
Binge Alcohol ³	0.1378		0.5963	
PAST YEAR				
Illicit Drug ²	0.2301		0.3300	
Marijuana	0.7295		0.3463	
Illicit Drug Other Than Marijuana ²	0.1929		0.8110	
Cigarettes	0.5494		0.0886	
Alcohol	0.2793		0.0661	
LIFETIME				
Illicit Drug ²	0.9639		0.3893	
Marijuana	0.5838		0.0981	
Illicit Drug Other Than Marijuana ²	0.6403		0.5862	
Cigarettes	0.2520		0.0022	
Alcohol	0.9937		0.0706	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.5C Risk Perceptions of Using Cocaine Once a Month, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Cocaine Once a Month (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Use Cocaine Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	112	106	112	106
AGE				
12-13	60	62	60	62
14-15	63	70	63	70
16-17	62	60	62	60
GENDER				
Male	74	72	74	72
Female	83	74	83	74
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	101	93	101	93
White	82	79	82	79
Black or African American	43	48	43	48
American Indian or Alaska Native	7	7	7	7
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	26	25	26	25
Two or More Races	13	12	13	12
Hispanic or Latino	48	47	48	47
COUNTY TYPE				
Large Metro	87	78	87	78
Small Metro	55	59	55	59
Nonmetro	42	39	42	39
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	49	46	50	47
Marijuana	43	40	41	40
Illicit Drug Other Than Marijuana ²	32	32	38	37
Cigarettes	52	52	48	48
Alcohol	59	59	53	59
Binge Alcohol ³	48	51	44	50
PAST YEAR				
Illicit Drug ²	65	65	61	58
Marijuana	55	57	52	51
Illicit Drug Other Than Marijuana ²	51	48	54	51
Cigarettes	65	63	58	59
Alcohol	74	78	70	73
LIFETIME				
Illicit Drug ²	76	71	69	66
Marijuana	64	68	58	55
Illicit Drug Other Than Marijuana ²	65	58	66	61
Cigarettes	80	74	70	70
Alcohol	79	80	74	75

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.5D Risk Perceptions of Using Cocaine Once a Month, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Cocaine Once a Month (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Use Cocaine Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.46	0.43	0.46	0.43
AGE				
12-13	0.73	0.76	0.73	0.76
14-15	0.78	0.82	0.78	0.82
16-17	0.76	0.74	0.76	0.74
GENDER				
Male	0.59	0.57	0.59	0.57
Female	0.69	0.61	0.69	0.61
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.49	0.45	0.49	0.45
White	0.54	0.52	0.54	0.52
Black or African American	1.18	1.29	1.18	1.29
American Indian or Alaska Native	4.16	4.32	4.16	4.32
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.68	2.52	2.68	2.52
Two or More Races	3.26	2.74	3.26	2.74
Hispanic or Latino	1.20	1.13	1.20	1.13
COUNTY TYPE				
Large Metro	0.66	0.60	0.66	0.60
Small Metro	0.76	0.78	0.76	0.78
Nonmetro	1.00	0.95	1.00	0.95
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.39	0.37	0.42	0.38
Marijuana	0.34	0.32	0.35	0.32
Illicit Drug Other Than Marijuana ²	0.25	0.26	0.32	0.30
Cigarettes	0.42	0.42	0.40	0.38
Alcohol	0.47	0.48	0.45	0.47
Binge Alcohol ³	0.38	0.42	0.37	0.40
PAST YEAR				
Illicit Drug ²	0.51	0.52	0.51	0.46
Marijuana	0.44	0.46	0.43	0.40
Illicit Drug Other Than Marijuana ²	0.40	0.39	0.46	0.41
Cigarettes	0.52	0.51	0.48	0.48
Alcohol	0.59	0.64	0.59	0.58
LIFETIME				
Illicit Drug ²	0.60	0.58	0.58	0.53
Marijuana	0.51	0.55	0.49	0.44
Illicit Drug Other Than Marijuana ²	0.51	0.47	0.55	0.49
Cigarettes	0.63	0.60	0.59	0.56
Alcohol	0.62	0.65	0.62	0.60

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.5P Risk Perceptions of Using Cocaine Once a Month, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Cocaine Once a Month (II), among Persons Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

(I) Demographic Characteristic and (II) Substance Use	Use Cocaine Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.0039		0.0039	
AGE				
12-13	0.4720		0.4720	
14-15	0.0044		0.0044	
16-17	0.2183		0.2183	
GENDER				
Male	0.0023		0.0023	
Female	0.2404		0.2404	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.0063		0.0063	
White	0.0030		0.0030	
Black or African American	0.2812		0.2812	
American Indian or Alaska Native	0.0380		0.0380	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.9102		0.9102	
Two or More Races	0.9729		0.9729	
Hispanic or Latino	0.3545		0.3545	
COUNTY TYPE				
Large Metro	0.1135		0.1135	
Small Metro	0.0000		0.0000	
Nonmetro	0.1849		0.1849	
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.0674		0.5857	
Marijuana	0.3626		0.7648	
Illicit Drug Other Than Marijuana ²	0.1987		0.3865	
Cigarettes	0.5421		0.7560	
Alcohol	0.9715		0.7600	
Binge Alcohol ³	0.1395		0.9559	
PAST YEAR				
Illicit Drug ²	0.1956		0.4182	
Marijuana	0.3829		0.6584	
Illicit Drug Other Than Marijuana ²	0.2918		0.4893	
Cigarettes	0.3496		0.6524	
Alcohol	0.9070		0.4809	
LIFETIME				
Illicit Drug ²	0.7631		0.3532	
Marijuana	0.6661		0.2972	
Illicit Drug Other Than Marijuana ²	0.7583		0.5276	
Cigarettes	0.0226		0.0742	
Alcohol	0.3751		0.3745	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.6C Risk Perceptions of Using Cocaine Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Cocaine Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Use Cocaine Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	88	92	88	92
AGE				
12-13	54	56	54	56
14-15	49	53	49	53
16-17	44	47	44	47
GENDER				
Male	57	63	57	63
Female	63	62	63	62
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	73	76	73	76
White	57	58	57	58
Black or African American	36	39	36	39
American Indian or Alaska Native	5	7	5	7
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	27	26	27	26
Two or More Races	8	9	8	9
Hispanic or Latino	47	47	47	47
COUNTY TYPE				
Large Metro	68	67	68	67
Small Metro	49	58	49	58
Nonmetro	32	32	32	32
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	58	60	32	36
Marijuana	52	50	26	30
Illicit Drug Other Than Marijuana ²	38	41	26	29
Cigarettes	62	66	33	36
Alcohol	72	74	36	38
Binge Alcohol ³	59	64	28	33
PAST YEAR				
Illicit Drug ²	82	80	37	42
Marijuana	69	70	32	35
Illicit Drug Other Than Marijuana ²	67	60	35	37
Cigarettes	80	78	39	44
Alcohol	90	93	46	47
LIFETIME				
Illicit Drug ²	96	88	43	47
Marijuana	81	78	35	38
Illicit Drug Other Than Marijuana ²	84	73	42	44
Cigarettes	97	91	44	48
Alcohol	98	98	46	49

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.6D Risk Perceptions of Using Cocaine Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Cocaine Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Use Cocaine Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.36	0.37	0.36	0.37
AGE				
12-13	0.66	0.68	0.66	0.68
14-15	0.60	0.62	0.60	0.62
16-17	0.54	0.58	0.54	0.58
GENDER				
Male	0.45	0.50	0.45	0.50
Female	0.52	0.51	0.52	0.51
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.35	0.37	0.35	0.37
White	0.37	0.38	0.37	0.38
Black or African American	1.00	1.04	1.00	1.04
American Indian or Alaska Native	3.19	4.10	3.19	4.10
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.83	2.67	2.83	2.67
Two or More Races	2.11	2.11	2.11	2.11
Hispanic or Latino	1.19	1.13	1.19	1.13
COUNTY TYPE				
Large Metro	0.52	0.51	0.52	0.51
Small Metro	0.67	0.77	0.67	0.77
Nonmetro	0.76	0.76	0.76	0.76
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.29	0.30	0.67	0.72
Marijuana	0.26	0.25	0.55	0.59
Illicit Drug Other Than Marijuana ²	0.19	0.21	0.55	0.58
Cigarettes	0.31	0.33	0.70	0.72
Alcohol	0.36	0.37	0.76	0.76
Binge Alcohol ³	0.30	0.32	0.60	0.65
PAST YEAR				
Illicit Drug ²	0.41	0.40	0.77	0.83
Marijuana	0.35	0.35	0.66	0.69
Illicit Drug Other Than Marijuana ²	0.34	0.30	0.74	0.74
Cigarettes	0.40	0.40	0.82	0.87
Alcohol	0.45	0.47	0.96	0.93
LIFETIME				
Illicit Drug ²	0.49	0.45	0.90	0.93
Marijuana	0.41	0.40	0.74	0.76
Illicit Drug Other Than Marijuana ²	0.42	0.37	0.88	0.87
Cigarettes	0.49	0.46	0.92	0.95
Alcohol	0.49	0.50	0.96	0.97

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.6P Risk Perceptions of Using Cocaine Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Cocaine Once or Twice a Week (II), among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2004 Versus 2003

(I) Demographic Characteristic and (II) Substance Use	Use Cocaine Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.0763		0.0763	
AGE				
12-13	0.5201		0.5201	
14-15	0.2498		0.2498	
16-17	0.1764		0.1764	
GENDER				
Male	0.1378		0.1378	
Female	0.2856		0.2856	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.1276		0.1276	
White	0.3841		0.3841	
Black or African American	0.1805		0.1805	
American Indian or Alaska Native	0.4900		0.4900	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.6728		0.6728	
Two or More Races	0.6557		0.6557	
Hispanic or Latino	0.4599		0.4599	
COUNTY TYPE				
Large Metro	1.0000		1.0000	
Small Metro	0.0105		0.0105	
Nonmetro	0.6797		0.6797	
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.0417		0.9723	
Marijuana	0.2011		0.6563	
Illicit Drug Other Than Marijuana ²	0.0855		0.8074	
Cigarettes	0.4931		0.7269	
Alcohol	0.9591		0.4830	
Binge Alcohol ³	0.3259		0.8841	
PAST YEAR				
Illicit Drug ²	0.0919		0.8668	
Marijuana	0.2875		0.9511	
Illicit Drug Other Than Marijuana ²	0.1523		0.9647	
Cigarettes	0.3128		0.6751	
Alcohol	0.6514		0.0180	
LIFETIME				
Illicit Drug ²	0.4216		0.7652	
Marijuana	0.2808		0.8188	
Illicit Drug Other Than Marijuana ²	0.9637		0.6403	
Cigarettes	0.0167		0.0732	
Alcohol	0.5763		0.0506	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.7C Risk Perceptions of Trying Heroin Once or Twice, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Trying Heroin Once or Twice (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Try Heroin Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	108	103	108	103
AGE				
12-13	61	60	61	60
14-15	60	67	60	67
16-17	59	57	59	57
GENDER				
Male	73	74	73	74
Female	79	73	79	73
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	101	91	101	91
White	80	80	80	80
Black or African American	43	43	43	43
American Indian or Alaska Native	9	9	9	9
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	27	27	27	27
Two or More Races	12	12	12	12
Hispanic or Latino	44	47	44	47
COUNTY TYPE				
Large Metro	81	79	81	79
Small Metro	58	56	58	56
Nonmetro	41	36	41	36
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	55	55	37	38
Marijuana	51	47	30	32
Illicit Drug Other Than Marijuana ²	37	38	31	30
Cigarettes	59	58	37	40
Alcohol	65	65	46	48
Binge Alcohol ³	54	57	36	39
PAST YEAR				
Illicit Drug ²	72	71	54	49
Marijuana	65	64	42	42
Illicit Drug Other Than Marijuana ²	56	56	48	44
Cigarettes	69	70	49	47
Alcohol	80	80	60	64
LIFETIME				
Illicit Drug ²	82	77	63	57
Marijuana	74	72	48	45
Illicit Drug Other Than Marijuana ²	71	68	56	56
Cigarettes	83	80	61	57
Alcohol	82	82	65	67

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.7D Risk Perceptions of Trying Heroin Once or Twice, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Trying Heroin Once or Twice (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Try Heroin Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.44	0.42	0.44	0.42
AGE				
12-13	0.74	0.75	0.74	0.75
14-15	0.74	0.79	0.74	0.79
16-17	0.72	0.70	0.72	0.70
GENDER				
Male	0.59	0.59	0.59	0.59
Female	0.66	0.60	0.66	0.60
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.49	0.44	0.49	0.44
White	0.52	0.52	0.52	0.52
Black or African American	1.19	1.16	1.19	1.16
American Indian or Alaska Native	5.28	5.25	5.28	5.25
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.85	2.83	2.85	2.83
Two or More Races	3.05	2.68	3.05	2.68
Hispanic or Latino	1.12	1.14	1.12	1.14
COUNTY TYPE				
Large Metro	0.62	0.60	0.62	0.60
Small Metro	0.80	0.74	0.80	0.74
Nonmetro	0.99	0.86	0.99	0.86
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.38	0.39	0.37	0.36
Marijuana	0.35	0.34	0.30	0.30
Illicit Drug Other Than Marijuana ²	0.26	0.27	0.31	0.29
Cigarettes	0.41	0.41	0.36	0.37
Alcohol	0.45	0.46	0.46	0.45
Binge Alcohol ³	0.37	0.41	0.35	0.37
PAST YEAR				
Illicit Drug ²	0.50	0.51	0.53	0.46
Marijuana	0.45	0.45	0.41	0.40
Illicit Drug Other Than Marijuana ²	0.39	0.40	0.48	0.42
Cigarettes	0.48	0.49	0.48	0.44
Alcohol	0.55	0.57	0.60	0.61
LIFETIME				
Illicit Drug ²	0.57	0.54	0.62	0.54
Marijuana	0.51	0.51	0.47	0.42
Illicit Drug Other Than Marijuana ²	0.50	0.48	0.56	0.52
Cigarettes	0.58	0.57	0.61	0.53
Alcohol	0.57	0.59	0.64	0.63

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.7P Risk Perceptions of Trying Heroin Once or Twice, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Trying Heroin Once or Twice (II), among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2004 Versus 2003

(I) Demographic Characteristic and (II) Substance Use	Try Heroin Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.0041		0.0041	
AGE				
12-13	0.3247		0.3247	
14-15	0.0033		0.0033	
16-17	0.2943		0.2943	
GENDER				
Male	0.0142		0.0142	
Female	0.1283		0.1283	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.0076		0.0076	
White	0.0147		0.0147	
Black or African American	0.2192		0.2192	
American Indian or Alaska Native	0.9515		0.9515	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.5137		0.5137	
Two or More Races	0.3637		0.3637	
Hispanic or Latino	0.2605		0.2605	
COUNTY TYPE				
Large Metro	0.1495		0.1495	
Small Metro	0.0035		0.0035	
Nonmetro	0.6636		0.6636	
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.0573		0.7708	
Marijuana	0.1890		0.2596	
Illicit Drug Other Than Marijuana ²	0.1070		0.9276	
Cigarettes	0.2838		0.3034	
Alcohol	0.5729		0.5587	
Binge Alcohol ³	0.2173		0.6988	
PAST YEAR				
Illicit Drug ²	0.2606		0.7312	
Marijuana	0.4296		0.8270	
Illicit Drug Other Than Marijuana ²	0.2869		0.9599	
Cigarettes	0.4477		0.9385	
Alcohol	0.9535		0.7881	
LIFETIME				
Illicit Drug ²	0.3482		0.7460	
Marijuana	0.4902		0.9042	
Illicit Drug Other Than Marijuana ²	0.8839		0.7081	
Cigarettes	0.0135		0.2939	
Alcohol	0.5144		0.5441	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.8C Risk Perceptions of Using Heroin Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Heroin Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Use Heroin Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	85	86	85	86
AGE				
12-13	55	56	55	56
14-15	47	52	47	52
16-17	35	40	35	40
GENDER				
Male	57	64	57	64
Female	62	55	62	55
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	72	75	72	75
White	53	59	53	59
Black or African American	36	39	36	39
American Indian or Alaska Native	5	7	5	7
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	25	23	25	23
Two or More Races	9	9	9	9
Hispanic or Latino	42	43	42	43
COUNTY TYPE				
Large Metro	69	65	69	65
Small Metro	40	51	40	51
Nonmetro	30	31	30	31
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	62	63	25	28
Marijuana	56	53	20	22
Illicit Drug Other Than Marijuana ²	43	44	22	23
Cigarettes	67	67	26	27
Alcohol	77	75	29	32
Binge Alcohol ³	62	66	22	27
PAST YEAR				
Illicit Drug ²	83	82	33	35
Marijuana	74	72	25	26
Illicit Drug Other Than Marijuana ²	68	65	30	32
Cigarettes	84	80	32	34
Alcohol	94	93	39	41
LIFETIME				
Illicit Drug ²	96	88	39	39
Marijuana	85	79	29	28
Illicit Drug Other Than Marijuana ²	84	77	36	38
Cigarettes	100	92	39	41
Alcohol	100	98	43	42

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.8D Risk Perceptions of Using Heroin Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Heroin Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Use Heroin Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.35	0.35	0.35	0.35
AGE				
12-13	0.67	0.69	0.67	0.69
14-15	0.58	0.62	0.58	0.62
16-17	0.43	0.49	0.43	0.49
GENDER				
Male	0.45	0.51	0.45	0.51
Female	0.52	0.45	0.52	0.45
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.35	0.37	0.35	0.37
White	0.35	0.39	0.35	0.39
Black or African American	0.99	1.05	0.99	1.05
American Indian or Alaska Native	2.88	4.20	2.88	4.20
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.68	2.40	2.68	2.40
Two or More Races	2.19	2.15	2.19	2.15
Hispanic or Latino	1.05	1.05	1.05	1.05
COUNTY TYPE				
Large Metro	0.53	0.50	0.53	0.50
Small Metro	0.56	0.68	0.56	0.68
Nonmetro	0.72	0.75	0.72	0.75
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.31	0.31	0.59	0.60
Marijuana	0.28	0.26	0.47	0.47
Illicit Drug Other Than Marijuana ²	0.21	0.22	0.52	0.51
Cigarettes	0.33	0.33	0.60	0.60
Alcohol	0.38	0.37	0.67	0.69
Binge Alcohol ³	0.31	0.33	0.52	0.58
PAST YEAR				
Illicit Drug ²	0.41	0.41	0.78	0.77
Marijuana	0.36	0.36	0.59	0.57
Illicit Drug Other Than Marijuana ²	0.34	0.32	0.71	0.69
Cigarettes	0.42	0.40	0.75	0.74
Alcohol	0.46	0.46	0.90	0.89
LIFETIME				
Illicit Drug ²	0.47	0.44	0.90	0.85
Marijuana	0.42	0.39	0.67	0.62
Illicit Drug Other Than Marijuana ²	0.42	0.38	0.84	0.82
Cigarettes	0.49	0.46	0.91	0.90
Alcohol	0.49	0.49	1.02	0.91

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.8P Risk Perceptions of Using Heroin Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Heroin Once or Twice a Week (II), among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2004 Versus 2003

(I) Demographic Characteristic and (II) Substance Use	Use Heroin Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.0183		0.0183	
AGE				
12-13	0.6759		0.6759	
14-15	0.0217		0.0217	
16-17	0.1037		0.1037	
GENDER				
Male	0.0144		0.0144	
Female	0.4206		0.4206	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.0249		0.0249	
White	0.0171		0.0171	
Black or African American	0.3225		0.3225	
American Indian or Alaska Native	0.6068		0.6068	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.6345		0.6345	
Two or More Races	0.5836		0.5836	
Hispanic or Latino	0.4148		0.4148	
COUNTY TYPE				
Large Metro	0.3973		0.3973	
Small Metro	0.0099		0.0099	
Nonmetro	0.4476		0.4476	
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.1008		0.9938	
Marijuana	0.3571		0.6214	
Illicit Drug Other Than Marijuana ²	0.1170		0.9418	
Cigarettes	0.6817		0.6949	
Alcohol	0.9749		0.8394	
Binge Alcohol ³	0.3475		0.5095	
PAST YEAR				
Illicit Drug ²	0.1385		0.6931	
Marijuana	0.3769		0.7611	
Illicit Drug Other Than Marijuana ²	0.2564		0.7445	
Cigarettes	0.4527		0.6180	
Alcohol	0.8959		0.1879	
LIFETIME				
Illicit Drug ²	0.4047		0.9138	
Marijuana	0.3342		0.8796	
Illicit Drug Other Than Marijuana ²	0.9518		0.9479	
Cigarettes	0.0184		0.1049	
Alcohol	0.8148		0.0210	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.9C Risk Perceptions of Trying LSD Once or Twice, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Trying LSD Once or Twice (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Try LSD Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	111	105	111	105
AGE				
12-13	54	60	54	60
14-15	60	61	60	61
16-17	68	58	68	58
GENDER				
Male	75	74	75	74
Female	76	76	76	76
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	102	89	102	89
White	81	78	81	78
Black or African American	45	41	45	41
American Indian or Alaska Native	8	6	8	6
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	27	25	27	25
Two or More Races	12	11	12	11
Hispanic or Latino	47	52	47	52
COUNTY TYPE				
Large Metro	84	80	84	80
Small Metro	55	58	55	58
Nonmetro	42	38	42	38
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	48	47	46	45
Marijuana	43	41	40	39
Illicit Drug Other Than Marijuana ²	31	31	36	36
Cigarettes	52	53	48	47
Alcohol	60	60	51	55
Binge Alcohol ³	49	51	41	50
PAST YEAR				
Illicit Drug ²	66	64	59	56
Marijuana	56	58	51	49
Illicit Drug Other Than Marijuana ²	51	49	52	50
Cigarettes	62	63	60	57
Alcohol	73	76	68	70
LIFETIME				
Illicit Drug ²	76	73	65	63
Marijuana	66	70	57	54
Illicit Drug Other Than Marijuana ²	64	60	60	61
Cigarettes	79	76	70	65
Alcohol	77	79	73	70

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.9D Risk Perceptions of Trying LSD Once or Twice, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Trying LSD Once or Twice (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Try LSD Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.46	0.44	0.46	0.44
AGE				
12-13	0.69	0.77	0.69	0.77
14-15	0.76	0.73	0.76	0.73
16-17	0.83	0.73	0.83	0.73
GENDER				
Male	0.62	0.60	0.62	0.60
Female	0.64	0.64	0.64	0.64
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.51	0.44	0.51	0.44
White	0.54	0.52	0.54	0.52
Black or African American	1.25	1.13	1.25	1.13
American Indian or Alaska Native	5.00	3.83	5.00	3.83
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.93	2.73	2.93	2.73
Two or More Races	3.22	2.67	3.22	2.67
Hispanic or Latino	1.21	1.30	1.21	1.30
COUNTY TYPE				
Large Metro	0.66	0.63	0.66	0.63
Small Metro	0.77	0.79	0.77	0.79
Nonmetro	1.03	0.93	1.03	0.93
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.38	0.37	0.41	0.40
Marijuana	0.33	0.32	0.35	0.34
Illicit Drug Other Than Marijuana ²	0.25	0.25	0.32	0.32
Cigarettes	0.40	0.41	0.43	0.41
Alcohol	0.47	0.48	0.46	0.48
Binge Alcohol ³	0.38	0.40	0.37	0.43
PAST YEAR				
Illicit Drug ²	0.52	0.50	0.53	0.49
Marijuana	0.44	0.46	0.46	0.42
Illicit Drug Other Than Marijuana ²	0.40	0.39	0.47	0.44
Cigarettes	0.48	0.49	0.54	0.49
Alcohol	0.57	0.60	0.61	0.61
LIFETIME				
Illicit Drug ²	0.59	0.58	0.58	0.55
Marijuana	0.52	0.55	0.51	0.47
Illicit Drug Other Than Marijuana ²	0.50	0.48	0.53	0.53
Cigarettes	0.62	0.60	0.62	0.57
Alcohol	0.60	0.62	0.65	0.61

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.9P Risk Perceptions of Trying LSD Once or Twice, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Trying LSD Once or Twice (II), among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2004 Versus 2003

(I) Demographic Characteristic and (II) Substance Use	Try LSD Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.2164		0.2164	
AGE				
12-13	0.8191		0.8191	
14-15	0.0397		0.0397	
16-17	0.9237		0.9237	
GENDER				
Male	0.0566		0.0566	
Female	0.8117		0.8117	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.1403		0.1403	
White	0.0251		0.0251	
Black or African American	0.2897		0.2897	
American Indian or Alaska Native	0.7422		0.7422	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.7850		0.7850	
Two or More Races	0.6727		0.6727	
Hispanic or Latino	0.8444		0.8444	
COUNTY TYPE				
Large Metro	0.8685		0.8685	
Small Metro	0.0603		0.0603	
Nonmetro	0.9406		0.9406	
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.3793		0.1879	
Marijuana	0.7796		0.5005	
Illicit Drug Other Than Marijuana ²	0.2765		0.3824	
Cigarettes	0.2404		0.6938	
Alcohol	0.8498		0.6868	
Binge Alcohol ³	0.3747		0.6334	
PAST YEAR				
Illicit Drug ²	0.3607		0.1938	
Marijuana	0.5437		0.4956	
Illicit Drug Other Than Marijuana ²	0.7596		0.2545	
Cigarettes	0.3018		0.6831	
Alcohol	0.7113		0.8749	
LIFETIME				
Illicit Drug ²	0.5318		0.4292	
Marijuana	0.8095		0.1514	
Illicit Drug Other Than Marijuana ²	0.8317		0.6946	
Cigarettes	0.0208		0.0836	
Alcohol	0.3086		0.6029	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.10C Risk Perceptions of Using LSD Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using LSD Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Use LSD Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	99	88	99	88
AGE				
12-13	55	55	55	55
14-15	53	50	53	50
16-17	48	46	48	46
GENDER				
Male	66	67	66	67
Female	67	59	67	59
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	84	78	84	78
White	64	62	64	62
Black or African American	42	38	42	38
American Indian or Alaska Native	5	7	5	7
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	28	27	28	27
Two or More Races	8	12	8	12
Hispanic or Latino	44	43	44	43
COUNTY TYPE				
Large Metro	76	70	76	70
Small Metro	53	50	53	50
Nonmetro	33	30	33	30
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	58	60	34	35
Marijuana	51	51	27	29
Illicit Drug Other Than Marijuana ²	39	40	27	28
Cigarettes	64	61	34	36
Alcohol	73	72	37	40
Binge Alcohol ³	59	63	29	34
PAST YEAR				
Illicit Drug ²	80	79	43	42
Marijuana	67	70	34	34
Illicit Drug Other Than Marijuana ²	65	60	38	36
Cigarettes	79	76	40	41
Alcohol	87	90	47	49
LIFETIME				
Illicit Drug ²	93	84	48	48
Marijuana	78	77	39	37
Illicit Drug Other Than Marijuana ²	83	71	45	43
Cigarettes	96	88	48	48
Alcohol	94	96	49	51

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.10D Risk Perceptions of Using LSD Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using LSD Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Use LSD Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.41	0.36	0.41	0.36
AGE				
12-13	0.69	0.70	0.69	0.70
14-15	0.66	0.61	0.66	0.61
16-17	0.58	0.57	0.58	0.57
GENDER				
Male	0.54	0.55	0.54	0.55
Female	0.57	0.49	0.57	0.49
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.42	0.39	0.42	0.39
White	0.43	0.42	0.43	0.42
Black or African American	1.19	1.03	1.19	1.03
American Indian or Alaska Native	2.98	4.30	2.98	4.30
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	3.01	2.86	3.01	2.86
Two or More Races	2.16	2.71	2.16	2.71
Hispanic or Latino	1.13	1.07	1.13	1.07
COUNTY TYPE				
Large Metro	0.59	0.55	0.59	0.55
Small Metro	0.74	0.68	0.74	0.68
Nonmetro	0.81	0.74	0.81	0.74
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.31	0.32	0.60	0.61
Marijuana	0.28	0.28	0.49	0.51
Illicit Drug Other Than Marijuana ²	0.21	0.22	0.49	0.48
Cigarettes	0.34	0.33	0.60	0.62
Alcohol	0.40	0.39	0.67	0.70
Binge Alcohol ³	0.32	0.34	0.52	0.60
PAST YEAR				
Illicit Drug ²	0.43	0.43	0.78	0.74
Marijuana	0.36	0.38	0.62	0.59
Illicit Drug Other Than Marijuana ²	0.35	0.32	0.69	0.63
Cigarettes	0.43	0.41	0.73	0.71
Alcohol	0.47	0.49	0.84	0.87
LIFETIME				
Illicit Drug ²	0.50	0.46	0.86	0.83
Marijuana	0.42	0.42	0.70	0.65
Illicit Drug Other Than Marijuana ²	0.45	0.39	0.80	0.76
Cigarettes	0.52	0.48	0.87	0.84
Alcohol	0.51	0.52	0.88	0.89

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.10P Risk Perceptions of Using LSD Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using LSD Once or Twice a Week (II), among Persons Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

(I) Demographic Characteristic and (II) Substance Use	Use LSD Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.3385		0.3385	
AGE				
12-13	0.1786		0.1786	
14-15	0.5957		0.5957	
16-17	0.8085		0.8085	
GENDER				
Male	0.0344		0.0344	
Female	0.4586		0.4586	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.2229		0.2229	
White	0.1086		0.1086	
Black or African American	0.6411		0.6411	
American Indian or Alaska Native	0.1581		0.1581	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.9140		0.9140	
Two or More Races	0.3963		0.3963	
Hispanic or Latino	0.7266		0.7266	
COUNTY TYPE				
Large Metro	0.8077		0.8077	
Small Metro	0.1495		0.1495	
Nonmetro	0.8419		0.8419	
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.1572		0.6629	
Marijuana	0.4474		0.9882	
Illicit Drug Other Than Marijuana ²	0.2396		0.6085	
Cigarettes	0.4468		0.9520	
Alcohol	0.9779		0.6898	
Binge Alcohol ³	0.4713		0.5805	
PAST YEAR				
Illicit Drug ²	0.2309		0.3713	
Marijuana	0.6346		0.2443	
Illicit Drug Other Than Marijuana ²	0.3110		0.8152	
Cigarettes	0.3865		0.5754	
Alcohol	0.8426		0.2144	
LIFETIME				
Illicit Drug ²	0.5006		0.4258	
Marijuana	0.4212		0.3476	
Illicit Drug Other Than Marijuana ²	0.9484		0.7090	
Cigarettes	0.0234		0.0478	
Alcohol	0.6065		0.1259	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.11C Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Have Four or Five Drinks of Alcoholic Beverage Nearly Every Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	112	112	112	112
AGE				
12-13	63	65	63	65
14-15	59	64	59	64
16-17	61	58	61	58
GENDER				
Male	76	81	76	81
Female	75	71	75	71
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	99	97	99	97
White	85	82	85	82
Black or African American	40	39	40	39
American Indian or Alaska Native	8	*	8	*
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	24	26	24	26
Two or More Races	12	12	12	12
Hispanic or Latino	52	55	52	55
COUNTY TYPE				
Large Metro	87	89	87	89
Small Metro	53	56	53	56
Nonmetro	43	41	43	41
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	45	47	50	47
Marijuana	39	38	43	43
Illicit Drug Other Than Marijuana ²	34	33	35	36
Cigarettes	45	50	51	54
Alcohol	59	59	54	57
Binge Alcohol ³	43	48	48	51
PAST YEAR				
Illicit Drug ²	65	65	64	62
Marijuana	52	52	56	56
Illicit Drug Other Than Marijuana ²	54	51	55	52
Cigarettes	60	63	59	62
Alcohol	80	78	67	65
LIFETIME				
Illicit Drug ²	77	71	69	66
Marijuana	64	59	62	62
Illicit Drug Other Than Marijuana ²	67	60	63	62
Cigarettes	71	75	70	67
Alcohol	88	85	67	65

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.11D Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Have Four or Five Drinks of Alcoholic Beverage Nearly Every Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.45	0.45	0.45	0.45
AGE				
12-13	0.76	0.79	0.76	0.79
14-15	0.72	0.75	0.72	0.75
16-17	0.74	0.72	0.74	0.72
GENDER				
Male	0.60	0.64	0.60	0.64
Female	0.61	0.57	0.61	0.57
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.48	0.47	0.48	0.47
White	0.55	0.53	0.55	0.53
Black or African American	1.07	1.03	1.07	1.03
American Indian or Alaska Native	4.34	*	4.34	*
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.48	2.67	2.48	2.67
Two or More Races	2.96	2.76	2.96	2.76
Hispanic or Latino	1.28	1.30	1.28	1.30
COUNTY TYPE				
Large Metro	0.66	0.67	0.66	0.67
Small Metro	0.72	0.74	0.72	0.74
Nonmetro	1.02	0.97	1.02	0.97
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.30	0.30	0.52	0.50
Marijuana	0.25	0.24	0.45	0.45
Illicit Drug Other Than Marijuana ²	0.22	0.22	0.37	0.38
Cigarettes	0.29	0.33	0.53	0.57
Alcohol	0.39	0.38	0.57	0.60
Binge Alcohol ³	0.28	0.31	0.50	0.53
PAST YEAR				
Illicit Drug ²	0.42	0.42	0.67	0.65
Marijuana	0.34	0.34	0.58	0.58
Illicit Drug Other Than Marijuana ²	0.35	0.33	0.57	0.54
Cigarettes	0.39	0.41	0.62	0.64
Alcohol	0.53	0.51	0.70	0.68
LIFETIME				
Illicit Drug ²	0.50	0.46	0.73	0.69
Marijuana	0.42	0.38	0.65	0.64
Illicit Drug Other Than Marijuana ²	0.44	0.39	0.66	0.65
Cigarettes	0.47	0.48	0.74	0.71
Alcohol	0.58	0.55	0.71	0.68

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.11P Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day (II), among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2004 Versus 2003

(I) Demographic Characteristic and (II) Substance Use	Have Four or Five Drinks of Alcoholic Beverage Nearly Every Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.7462		0.7462	
AGE				
12-13	0.3717		0.3717	
14-15	0.3752		0.3752	
16-17	0.2154		0.2154	
GENDER				
Male	0.3637		0.3637	
Female	0.1349		0.1349	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.9440		0.9440	
White	0.8559		0.8559	
Black or African American	0.2703		0.2703	
American Indian or Alaska Native	*		*	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.9443		0.9443	
Two or More Races	0.5055		0.5055	
Hispanic or Latino	0.3894		0.3894	
COUNTY TYPE				
Large Metro	0.8903		0.8903	
Small Metro	0.4562		0.4562	
Nonmetro	0.8311		0.8311	
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.3745		0.1842	
Marijuana	0.4744		0.5945	
Illicit Drug Other Than Marijuana ²	0.8685		0.1101	
Cigarettes	0.6038		0.1711	
Alcohol	0.7701		0.9498	
Binge Alcohol ³	0.2898		0.5804	
PAST YEAR				
Illicit Drug ²	0.3705		0.2216	
Marijuana	0.6906		0.3191	
Illicit Drug Other Than Marijuana ²	0.4327		0.5580	
Cigarettes	0.4685		0.0205	
Alcohol	0.8363		0.4652	
LIFETIME				
Illicit Drug ²	0.5323		0.6041	
Marijuana	0.6168		0.3195	
Illicit Drug Other Than Marijuana ²	0.8128		0.8406	
Cigarettes	0.1389		0.0038	
Alcohol	0.7445		0.0740	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.12C Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Have Five or More Drinks of Alcoholic Beverage Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	103	101	103	101
AGE				
12-13	60	65	60	65
14-15	58	62	58	62
16-17	59	55	59	55
GENDER				
Male	72	78	72	78
Female	76	76	76	76
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	92	94	92	94
White	74	79	74	79
Black or African American	44	41	44	41
American Indian or Alaska Native	7	6	7	6
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	27	30	27	30
Two or More Races	13	11	13	11
Hispanic or Latino	49	52	49	52
COUNTY TYPE				
Large Metro	79	77	79	77
Small Metro	56	54	56	54
Nonmetro	41	42	41	42
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	33	36	57	57
Marijuana	27	27	51	52
Illicit Drug Other Than Marijuana ²	25	27	40	42
Cigarettes	37	39	60	61
Alcohol	44	46	69	68
Binge Alcohol ³	32	38	57	60
PAST YEAR				
Illicit Drug ²	47	51	76	72
Marijuana	39	40	66	67
Illicit Drug Other Than Marijuana ²	39	42	63	58
Cigarettes	49	48	71	72
Alcohol	62	61	84	81
LIFETIME				
Illicit Drug ²	58	58	82	75
Marijuana	48	48	73	71
Illicit Drug Other Than Marijuana ²	49	48	74	69
Cigarettes	62	62	84	83
Alcohol	66	64	84	82

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.12D Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2003 and 2004

(I) Demographic Characteristic and (II) Substance Use	Have Five or More Drinks of Alcoholic Beverage Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.42	0.40	0.42	0.40
AGE				
12-13	0.73	0.79	0.73	0.79
14-15	0.70	0.72	0.70	0.72
16-17	0.71	0.68	0.71	0.68
GENDER				
Male	0.57	0.61	0.57	0.61
Female	0.63	0.62	0.63	0.62
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.44	0.45	0.44	0.45
White	0.48	0.52	0.48	0.52
Black or African American	1.19	1.09	1.19	1.09
American Indian or Alaska Native	4.02	3.82	4.02	3.82
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.80	3.04	2.80	3.04
Two or More Races	3.34	2.60	3.34	2.60
Hispanic or Latino	1.22	1.25	1.22	1.25
COUNTY TYPE				
Large Metro	0.60	0.58	0.60	0.58
Small Metro	0.77	0.71	0.77	0.71
Nonmetro	0.97	0.99	0.97	0.99
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.34	0.38	0.37	0.37
Marijuana	0.29	0.28	0.33	0.34
Illicit Drug Other Than Marijuana ²	0.26	0.28	0.26	0.27
Cigarettes	0.39	0.41	0.39	0.39
Alcohol	0.46	0.48	0.45	0.44
Binge Alcohol ³	0.34	0.40	0.38	0.39
PAST YEAR				
Illicit Drug ²	0.49	0.54	0.50	0.47
Marijuana	0.41	0.42	0.43	0.43
Illicit Drug Other Than Marijuana ²	0.41	0.44	0.42	0.38
Cigarettes	0.52	0.51	0.47	0.47
Alcohol	0.65	0.64	0.55	0.52
LIFETIME				
Illicit Drug ²	0.61	0.61	0.54	0.49
Marijuana	0.50	0.51	0.48	0.46
Illicit Drug Other Than Marijuana ²	0.51	0.51	0.48	0.44
Cigarettes	0.65	0.65	0.55	0.54
Alcohol	0.69	0.67	0.55	0.53

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 3.12P Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week (II), among Persons Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

(I) Demographic Characteristic and (II) Substance Use	Have Five or More Drinks of Alcoholic Beverage Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2003	2004	2003	2004
(I) Demographic Characteristic				
TOTAL	0.4992		0.4992	
AGE				
12-13	0.7863		0.7863	
14-15	0.8668		0.8668	
16-17	0.2023		0.2023	
GENDER				
Male	0.0834		0.0834	
Female	0.4039		0.4039	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.3630		0.3630	
White	0.2106		0.2106	
Black or African American	0.7021		0.7021	
American Indian or Alaska Native	0.0310		0.0310	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.5576		0.5576	
Two or More Races	0.3710		0.3710	
Hispanic or Latino	0.7774		0.7774	
COUNTY TYPE				
Large Metro	0.7804		0.7804	
Small Metro	0.1114		0.1114	
Nonmetro	0.9316		0.9316	
(II) Substance Use				
PAST MONTH				
Illicit Drug ²	0.5959		0.1018	
Marijuana	0.3667		0.5529	
Illicit Drug Other Than Marijuana ²	0.5929		0.0540	
Cigarettes	0.8445		0.3195	
Alcohol	0.2834		0.2530	
Binge Alcohol ³	0.0395		0.9313	
PAST YEAR				
Illicit Drug ²	0.4551		0.1472	
Marijuana	0.8303		0.2500	
Illicit Drug Other Than Marijuana ²	0.4769		0.4072	
Cigarettes	0.9278		0.1344	
Alcohol	0.4459		0.7283	
LIFETIME				
Illicit Drug ²	0.6063		0.4417	
Marijuana	0.8039		0.2348	
Illicit Drug Other Than Marijuana ²	0.6509		0.8209	
Cigarettes	0.4848		0.0012	
Alcohol	0.5127		0.1500	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.