

Table 2.72C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	783	812	828	818	803	794
GEOGRAPHIC DIVISION						
Northeast	328	359	301	348	304	334
New England	138	169	169	169	165	156
Middle Atlantic	298	317	249	302	254	294
Midwest	298	274	350	325	322	310
East North Central	234	225	295	258	266	253
West North Central	186	156	188	197	180	178
South	485	504	545	539	558	510
South Atlantic	365	399	382	450	397	428
East South Central	190	182	253	189	256	170
West South Central	256	244	303	227	302	219
West	428	451	410	382	369	393
Mountain	175	167	171	149	149	152
Pacific	391	423	362	349	328	360
COUNTY TYPE						
Large Metro	621	649	600	606	571	576
Small Metro	399	385	483	435	479	435
250K – 1 Mil. Pop.	322	297	402	369	402	359
< 250K Pop.	239	263	269	241	247	246
Nonmetro	264	276	322	353	304	336
Urbanized	181	185	227	228	222	210
Less Urbanized	183	206	210	237	197	236
Completely Rural	86	78	121	79	105	73

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.72D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.33	0.34	0.35	0.34	0.34	0.33
GEOGRAPHIC DIVISION						
Northeast	0.72	0.79	0.66	0.76	0.67	0.73
New England	1.16	1.42	1.42	1.41	1.39	1.31
Middle Atlantic	0.89	0.94	0.75	0.90	0.76	0.88
Midwest	0.55	0.51	0.65	0.60	0.60	0.57
East North Central	0.62	0.59	0.78	0.68	0.71	0.67
West North Central	1.15	0.96	1.17	1.21	1.12	1.10
South	0.57	0.59	0.64	0.63	0.66	0.59
South Atlantic	0.82	0.88	0.86	1.00	0.89	0.95
East South Central	1.33	1.26	1.78	1.31	1.80	1.18
West South Central	0.98	0.92	1.16	0.86	1.15	0.83
West	0.80	0.83	0.76	0.70	0.69	0.72
Mountain	1.12	1.04	1.09	0.93	0.96	0.95
Pacific	1.02	1.09	0.95	0.90	0.86	0.93
COUNTY TYPE						
Large Metro	0.49	0.51	0.48	0.47	0.45	0.45
Small Metro	0.56	0.54	0.68	0.61	0.67	0.61
250K – 1 Mil. Pop.	0.69	0.63	0.86	0.78	0.86	0.76
< 250K Pop.	0.98	1.08	1.10	0.98	1.01	1.01
Nonmetro	0.66	0.68	0.80	0.87	0.76	0.83
Urbanized	1.01	1.11	1.27	1.37	1.24	1.26
Less Urbanized	1.01	1.01	1.15	1.17	1.08	1.16
Completely Rural	2.10	2.18	2.96	2.23	2.57	2.05

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.72P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.0065		0.2289		0.2089	
GEOGRAPHIC DIVISION						
Northeast	0.3354		0.5015		0.3165	
New England	0.9061		0.2322		0.8664	
Middle Atlantic	0.3262		0.8232		0.2547	
Midwest	0.3145		0.2275		0.7955	
East North Central	0.2841		0.5131		0.8504	
West North Central	0.8008		0.2522		0.4744	
South	0.1821		0.9868		0.9748	
South Atlantic	0.3253		0.1681		0.1854	
East South Central	0.2183		0.2525		0.4592	
West South Central	0.8320		0.3451		0.1613	
West	0.0435		0.3715		0.1437	
Mountain	0.9727		0.0421		0.0075	
Pacific	0.0269		0.8696		0.5635	
COUNTY TYPE						
Large Metro	0.0319		0.5773		0.2334	
Small Metro	0.1979		0.3146		0.6247	
250K – 1 Mil. Pop.	0.0022		0.1094		0.2069	
< 250K Pop.	0.0769		0.5363		0.2704	
Nonmetro	0.3058		0.6518		0.8302	
Urbanized	0.2315		0.9703		0.6841	
Less Urbanized	0.8054		0.9494		0.4416	
Completely Rural	0.7923		0.0656		0.0493	

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.73C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	113	106	93	96	80	80
GEOGRAPHIC DIVISION						
Northeast	44	41	39	41	32	36
New England	23	20	18	19	16	18
Middle Atlantic	37	36	35	36	28	32
Midwest	43	41	40	37	37	31
East North Central	34	33	30	29	30	22
West North Central	27	24	26	23	22	21
South	70	66	58	59	51	49
South Atlantic	46	49	39	44	36	38
East South Central	30	24	27	20	23	18
West South Central	44	36	35	33	28	26
West	64	60	47	51	38	41
Mountain	28	25	24	26	19	21
Pacific	57	54	40	45	32	35
COUNTY TYPE						
Large Metro	84	80	69	70	60	56
Small Metro	58	57	46	50	40	43
250K – 1 Mil. Pop.	49	44	40	40	35	34
< 250K Pop.	33	33	28	29	24	24
Nonmetro	44	48	39	44	31	37
Urbanized	28	24	25	23	20	19
Less Urbanized	29	36	27	32	22	28
Completely Rural	14	15	12	11	10	9

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.73D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.45	0.42	0.37	0.38	0.32	0.32
GEOGRAPHIC DIVISION						
Northeast	0.97	0.91	0.87	0.90	0.71	0.80
New England	1.96	1.69	1.50	1.59	1.38	1.50
Middle Atlantic	1.11	1.08	1.06	1.08	0.83	0.94
Midwest	0.76	0.73	0.71	0.66	0.66	0.54
East North Central	0.86	0.83	0.76	0.73	0.75	0.56
West North Central	1.57	1.46	1.55	1.38	1.30	1.27
South	0.78	0.73	0.65	0.65	0.57	0.55
South Atlantic	1.03	1.06	0.87	0.96	0.81	0.82
East South Central	2.07	1.67	1.85	1.36	1.58	1.25
West South Central	1.48	1.22	1.17	1.12	0.93	0.86
West	1.09	0.99	0.80	0.86	0.64	0.68
Mountain	1.63	1.46	1.37	1.46	1.08	1.19
Pacific	1.38	1.27	0.97	1.05	0.78	0.83
COUNTY TYPE						
Large Metro	0.63	0.60	0.52	0.52	0.45	0.42
Small Metro	0.78	0.74	0.62	0.66	0.53	0.57
250K – 1 Mil. Pop.	0.99	0.87	0.80	0.79	0.71	0.67
< 250K Pop.	1.36	1.31	1.15	1.12	0.99	0.95
Nonmetro	1.03	1.14	0.90	1.03	0.73	0.88
Urbanized	1.53	1.43	1.35	1.35	1.09	1.13
Less Urbanized	1.46	1.68	1.35	1.50	1.13	1.30
Completely Rural	3.23	4.04	2.79	3.03	2.29	2.49

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.73P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.0037		0.3724		0.9564	
GEOGRAPHIC DIVISION						
Northeast	0.1134		0.4359		0.7051	
New England	0.8904		0.2752		0.6617	
Middle Atlantic	0.0811		0.7582		0.8552	
Midwest	0.3391		0.2796		0.4166	
East North Central	0.4288		0.3447		0.5511	
West North Central	0.6024		0.5929		0.5887	
South	0.0098		0.1504		0.3731	
South Atlantic	0.1448		0.4173		0.8383	
East South Central	0.3429		0.6838		0.5240	
West South Central	0.0435		0.0660		0.0616	
West	0.5756		0.8375		0.1546	
Mountain	0.0938		0.3384		0.7902	
Pacific	0.9088		0.4640		0.0694	
COUNTY TYPE						
Large Metro	0.1077		0.7179		0.9277	
Small Metro	0.0303		0.7056		0.9176	
250K – 1 Mil. Pop.	0.0938		0.7650		0.6985	
< 250K Pop.	0.1581		0.2690		0.4308	
Nonmetro	0.1658		0.2910		0.8678	
Urbanized	0.0986		0.3215		0.6540	
Less Urbanized	0.3863		0.6313		0.6736	
Completely Rural	0.9873		0.3555		0.2320	

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.74C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	142	143	152	162	151	161
GEOGRAPHIC DIVISION						
Northeast	52	54	58	69	60	67
New England	23	28	26	31	20	26
Middle Atlantic	46	46	52	61	56	62
Midwest	56	52	69	67	68	70
East North Central	45	43	59	56	59	61
West North Central	33	30	35	37	33	36
South	93	92	92	100	92	98
South Atlantic	74	57	66	69	68	71
East South Central	33	33	39	41	40	37
West South Central	46	63	51	59	48	58
West	77	78	79	81	76	77
Mountain	29	30	41	35	35	30
Pacific	71	71	68	72	67	70
COUNTY TYPE						
Large Metro	108	109	120	118	117	114
Small Metro	79	76	78	90	77	94
250K – 1 Mil. Pop.	67	61	66	76	63	78
< 250K Pop.	43	52	44	55	44	53
Nonmetro	45	41	50	54	53	57
Urbanized	28	31	33	37	36	39
Less Urbanized	32	25	36	34	38	35
Completely Rural	10	8	14	14	14	14

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.74D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.45	0.44	0.48	0.50	0.48	0.50
GEOGRAPHIC DIVISION						
Northeast	0.92	0.95	1.03	1.21	1.06	1.19
New England	1.59	1.86	1.78	2.05	1.39	1.74
Middle Atlantic	1.12	1.09	1.24	1.46	1.35	1.48
Midwest	0.76	0.70	0.94	0.90	0.93	0.95
East North Central	0.89	0.83	1.17	1.09	1.17	1.18
West North Central	1.45	1.28	1.55	1.59	1.47	1.54
South	0.82	0.79	0.81	0.86	0.81	0.85
South Atlantic	1.32	1.00	1.18	1.21	1.21	1.24
East South Central	1.70	1.70	2.00	2.08	2.05	1.90
West South Central	1.20	1.62	1.35	1.53	1.25	1.49
West	1.03	1.03	1.07	1.07	1.02	1.03
Mountain	1.29	1.31	1.82	1.53	1.54	1.31
Pacific	1.37	1.36	1.32	1.37	1.30	1.34
COUNTY TYPE						
Large Metro	0.65	0.65	0.72	0.71	0.70	0.68
Small Metro	0.77	0.71	0.76	0.85	0.76	0.89
250K – 1 Mil. Pop.	1.01	0.88	0.99	1.10	0.94	1.13
< 250K Pop.	1.20	1.39	1.25	1.47	1.25	1.42
Nonmetro	0.92	0.83	1.01	1.09	1.08	1.15
Urbanized	1.18	1.36	1.40	1.59	1.52	1.70
Less Urbanized	1.46	1.09	1.65	1.47	1.75	1.52
Completely Rural	2.82	2.44	3.78	4.28	3.91	4.43

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.74P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.0447		0.4875		0.8192	
GEOGRAPHIC DIVISION						
Northeast	0.5644		0.4837		0.2531	
New England	0.4559		0.2448		0.0349	
Middle Atlantic	0.7731		0.8243		0.7374	
Midwest	0.7455		0.1758		0.1165	
East North Central	0.9554		0.2405		0.1593	
West North Central	0.5447		0.5046		0.4876	
South	0.1394		0.6522		0.1740	
South Atlantic	0.8115		0.5500		0.8584	
East South Central	0.6335		0.6027		0.8458	
West South Central	0.0147		0.0454		0.0080	
West	0.0825		0.8743		0.2234	
Mountain	0.7746		0.2807		0.5771	
Pacific	0.0767		0.6658		0.1033	
COUNTY TYPE						
Large Metro	0.2238		0.5095		0.8318	
Small Metro	0.1348		0.9885		0.5253	
250K – 1 Mil. Pop.	0.0238		0.4753		0.1697	
< 250K Pop.	0.5762		0.2554		0.3407	
Nonmetro	0.2772		0.6014		0.4447	
Urbanized	0.3053		0.4075		0.3167	
Less Urbanized	0.5396		0.7791		0.9034	
Completely Rural	0.7847		0.3974		0.8096	

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.75C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	729	768	774	768	748	752
GEOGRAPHIC DIVISION						
Northeast	325	338	278	322	284	315
New England	136	163	156	145	158	142
Middle Atlantic	296	296	229	286	234	280
Midwest	267	265	327	302	297	292
East North Central	212	215	270	242	241	241
West North Central	163	155	183	181	174	164
South	449	467	501	518	512	493
South Atlantic	330	368	362	439	369	421
East South Central	181	166	216	173	219	159
West South Central	241	232	275	214	281	199
West	392	434	396	350	354	366
Mountain	155	161	168	131	148	144
Pacific	359	405	348	323	312	335
COUNTY TYPE						
Large Metro	581	617	576	578	540	557
Small Metro	363	355	447	403	449	413
250K – 1 Mil. Pop.	297	282	375	335	381	337
< 250K Pop.	215	219	251	228	232	233
Nonmetro	253	269	305	331	288	318
Urbanized	169	173	214	211	212	196
Less Urbanized	179	203	194	225	182	226
Completely Rural	88	74	119	73	103	69

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.75D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.40	0.42	0.43	0.42	0.41	0.41
GEOGRAPHIC DIVISION						
Northeast	0.92	0.96	0.79	0.91	0.81	0.89
New England	1.47	1.75	1.69	1.56	1.70	1.53
Middle Atlantic	1.14	1.14	0.88	1.10	0.90	1.08
Midwest	0.66	0.65	0.80	0.74	0.73	0.71
East North Central	0.74	0.75	0.95	0.84	0.84	0.84
West North Central	1.34	1.27	1.51	1.47	1.43	1.33
South	0.70	0.71	0.78	0.79	0.79	0.75
South Atlantic	0.96	1.05	1.05	1.26	1.07	1.21
East South Central	1.67	1.51	1.99	1.57	2.02	1.45
West South Central	1.25	1.18	1.42	1.09	1.45	1.01
West	0.97	1.05	0.98	0.85	0.87	0.89
Mountain	1.33	1.35	1.44	1.10	1.27	1.20
Pacific	1.24	1.39	1.21	1.11	1.08	1.15
COUNTY TYPE						
Large Metro	0.60	0.63	0.60	0.59	0.56	0.57
Small Metro	0.68	0.66	0.83	0.76	0.83	0.77
250K – 1 Mil. Pop.	0.84	0.80	1.06	0.95	1.08	0.96
< 250K Pop.	1.16	1.20	1.36	1.25	1.26	1.28
Nonmetro	0.82	0.86	0.98	1.06	0.93	1.01
Urbanized	1.24	1.36	1.57	1.66	1.55	1.55
Less Urbanized	1.27	1.29	1.38	1.42	1.29	1.43
Completely Rural	2.66	2.57	3.62	2.56	3.13	2.41

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.75P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.0369		0.1774		0.1929	
GEOGRAPHIC DIVISION						
Northeast	0.4859		0.3089		0.1647	
New England	0.9597		0.0438		0.6513	
Middle Atlantic	0.4482		0.7396		0.1882	
Midwest	0.2813		0.1266		0.5359	
East North Central	0.2922		0.3571		0.8892	
West North Central	0.6859		0.1907		0.3948	
South	0.4463		0.7940		0.7564	
South Atlantic	0.3966		0.1700		0.1762	
East South Central	0.2893		0.1496		0.3876	
West South Central	0.5516		0.7334		0.4772	
West	0.0875		0.3007		0.1521	
Mountain	0.8248		0.0137		0.0045	
Pacific	0.0482		0.8555		0.6598	
COUNTY TYPE						
Large Metro	0.0545		0.5153		0.2263	
Small Metro	0.5921		0.2599		0.6841	
250K – 1 Mil. Pop.	0.0179		0.1011		0.2748	
< 250K Pop.	0.0252		0.6303		0.3157	
Nonmetro	0.4568		0.6639		0.7148	
Urbanized	0.3608		0.9305		0.5017	
Less Urbanized	0.9272		0.7923		0.4351	
Completely Rural	0.7487		0.0572		0.0557	

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.76C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	755	795	810	816	784	793
GEOGRAPHIC DIVISION						
Northeast	334	347	296	345	296	332
New England	142	161	165	164	162	154
Middle Atlantic	303	307	245	302	247	294
Midwest	281	268	348	326	316	311
East North Central	222	219	292	255	261	252
West North Central	173	154	189	203	179	181
South	455	488	524	541	537	510
South Atlantic	337	385	372	450	384	428
East South Central	180	175	231	189	237	172
West South Central	244	240	291	233	293	218
West	415	450	403	378	365	392
Mountain	163	172	168	147	146	154
Pacific	381	419	355	345	324	358
COUNTY TYPE						
Large Metro	601	638	586	601	554	578
Small Metro	381	366	472	434	471	435
250K – 1 Mil. Pop.	308	290	395	367	397	361
< 250K Pop.	227	239	263	242	243	245
Nonmetro	258	273	319	357	302	337
Urbanized	174	180	224	230	221	210
Less Urbanized	180	205	207	238	195	235
Completely Rural	89	74	120	80	104	74

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.76D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.35	0.37	0.38	0.38	0.37	0.37
GEOGRAPHIC DIVISION						
Northeast	0.82	0.85	0.72	0.84	0.73	0.81
New England	1.33	1.49	1.54	1.52	1.51	1.43
Middle Atlantic	1.01	1.02	0.81	1.00	0.82	0.97
Midwest	0.59	0.55	0.72	0.67	0.66	0.64
East North Central	0.66	0.65	0.87	0.75	0.77	0.74
West North Central	1.20	1.05	1.31	1.39	1.24	1.24
South	0.60	0.63	0.69	0.70	0.71	0.66
South Atlantic	0.84	0.95	0.93	1.11	0.96	1.05
East South Central	1.41	1.35	1.81	1.46	1.86	1.33
West South Central	1.06	1.02	1.26	0.99	1.27	0.92
West	0.87	0.93	0.84	0.78	0.76	0.81
Mountain	1.17	1.21	1.21	1.03	1.05	1.08
Pacific	1.12	1.22	1.04	1.00	0.95	1.04
COUNTY TYPE						
Large Metro	0.53	0.55	0.52	0.52	0.49	0.50
Small Metro	0.60	0.57	0.74	0.68	0.74	0.68
250K – 1 Mil. Pop.	0.73	0.69	0.94	0.87	0.95	0.86
< 250K Pop.	1.03	1.09	1.19	1.11	1.10	1.12
Nonmetro	0.72	0.75	0.89	0.98	0.84	0.93
Urbanized	1.09	1.20	1.40	1.53	1.38	1.40
Less Urbanized	1.11	1.13	1.27	1.31	1.20	1.30
Completely Rural	2.43	2.31	3.27	2.51	2.84	2.33

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.76P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.0172		0.2632		0.2030	
GEOGRAPHIC DIVISION						
Northeast	0.4498		0.4367		0.2938	
New England	0.9132		0.1495		0.9258	
Middle Atlantic	0.4438		0.7997		0.2462	
Midwest	0.3307		0.2680		0.8466	
East North Central	0.3116		0.5764		0.8023	
West North Central	0.7813		0.2732		0.5002	
South	0.3090		0.8631		0.9664	
South Atlantic	0.4269		0.1395		0.1726	
East South Central	0.2624		0.2121		0.4018	
West South Central	0.9595		0.4557		0.2145	
West	0.0499		0.3499		0.1096	
Mountain	0.8748		0.0503		0.0092	
Pacific	0.0253		0.8004		0.4552	
COUNTY TYPE						
Large Metro	0.0403		0.5774		0.2094	
Small Metro	0.4003		0.3537		0.6745	
250K – 1 Mil. Pop.	0.0055		0.1067		0.2090	
< 250K Pop.	0.0273		0.4208		0.2083	
Nonmetro	0.3686		0.7284		0.8166	
Urbanized	0.2910		0.9005		0.6977	
Less Urbanized	0.8615		0.9131		0.4892	
Completely Rural	0.7769		0.0850		0.0722	

*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.