

**Table 6.4C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Use of Illicit Drugs: Standard Errors of Numbers in Thousands, 2003 and 2004**

Drug	ILLCIT DRUG USE			
	Yes		No	
	2003	2004	2003	2004
<b>ILLCIT DRUG<sup>1</sup></b>	169	332	402	799
Marijuana and Hashish	133	262	413	823
Cocaine	78	132	423	853
Crack	45	*	428	862
Heroin	*	*	432	864
Hallucinogens	49	86	425	866
LSD	14	*	431	871
PCP	*	*	432	872
Ecstasy	33	60	430	872
Inhalants	26	*	430	867
Nonmedical Use of Psychotherapeutics <sup>2</sup>	129	237	420	831
Pain Relievers	104	187	423	837
OxyContin <sup>®</sup>	--	*	--	865
Tranquilizers	83	170	423	839
Stimulants	51	95	430	865
Methamphetamine	39	*	429	874
Sedatives	*	*	429	868
<b>ILLCIT DRUG OTHER THAN MARIJUANA<sup>1</sup></b>	138	276	409	821

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.4D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Use of Illicit Drugs: Standard Errors of Percentages, 2003 and 2004**

Drug	ILLCIT DRUG USE			
	Yes		No	
	2003	2004	2003	2004
<b>ILLCIT DRUG<sup>1</sup></b>	0.57	1.15	0.22	0.43
Marijuana and Hashish	0.62	1.26	0.22	0.42
Cocaine	1.42	2.41	0.20	0.41
Crack	3.42	*	0.20	0.40
Heroin	*	*	0.20	0.40
Hallucinogens	1.55	2.76	0.20	0.41
LSD	3.36	*	0.20	0.41
PCP	*	*	0.20	0.41
Ecstasy	1.83	3.77	0.20	0.41
Inhalants	2.67	*	0.20	0.40
Nonmedical Use of Psychotherapeutics <sup>2</sup>	1.01	1.85	0.21	0.41
Pain Relievers	1.07	1.97	0.21	0.41
OxyContin <sup>®</sup>	--	*	--	0.40
Tranquilizers	1.87	3.32	0.20	0.40
Stimulants	2.32	4.28	0.20	0.41
Methamphetamine	3.38	*	0.20	0.41
Sedatives	*	*	0.20	0.40
<b>ILLCIT DRUG OTHER THAN MARIJUANA<sup>1</sup></b>	0.82	1.62	0.21	0.41

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.4P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Use of Illicit Drugs: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Drug	ILLCIT DRUG USE			
	Yes		No	
	2003	2004	2003	2004
<b>ILLCIT DRUG<sup>1</sup></b>	0.0518		0.2757	
Marijuana and Hashish	0.0297		0.2617	
Cocaine	0.1897		0.1473	
Crack	*		0.1192	
Heroin	*		0.1328	
Hallucinogens	0.2429		0.1275	
LSD	*		0.1192	
PCP	*		0.1119	
Ecstasy	0.2407		0.1178	
Inhalants	*		0.1197	
Nonmedical Use of Psychotherapeutics <sup>2</sup>	0.0850		0.2345	
Pain Relievers	0.2643		0.1440	
OxyContin <sup>®</sup>	--		--	
Tranquilizers	0.0886		0.2263	
Stimulants	0.1720		0.1392	
Methamphetamine	*		0.1223	
Sedatives	*		0.0830	
<b>ILLCIT DRUG OTHER THAN MARIJUANA<sup>1</sup></b>	0.1330		0.2173	

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.5C Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Numbers in Thousands, 2003 and 2004**

Drug	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>ILLCIT DRUG<sup>2</sup></b>	574	819	195	371	506	732
Marijuana and Hashish	457	683	150	289	410	616
Cocaine	232	323	84	147	213	289
Crack	123	215	51	105	107	187
Heroin	47	101	16	88	45	50
Hallucinogens	141	270	54	84	128	256
LSD	41	55	15	39	38	40
PCP	20	41	7	30	19	25
Ecstasy	102	212	34	55	96	202
Inhalants	67	125	26	95	63	94
Nonmedical Use of Psychotherapeutics <sup>3</sup>	361	530	143	275	323	456
Pain Relievers	304	441	119	215	267	385
OxyContin <sup>®</sup>	--	132	--	102	--	85
Tranquilizers	195	326	93	216	173	252
Stimulants	124	196	60	123	105	151
Methamphetamine	93	153	48	87	74	125
Sedatives	83	149	53	51	67	140
<b>ILLCIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	425	627	153	313	382	552

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.5D Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Percentages, 2003 and 2004**

Drug	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>ILLCIT DRUG<sup>2</sup></b>	0.25	0.36	0.95	1.61	0.25	0.36
Marijuana and Hashish	0.21	0.31	0.75	1.31	0.20	0.31
Cocaine	0.11	0.14	0.42	0.68	0.11	0.14
Crack	0.06	0.10	0.26	0.48	0.06	0.10
Heroin	0.02	0.05	0.08	0.40	0.02	0.03
Hallucinogens	0.07	0.12	0.27	0.39	0.07	0.13
LSD	0.02	0.03	0.07	0.18	0.02	0.02
PCP	0.01	0.02	0.04	0.14	0.01	0.01
Ecstasy	0.05	0.10	0.17	0.26	0.05	0.10
Inhalants	0.03	0.06	0.13	0.44	0.03	0.05
Nonmedical Use of Psychotherapeutics <sup>3</sup>	0.17	0.24	0.72	1.24	0.16	0.23
Pain Relievers	0.14	0.20	0.60	0.97	0.14	0.20
OxyContin <sup>®</sup>	--	0.06	--	0.47	--	0.04
Tranquilizers	0.09	0.15	0.47	0.96	0.09	0.13
Stimulants	0.06	0.09	0.31	0.57	0.05	0.08
Methamphetamine	0.04	0.07	0.25	0.41	0.04	0.06
Sedatives	0.04	0.07	0.27	0.24	0.03	0.07
<b>ILLCIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	0.19	0.28	0.74	1.38	0.19	0.28

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.5P Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Drug	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>ILLCIT DRUG<sup>2</sup></b>	0.2095		0.8702		0.0844	
Marijuana and Hashish	0.2489		0.6255		0.0866	
Cocaine	0.8936		0.5588		0.5539	
Crack	0.6208		0.6311		0.8285	
Heroin	0.3763		0.1488		0.6350	
Hallucinogens	0.8316		0.5852		0.6050	
LSD	0.7189		0.2136		0.1463	
PCP	0.7253		0.3859		0.6889	
Ecstasy	0.3334		0.8397		0.2516	
Inhalants	0.3913		0.5214		0.6806	
Nonmedical Use of Psychotherapeutics <sup>3</sup>	0.9696		0.4286		0.4723	
Pain Relievers	0.5090		0.9837		0.3341	
OxyContin <sup>®</sup>	--		--		--	
Tranquilizers	0.1216		0.0864		0.6886	
Stimulants	0.9291		0.4385		0.5090	
Methamphetamine	0.8392		0.6740		0.5071	
Sedatives	0.7989		0.0846		0.6614	
<b>ILLCIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	0.8371		0.6617		0.4767	

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.6C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year and Past Month Tobacco Product and Alcohol Use: Standard Errors of Numbers in Thousands, 2003 and 2004**

Substance	TOBACCO PRODUCT OR ALCOHOL USE			
	Yes		No	
	2003	2004	2003	2004
<b>PAST YEAR</b>				
Tobacco Product <sup>1</sup>	287	481	318	720
Cigarettes	274	465	326	731
Smokeless Tobacco	77	137	426	862
Cigars	146	192	420	853
Pipe Tobacco <sup>2</sup>	--	--	--	--
Alcohol	339	582	259	640
Binge Alcohol Use <sup>3</sup>	--	--	--	--
Heavy Alcohol Use <sup>3</sup>	--	--	--	--
<b>PAST MONTH</b>				
Tobacco Product <sup>1</sup>	273	462	330	738
Cigarettes	258	448	341	748
Smokeless Tobacco	62	106	425	869
Cigars	107	147	422	859
Pipe Tobacco	59	65	431	869
Alcohol	291	524	320	690
Binge Alcohol Use <sup>3</sup>	195	386	383	782
Heavy Alcohol Use <sup>3</sup>	116	175	414	846

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.6D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year and Past Month Tobacco Product and Alcohol Use: Standard Errors of Percentages, 2003 and 2004**

Substance	TOBACCO PRODUCT OR ALCOHOL USE			
	Yes		No	
	2003	2004	2003	2004
<b>PAST YEAR</b>				
Tobacco Product <sup>1</sup>	0.37	0.62	0.24	0.52
Cigarettes	0.42	0.72	0.22	0.49
Smokeless Tobacco	0.82	1.59	0.21	0.42
Cigars	0.64	0.84	0.22	0.44
Pipe Tobacco <sup>2</sup>	--	--	--	--
Alcohol	0.23	0.40	0.39	0.93
Binge Alcohol Use <sup>3</sup>	--	--	--	--
Heavy Alcohol Use <sup>3</sup>	--	--	--	--
<b>PAST MONTH</b>				
Tobacco Product <sup>1</sup>	0.41	0.70	0.23	0.49
Cigarettes	0.45	0.80	0.22	0.47
Smokeless Tobacco	0.87	1.69	0.21	0.42
Cigars	0.91	1.14	0.21	0.42
Pipe Tobacco	4.06	3.81	0.20	0.41
Alcohol	0.25	0.46	0.33	0.69
Binge Alcohol Use <sup>3</sup>	0.38	0.75	0.24	0.48
Heavy Alcohol Use <sup>3</sup>	0.75	1.12	0.21	0.42

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).



**Table 6.6P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year and Past Month Tobacco Product and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Substance	TOBACCO PRODUCT OR ALCOHOL USE			
	Yes		No	
	2003	2004	2003	2004
<b>PAST YEAR</b>				
Tobacco Product <sup>1</sup>	0.8965		0.0252	
Cigarettes	0.9735		0.0405	
Smokeless Tobacco	0.2078		0.1528	
Cigars	0.4074		0.0637	
Pipe Tobacco <sup>2</sup>	--		--	
Alcohol	0.7068		0.0495	
Binge Alcohol Use <sup>3</sup>	--		--	
Heavy Alcohol Use <sup>3</sup>	--		--	
<b>PAST MONTH</b>				
Tobacco Product <sup>1</sup>	0.7093		0.0620	
Cigarettes	0.6090		0.0725	
Smokeless Tobacco	0.2054		0.1461	
Cigars	0.5227		0.0836	
Pipe Tobacco	0.5700		0.0968	
Alcohol	0.2192		0.2720	
Binge Alcohol Use <sup>3</sup>	0.0730		0.3521	
Heavy Alcohol Use <sup>3</sup>	0.3711		0.1362	

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.7C Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Numbers in Thousands, 2003 and 2004**

Substance	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>PAST YEAR</b>						
Tobacco Product <sup>2</sup>	1,042	1,536	325	516	935	1,430
Cigarettes	956	1,403	313	507	849	1,289
Smokeless Tobacco	314	443	80	142	302	429
Cigars	498	778	158	188	460	764
Pipe Tobacco <sup>3</sup>	--	--	--	--	--	--
Alcohol	1,725	2,223	371	613	1,610	2,110
Binge Alcohol Use <sup>4</sup>	--	--	--	--	--	--
Heavy Alcohol Use <sup>4</sup>	--	--	--	--	--	--
<b>PAST MONTH</b>						
Tobacco Product <sup>2</sup>	972	1,451	307	496	876	1,350
Cigarettes	901	1,312	294	487	801	1,205
Smokeless Tobacco	288	356	64	108	280	346
Cigars	353	599	115	145	327	589
Pipe Tobacco	130	228	66	67	115	217
Alcohol	1,507	2,030	316	558	1,413	1,913
Binge Alcohol Use <sup>4</sup>	787	1,147	214	411	727	1,078
Heavy Alcohol Use <sup>4</sup>	401	554	124	182	374	529

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>3</sup> Information about past year use of pipe tobacco was not collected.

<sup>4</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.7D Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Percentages, 2003 and 2004**

Substance	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>PAST YEAR</b>						
Tobacco Product <sup>2</sup>	0.38	0.61	1.18	2.10	0.39	0.63
Cigarettes	0.38	0.58	1.18	2.06	0.38	0.60
Smokeless Tobacco	0.15	0.20	0.40	0.66	0.15	0.22
Cigars	0.21	0.33	0.76	0.92	0.23	0.37
Pipe Tobacco <sup>3</sup>	--	--	--	--	--	--
Alcohol	0.42	0.63	1.07	2.34	0.45	0.65
Binge Alcohol Use <sup>4</sup>	--	--	--	--	--	--
Heavy Alcohol Use <sup>4</sup>	--	--	--	--	--	--
<b>PAST MONTH</b>						
Tobacco Product <sup>2</sup>	0.37	0.58	1.17	2.05	0.37	0.60
Cigarettes	0.36	0.54	1.16	2.01	0.36	0.56
Smokeless Tobacco	0.13	0.16	0.32	0.51	0.14	0.17
Cigars	0.16	0.26	0.56	0.69	0.16	0.29
Pipe Tobacco	0.06	0.11	0.34	0.31	0.06	0.11
Alcohol	0.43	0.67	1.16	2.19	0.45	0.69
Binge Alcohol Use <sup>4</sup>	0.32	0.49	0.96	1.76	0.33	0.52
Heavy Alcohol Use <sup>4</sup>	0.18	0.25	0.60	0.84	0.18	0.27

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>3</sup> Information about past year use of pipe tobacco was not collected.

<sup>4</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.7P Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: P Values from Tests of Differences of Percentages, 2004 Versus 2003**

Substance	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>	
			Yes	No
	2003	2004	2003	2004
<b>PAST YEAR</b>				
Tobacco Product <sup>2</sup>	0.2842		0.0281	0.5670
Cigarettes	0.4752		0.0737	0.7357
Smokeless Tobacco	0.1255		0.6770	0.0896
Cigars	0.7987		0.0882	0.8459
Pipe Tobacco <sup>3</sup>	--		--	--
Alcohol	0.4591		0.0691	0.8715
Binge Alcohol Use <sup>4</sup>	--		--	--
Heavy Alcohol Use <sup>4</sup>	--		--	--
<b>PAST MONTH</b>				
Tobacco Product <sup>2</sup>	0.1979		0.1083	0.3138
Cigarettes	0.1499		0.1373	0.2170
Smokeless Tobacco	0.0178		0.7562	0.0127
Cigars	0.1439		0.5651	0.0971
Pipe Tobacco	0.3948		0.7233	0.3135
Alcohol	0.3882		0.6810	0.4374
Binge Alcohol Use <sup>4</sup>	0.8441		0.4927	0.5998
Heavy Alcohol Use <sup>4</sup>	0.8626		0.9290	0.7757

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>3</sup> Information about past year use of pipe tobacco was not collected.

<sup>4</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.8C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug Use and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004**

Demographic Characteristic	Total		ILLCIT DRUG USE <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	432	872	169	332	402	799
<b>AGE</b>						
18-25	100	164	67	98	71	128
26-49	280	482	140	273	254	373
50 or Older	307	713	71	*	300	698
<b>GENDER</b>						
Male	264	618	109	234	245	577
Female	345	601	133	210	319	550
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	415	736	159	312	379	652
White	388	670	147	270	354	609
Black or African American	143	206	60	104	131	170
American Indian or Alaska Native	21	*	*	*	19	*
Native Hawaiian or Other Pacific Islander	15	*	*	*	10	*
Asian	83	137	*	*	76	97
Two or More Races	48	36	13	*	46	28
Hispanic or Latino	151	485	60	103	145	476
<b>EDUCATION</b>						
< High School	223	519	83	129	201	508
High School Graduate	244	465	98	202	221	412
Some College	212	369	92	149	198	338
College Graduate	186	324	73	156	182	286
<b>CURRENT EMPLOYMENT</b>						
Full-Time	291	438	131	197	262	379
Part-Time	151	261	60	158	143	181
Unemployed	85	183	46	123	73	126
Other <sup>2</sup>	271	645	86	139	251	633

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.8D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug Use and Demographic Characteristics: Standard Errors of Percentages, 2003 and 2004**

Demographic Characteristic	Total		ILLCIT DRUG USE <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.20	0.41	0.57	1.15	0.22	0.43
<b>AGE</b>						
18-25	0.32	0.51	0.61	0.92	0.34	0.59
26-49	0.28	0.48	0.89	1.80	0.30	0.44
50 or Older	0.38	0.86	2.59	*	0.38	0.87
<b>GENDER</b>						
Male	0.26	0.60	0.64	1.40	0.29	0.66
Female	0.31	0.54	1.07	1.75	0.33	0.55
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.22	0.39	0.61	1.22	0.24	0.40
White	0.26	0.44	0.69	1.31	0.27	0.47
Black or African American	0.60	0.85	1.71	2.92	0.65	0.82
American Indian or Alaska Native	1.92	*	*	*	2.11	*
Native Hawaiian or Other Pacific Islander	3.53	*	*	*	2.89	*
Asian	0.94	1.51	*	*	0.92	1.15
Two or More Races	2.22	1.90	2.98	*	2.64	1.88
Hispanic or Latino	0.58	1.81	1.69	3.27	0.65	2.01
<b>EDUCATION</b>						
< High School	0.59	1.43	1.50	2.60	0.62	1.62
High School Graduate	0.36	0.68	1.02	2.16	0.38	0.70
Some College	0.40	0.66	1.07	1.78	0.45	0.72
College Graduate	0.34	0.59	1.27	2.60	0.37	0.58
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.25	0.37	0.76	1.18	0.26	0.38
Part-Time	0.53	0.98	1.13	3.21	0.61	0.83
Unemployed	1.05	2.28	1.97	5.27	1.26	2.21
Other <sup>2</sup>	0.46	1.02	1.86	2.86	0.46	1.08

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.8P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug Use and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Demographic Characteristic	Total		ILLCIT DRUG USE <sup>1</sup>	
			Yes	No
	2003	2004	2003	2004
<b>TOTAL</b>	0.1052		0.0518	0.2757
<b>AGE</b>				
18-25	0.7137		0.9698	0.8867
26-49	0.9705		0.1012	0.3312
50 or Older	0.0279		*	0.0558
<b>GENDER</b>				
Male	0.1422		0.0356	0.4226
Female	0.4026		0.4286	0.4516
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic or Latino	0.1986		0.0582	0.4999
White	0.1493		0.0298	0.4420
Black or African American	0.7694		0.4899	0.9510
American Indian or Alaska Native	*		*	*
Native Hawaiian or Other Pacific Islander	*		*	*
Asian	0.7244		*	0.8376
Two or More Races	0.1845		*	0.0556
Hispanic or Latino	0.3406		0.6725	0.3421
<b>EDUCATION</b>				
< High School	0.3207		0.0914	0.5261
High School Graduate	0.3566		0.1612	0.6622
Some College	0.7171		0.8910	0.8951
College Graduate	0.0560		0.2104	0.1579
<b>CURRENT EMPLOYMENT</b>				
Full-Time	0.2496		0.4431	0.4388
Part-Time	0.2624		0.0122	0.7401
Unemployed	0.1365		0.0273	0.9736
Other <sup>2</sup>	0.0320		0.4116	0.0426

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.9C Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004**

Demographic Characteristic	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	574	819	195	371	506	732
<b>AGE</b>						
18-25	212	303	81	114	180	264
26-49	448	644	164	309	389	567
50 or Older	229	375	76	165	212	334
<b>GENDER</b>						
Male	404	614	122	256	367	560
Female	337	479	153	253	293	399
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	527	757	184	353	463	668
White	468	691	168	308	413	605
Black or African American	218	276	75	111	183	261
American Indian or Alaska Native	28	75	*	*	25	*
Native Hawaiian or Other Pacific Islander	17	*	*	*	12	*
Asian	77	141	*	*	72	100
Two or More Races	53	85	*	*	51	81
Hispanic or Latino	195	273	65	113	180	248
<b>EDUCATION</b>						
< High School	229	282	99	154	195	234
High School Graduate	287	451	110	235	256	384
Some College	277	429	105	156	244	403
College Graduate	259	428	77	171	244	389
<b>CURRENT EMPLOYMENT</b>						
Full-Time	424	627	151	208	372	589
Part-Time	189	298	66	197	173	222
Unemployed	114	237	47	*	110	170
Other <sup>2</sup>	214	311	96	164	189	263

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).



**Table 6.9D Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress and Demographic Characteristics: Standard Errors of Percentages, 2003 and 2004**

Demographic Characteristic	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.25	0.36	0.95	1.61	0.25	0.36
<b>AGE</b>						
18-25	0.48	0.73	1.24	2.05	0.49	0.72
26-49	0.42	0.58	1.40	2.27	0.41	0.59
50 or Older	0.28	0.44	1.57	2.48	0.27	0.43
<b>GENDER</b>						
Male	0.37	0.55	1.72	3.29	0.36	0.55
Female	0.29	0.43	1.14	1.72	0.29	0.41
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.27	0.38	1.01	1.67	0.26	0.38
White	0.30	0.44	1.12	1.82	0.29	0.44
Black or African American	0.78	1.11	3.40	4.67	0.72	1.14
American Indian or Alaska Native	2.06	5.32	*	*	2.06	*
Native Hawaiian or Other Pacific Islander	4.05	*	*	*	3.53	*
Asian	0.88	1.54	*	*	0.88	1.19
Two or More Races	2.38	4.09	*	*	2.60	4.26
Hispanic or Latino	0.67	0.95	2.64	4.50	0.70	0.98
<b>EDUCATION</b>						
< High School	0.59	0.80	2.12	4.11	0.57	0.76
High School Graduate	0.40	0.65	1.59	3.00	0.40	0.62
Some College	0.49	0.72	1.86	2.71	0.49	0.75
College Graduate	0.46	0.72	2.16	3.46	0.46	0.72
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.34	0.49	1.43	1.99	0.33	0.50
Part-Time	0.61	1.07	2.33	4.58	0.63	0.93
Unemployed	1.33	2.69	3.48	*	1.47	2.53
Other <sup>2</sup>	0.36	0.49	1.55	2.29	0.35	0.48

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.9P Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Demographic Characteristic	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.2095		0.8702		0.0844	
<b>AGE</b>						
18-25	0.0742		0.5023		0.0834	
26-49	0.3365		0.1735		0.0961	
50 or Older	0.7566		0.4474		0.8474	
<b>GENDER</b>						
Male	0.3862		0.6220		0.1542	
Female	0.3013		0.7022		0.2457	
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.4316		0.5418		0.1716	
White	0.3427		0.5516		0.1172	
Black or African American	0.9761		0.5807		0.8270	
American Indian or Alaska Native	0.0835		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.8129		*		0.5664	
Two or More Races	0.6232		*		0.9651	
Hispanic or Latino	0.1265		0.3652		0.1537	
<b>EDUCATION</b>						
< High School	0.2763		0.9447		0.1047	
High School Graduate	0.5016		0.6132		0.2359	
Some College	0.1052		0.4851		0.1613	
College Graduate	0.5914		0.6058		0.8570	
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.2950		0.5998		0.4330	
Part-Time	0.8838		0.0499		0.1369	
Unemployed	0.8779		*		0.3250	
Other <sup>2</sup>	0.8333		0.3930		0.7607	

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).