

**Table 2.19A Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	172,843	171,827	83,415	83,066	70,757	70,257
12	398	339	188	169	89	61
13	782	757	410	419	184	222
14	1,206	1,231	748	767	428	447
15	1,727	1,653	1,120	1,110	671	697
16	2,116	1,965	1,463	1,404	990	946
17	2,400	2,304	1,705	1,692	1,248	1,262
18	3,018	3,026	2,285	2,377	1,759	1,874
19	2,813	2,745	2,036	2,123	1,632	1,690
20	3,053	2,911	2,303	2,240	1,950	1,800
21	2,930	3,073	2,129	2,253	1,834	1,878
22	3,186	3,096	2,305	2,274	1,928	1,908
23	3,095	2,912	2,187	2,105	1,843	1,762
24	2,833 <sup>a</sup>	3,129	1,992	2,167	1,700	1,809
25	2,795	2,765	1,837	1,942	1,555	1,637
26-29	11,565	11,703	7,246	7,418	6,151	6,314
30-34	15,582	14,642	8,621	8,044	7,410 <sup>a</sup>	6,708
35-39	16,424	16,161	8,520	8,521	7,291	7,270
40-44	18,091	18,332	9,339	9,188	8,360	8,018
45-49	17,133	17,261	8,049	7,949	7,129	7,120
50-54	15,669	15,622	6,547	6,090	5,776	5,427
55-59	12,062	12,865	4,478	4,662	3,939	4,196
60-64	9,479	9,865	2,645	3,014	2,361	2,737
65 or Older	24,484	23,468	5,261	5,137	4,529	4,472

\*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 2.19B Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	72.7 <sup>b</sup>	71.4	35.1	34.5	29.8	29.2
12	9.7	8.5	4.6	4.2	2.2	1.5
13	18.1	17.2	9.5	9.5	4.2	5.1
14	29.2	28.1	18.1	17.5	10.4	10.2
15	41.8 <sup>a</sup>	38.5	27.1	25.9	16.2	16.2
16	50.5	47.7	34.9	34.1	23.6	23.0
17	58.6	56.7	41.7	41.6	30.5	31.0
18	65.5	65.8	49.6	51.7	38.2	40.8
19	72.4	69.7	52.4	53.9	42.0	42.9
20	75.8	73.0	57.2	56.1	48.4	45.1
21	74.7	75.3	54.3	55.2	46.7	46.0
22	79.3	77.0	57.4	56.6	48.0	47.5
23	77.3	74.2	54.6	53.6	46.0	44.9
24	77.9	77.3	54.8	53.5	46.8	44.7
25	77.0	76.9	50.6	54.0	42.8	45.5
26-29	78.3	76.3	49.1	48.3	41.7	41.1
30-34	77.2	74.6	42.7	41.0	36.7	34.2
35-39	78.5	77.2	40.7	40.7	34.8	34.7
40-44	79.7	80.8	41.2	40.5	36.8	35.4
45-49	80.0	80.3	37.6	37.0	33.3	33.1
50-54	78.4	80.6	32.7	31.4	28.9	28.0
55-59	80.9	78.4	30.0	28.4	26.4	25.6
60-64	80.6	78.7	22.5	24.0	20.1	21.8
65 or Older	71.4 <sup>a</sup>	67.7	15.3	14.8	13.2	12.9

\*Low precision; no estimate reported.

NOTE: Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 2.20A Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	163,240	161,842	69,853	69,909	60,434	59,896
12	322	277	146	132	70	45
13	662	653	325	362	143 <sup>a</sup>	190
14	1,056	1,097	616	656	347	368
15	1,539	1,467	928	913	561	578
16	1,957	1,790	1,272	1,193	842	787
17	2,217	2,079	1,449	1,390	1,080	1,031
18	2,763	2,722	1,972	1,983	1,551	1,597
19	2,636	2,554	1,766	1,855	1,445	1,464
20	2,866	2,707	2,063	1,955	1,775	1,608
21	2,774	2,912	1,891	2,021	1,647	1,680
22	3,034	2,922	2,039	2,020	1,749	1,725
23	2,927	2,731	1,960	1,875	1,666	1,595
24	2,667 <sup>a</sup>	2,942	1,780	1,903	1,532	1,605
25	2,601	2,628	1,642	1,679	1,400	1,456
26-29	10,788	11,046	6,304	6,578	5,434	5,643
30-34	14,478	13,664	7,155	6,724	6,233	5,702
35-39	15,519	15,262	6,875	7,148	6,078	6,202
40-44	17,159	17,457	7,724	7,748	7,063	6,891
45-49	16,446	16,422	6,779	6,624	6,181	6,048
50-54	15,182	14,946	5,473	5,216	4,993	4,777
55-59	11,518	12,251	3,502	3,752	3,258	3,416
60-64	9,094	9,376	2,148	2,486	1,944	2,285
65 or Older	23,038	21,936	4,043	3,698	3,444	3,203

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 2.20B Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	68.7 <sup>b</sup>	67.3	29.4	29.1	25.4	24.9
12	7.8	7.0	3.5	3.3	1.7	1.1
13	15.3	14.9	7.5	8.2	3.3 <sup>a</sup>	4.3
14	25.6	25.1	14.9	15.0	8.4	8.4
15	37.2 <sup>a</sup>	34.2	22.5	21.3	13.6	13.5
16	46.7 <sup>a</sup>	43.5	30.4	29.0	20.1	19.1
17	54.2	51.1	35.4	34.2	26.4	25.4
18	60.0	59.2	42.8	43.1	33.7	34.8
19	67.8	64.9	45.4	47.1	37.2	37.2
20	71.2	67.8	51.2	49.0	44.1 <sup>a</sup>	40.3
21	70.7	71.4	48.2	49.5	42.0	41.2
22	75.6	72.7	50.8	50.2	43.5	42.9
23	73.1 <sup>a</sup>	69.6	48.9	47.7	41.6	40.6
24	73.4	72.7	49.0	47.0	42.1	39.7
25	71.7	73.0	45.2	46.7	38.6	40.5
26-29	73.1	72.0	42.7	42.9	36.8	36.8
30-34	71.7	69.6	35.4	34.3	30.9	29.0
35-39	74.2	72.9	32.9	34.1	29.0	29.6
40-44	75.6	77.0	34.0	34.2	31.1	30.4
45-49	76.8	76.4	31.7	30.8	28.9	28.1
50-54	75.9	77.1	27.4	26.9	25.0	24.6
55-59	77.2	74.7	23.5	22.9	21.8	20.8
60-64	77.4	74.8	18.3	19.8	16.5	18.2
65 or Older	67.2 <sup>a</sup>	63.3	11.8	10.7	10.0	9.2

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 2.21A Daily Cigarette Use in Lifetime and Past Month Smoking Status, by Detailed Age Category: Numbers in Thousands, 2003 and 2004**

Age Category	Daily Cigarette Use in Lifetime		Daily Cigarette Use in Lifetime and Cigarette Use in Past Month		Daily Cigarette Use in Lifetime but No Cigarette Use in Past Month		Percentage Who Did Not Smoke Cigarettes in the Past Month among Lifetime Daily Smokers	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	94,759	93,409	51,241	50,278	43,517	43,130	45.9	46.2
12	24	13	14	10	9	3	*	*
13	72	71	45	48	26	22	*	*
14	180	159	138	134	42	25	23.2	15.6
15	295	325	241	271	53	54	18.1	16.8
16	496	464	436	401	60	63	12.0	13.6
17	785 <sup>a</sup>	676	691 <sup>a</sup>	589	94	87	12.0	12.9
18	1,136	1,140	1,015	1,014	121	126	10.7	11.0
19	1,173	1,134	1,000	993	173	141	14.8	12.4
20	1,454	1,316	1,284	1,169	171	148	11.7	11.2
21	1,464	1,506	1,277	1,269	187	237	12.8	15.7
22	1,593	1,484	1,366	1,262	227	222	14.2	14.9
23	1,625	1,534	1,318	1,260	306	274	18.8	17.9
24	1,526	1,643	1,249	1,282	277 <sup>a</sup>	361	18.1	22.0
25	1,452	1,456	1,151	1,178	301	278	20.7	19.1
26-29	5,827	6,042	4,480	4,629	1,347	1,414	23.1	23.4
30-34	7,610 <sup>a</sup>	6,850	5,334	4,745	2,277	2,105	29.9	30.7
35-39	8,178	7,935	5,435	5,307	2,743	2,628	33.5	33.1
40-44	10,132	10,037	6,351	6,338	3,782	3,699	37.3	36.9
45-49	10,161	10,137	5,688	5,613	4,473	4,524	44.0	44.6
50-54	9,556	9,475	4,631	4,377	4,926	5,097	51.5	53.8
55-59	7,739	8,220	3,041	3,294	4,698	4,925	60.7	59.9
60-64	6,135	6,518	1,767	2,199	4,368	4,319	71.2	66.3
65 or Older	16,145	15,274	3,289	2,897	12,856	12,378	79.6	81.0

\*Low precision; no estimate reported.

NOTE: Daily Cigarette Use in Lifetime is defined as ever smoking every day for at least 30 days.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 2.21B Daily Cigarette Use in Lifetime and Past Month Smoking Status, by Detailed Age Category: Percentages, 2003 and 2004**

Age Category	Daily Cigarette Use in Lifetime		Daily Cigarette Use in Lifetime and Cigarette Use in Past Month		Daily Cigarette Use in Lifetime but No Cigarette Use in Past Month		Percentage Who Did Not Smoke Cigarettes in the Past Month among Lifetime Daily Smokers	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	39.9 <sup>a</sup>	38.8	21.6	20.9	18.3	17.9	45.9	46.2
12	0.6	0.3	0.3	0.3	0.2	0.1	*	*
13	1.7	1.6	1.0	1.1	0.6	0.5	*	*
14	4.3	3.6	3.3	3.1	1.0 <sup>a</sup>	0.6	23.2	15.6
15	7.1	7.6	5.8	6.3	1.3	1.3	18.1	16.8
16	11.8	11.3	10.4	9.7	1.4	1.5	12.0	13.6
17	19.2 <sup>a</sup>	16.6	16.9 <sup>a</sup>	14.5	2.3	2.2	12.0	12.9
18	24.7	24.8	22.0	22.1	2.6	2.7	10.7	11.0
19	30.2	28.8	25.7	25.2	4.5	3.6	14.8	12.4
20	36.1	33.0	31.9	29.3	4.2	3.7	11.7	11.2
21	37.3	36.9	32.5	31.1	4.8	5.8	12.8	15.7
22	39.7	36.9	34.0	31.4	5.6	5.5	14.2	14.9
23	40.6	39.1	32.9	32.1	7.6	7.0	18.8	17.9
24	42.0	40.6	34.4	31.7	7.6	8.9	18.1	22.0
25	40.0	40.5	31.7	32.7	8.3	7.7	20.7	19.1
26-29	39.5	39.4	30.3	30.2	9.1	9.2	23.1	23.4
30-34	37.7	34.9	26.4	24.2	11.3	10.7	29.9	30.7
35-39	39.1	37.9	26.0	25.3	13.1	12.5	33.5	33.1
40-44	44.7	44.3	28.0	27.9	16.7	16.3	37.3	36.9
45-49	47.4	47.1	26.6	26.1	20.9	21.0	44.0	44.6
50-54	47.8	48.9	23.2	22.6	24.6	26.3	51.5	53.8
55-59	51.9	50.1	20.4	20.1	31.5	30.0	60.7	59.9
60-64	52.2	52.0	15.0	17.5	37.2	34.4	71.2	66.3
65 or Older	47.1	44.0	9.6	8.4	37.5	35.7	79.6	81.0

\*Low precision; no estimate reported.

NOTE: Daily Cigarette Use in Lifetime is defined as ever smoking every day for at least 30 days.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 2.22A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	46,065	44,625	10,347	9,756	7,725	7,154
12	75	76	25	39	11	13
13	177	139	79	78	33	36
14	236	223	135	131	57	61
15	332	364	200	220	98	108
16	463	431	256	281	121 <sup>a</sup>	164
17	615	558	339	335	191	188
18	739	817	386	445	198 <sup>a</sup>	264
19	721	685	321	354	175	194
20	887	777	390	333	208	190
21	875	934	302	314	184	208
22	945	928	295	318	184	188
23	981	884	320	253	208	172
24	964	978	273	293	197	196
25	864	874	198	252	131	177
26-29	4,092	4,159	1,049	955	789	686
30-34	5,749 <sup>b</sup>	4,978	1,314 <sup>a</sup>	1,010	1,054 <sup>a</sup>	797
35-39	5,658	5,442	1,247 <sup>a</sup>	939	1,025 <sup>a</sup>	759
40-44	5,161	4,940	853	779	748	654
45-49	3,992	4,382	527	589	430	447
50-54	3,155	3,108	395	418	343	342
55-59	2,682	2,511	438	377	417	308
60-64	1,709	1,717	267	267	214	228
65 or Older	4,993	4,719	738	777	707	774

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 2.22B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	19.4 <sup>a</sup>	18.6	4.4	4.1	3.3	3.0
12	1.8	1.9	0.6	1.0	0.3	0.3
13	4.1	3.2	1.8	1.8	0.8	0.8
14	5.7	5.1	3.3	3.0	1.4	1.4
15	8.0	8.5	4.8	5.1	2.4	2.5
16	11.1	10.5	6.1	6.8	2.9 <sup>a</sup>	4.0
17	15.0	13.7	8.3	8.2	4.7	4.6
18	16.0	17.8	8.4	9.7	4.3 <sup>a</sup>	5.7
19	18.6	17.4	8.3	9.0	4.5	4.9
20	22.0	19.5	9.7	8.4	5.2	4.8
21	22.3	22.9	7.7	7.7	4.7	5.1
22	23.5	23.1	7.3	7.9	4.6	4.7
23	24.5	22.5	8.0	6.4	5.2	4.4
24	26.5	24.2	7.5	7.2	5.4	4.9
25	23.8	24.3	5.5	7.0	3.6	4.9
26-29	27.7	27.1	7.1	6.2	5.3	4.5
30-34	28.5 <sup>a</sup>	25.4	6.5 <sup>a</sup>	5.1	5.2	4.1
35-39	27.0	26.0	6.0 <sup>a</sup>	4.5	4.9 <sup>a</sup>	3.6
40-44	22.7	21.8	3.8	3.4	3.3	2.9
45-49	18.6	20.4	2.5	2.7	2.0	2.1
50-54	15.8	16.0	2.0	2.2	1.7	1.8
55-59	18.0	15.3	2.9	2.3	2.8	1.9
60-64	14.5	13.7	2.3	2.1	1.8	1.8
65 or Older	14.6	13.6	2.2	2.2	2.1	2.2

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.



**Table 2.23A Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	88,096	87,409	25,386	25,974	12,837	13,727
12	130 <sup>a</sup>	81	69	44	21	26
13	243	252	123	141	45	53
14	458	452	292	285	115	125
15	695	748	471	494	206	204
16	979	943	670	695	331	335
17	1,272	1,267	864	911	396	458
18	1,682	1,779	1,148 <sup>a</sup>	1,330	576 <sup>b</sup>	766
19	1,619	1,631	999	1,102	516	595
20	1,882	1,768	1,043	1,094	561	529
21	1,759	1,815	863	962	450	541
22	1,977	1,904	939	868	453	439
23	1,900	1,752	782	809	386	421
24	1,770	1,909	729	844	355	437
25	1,738	1,654	693	716	325	375
26-29	6,707	6,942	2,237	2,447	1,214	1,196
30-34	8,196	7,808	2,537	2,342	1,206	1,087
35-39	8,333	8,350	2,383	2,294	1,052	1,154
40-44	8,910	9,076	2,475	2,277	1,322	1,204
45-49	9,042	8,824	2,035	2,296	1,091	1,293
50-54	7,926	7,809	1,718	1,399	869	864
55-59	6,178	6,530	1,088	1,196	547	741
60-64	4,244	4,911	423	639	250	388
65 or Older	10,456	9,203	805	791	549	497

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 2.23B Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	37.1	36.3	10.7	10.8	5.4	5.7
12	3.1 <sup>a</sup>	2.0	1.7	1.1	0.5	0.7
13	5.6	5.7	2.8	3.2	1.0	1.2
14	11.1	10.3	7.1	6.5	2.8	2.9
15	16.8	17.4	11.4	11.5	5.0	4.8
16	23.4	22.9	16.0	16.9	7.9	8.1
17	31.1	31.2	21.1	22.4	9.7	11.3
18	36.5	38.7	24.9 <sup>b</sup>	28.9	12.5 <sup>b</sup>	16.7
19	41.6	41.4	25.7	28.0	13.3	15.1
20	46.7	44.3	25.9	27.4	13.9	13.3
21	44.8	44.5	22.0	23.6	11.5	13.3
22	49.2	47.4	23.4	21.6	11.3	10.9
23	47.5	44.6	19.5	20.6	9.7	10.7
24	48.7	47.2	20.1	20.9	9.8	10.8
25	47.9	46.0	19.1	19.9	9.0	10.4
26-29	45.4	45.2	15.2	15.9	8.2	7.8
30-34	40.6	39.8	12.6	11.9	6.0	5.5
35-39	39.8	39.9	11.4	11.0	5.0	5.5
40-44	39.3	40.0	10.9	10.0	5.8	5.3
45-49	42.2	41.0	9.5	10.7	5.1	6.0
50-54	39.6	40.3	8.6	7.2	4.3	4.5
55-59	41.4	39.8	7.3	7.3	3.7	4.5
60-64	36.1	39.2	3.6	5.1	2.1	3.1
65 or Older	30.5 <sup>a</sup>	26.5	2.3	2.3	1.6	1.4

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 2.24A Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	197,533	198,220	154,540	156,686	118,965	120,934
12	531 <sup>a</sup>	441	315	260	119	92
13	1,006	969	713	685	263	270
14	1,463	1,598	1,126	1,247	542	540
15	2,181	2,178	1,756	1,773	863	878
16	2,634	2,493	2,211	2,102	1,195	1,142
17	2,898	2,916	2,443	2,469	1,443	1,515
18	3,523	3,514	3,053	3,120	2,012	2,130
19	3,199	3,204	2,801	2,864	2,038	2,027
20	3,501	3,395	3,137	3,047	2,401	2,246
21	3,522	3,681	3,271	3,436	2,734	2,846
22	3,669	3,612	3,359	3,362	2,803	2,776
23	3,610	3,460	3,295	3,118	2,699	2,504
24	3,300 <sup>a</sup>	3,632	2,920 <sup>b</sup>	3,250	2,380	2,580
25	3,299	3,254	2,942	2,931	2,400	2,365
26-29	13,307	13,908	11,620	12,046	9,107	9,705
30-34	18,268	17,573	15,686	14,775	11,954	11,457
35-39	19,093	18,984	15,709	15,806	12,456	12,687
40-44	20,706	20,747	16,731	17,017	13,293	13,623
45-49	19,435	19,687	15,234	15,359	12,348	12,355
50-54	18,061	17,304	13,993	13,368	10,796	10,779
55-59	13,244	14,440	9,864	10,289	7,886	7,922
60-64	10,093	10,675	6,823	7,852	5,432	6,255
65 or Older	26,992	26,554	15,541	16,511	11,801	12,242

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 2.24B Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	83.1	82.4	65.0	65.1	50.1	50.3
12	12.9	11.1	7.6	6.6	2.9	2.3
13	23.2	22.1	16.4	15.6	6.1	6.1
14	35.4	36.5	27.3	28.5	13.1	12.3
15	52.8	50.7	42.5	41.3	20.9	20.4
16	62.8	60.6	52.7	51.1	28.5	27.7
17	70.8	71.7	59.7	60.7	35.3	37.3
18	76.5	76.5	66.3	67.9	43.7	46.3
19	82.3	81.4	72.1	72.7	52.4	51.5
20	86.9	85.1	77.9	76.4	59.6	56.3
21	89.8	90.2	83.4	84.2	69.7	69.8
22	91.3	89.9	83.7	83.6	69.8	69.1
23	90.2	88.1	82.3 <sup>a</sup>	79.4	67.4 <sup>a</sup>	63.8
24	90.8	89.7	80.3	80.3	65.5	63.7
25	90.9	90.4	81.1	81.4	66.1	65.7
26-29	90.1	90.6	78.7	78.5	61.7	63.2
30-34	90.5	89.5	77.7	75.3	59.2	58.4
35-39	91.2	90.6	75.1	75.5	59.5	60.6
40-44	91.3	91.5	73.7	75.0	58.6	60.1
45-49	90.8	91.5	71.1	71.4	57.7	57.5
50-54	90.3	89.3	70.0	69.0	54.0	55.6
55-59	88.8	88.0	66.1	62.7	52.9	48.3
60-64	85.9	85.1	58.0	62.6	46.2	49.9
65 or Older	78.7	76.6	45.3	47.6	34.4	35.3

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 2.25A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Numbers in Thousands, 2003 and 2004**

Age Category	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	118,965	120,934	53,770	54,725	16,144	16,689
12	119	92	36	42	5	4
13	263	270	95	129	7	16
14	542	540	292	244	52	43
15	863	878	482	544	129	93
16	1,195	1,142	753	754	180	187
17	1,443	1,515	1,001	1,080	280	328
18	2,012	2,130	1,454	1,522	464	507
19	2,038	2,027	1,412	1,472	535	548
20	2,401	2,246	1,666	1,610	647	649
21	2,734	2,846	1,876	1,965	735	781
22	2,803	2,776	1,886	1,882	699	715
23	2,699	2,504	1,845	1,694	662	562
24	2,380	2,580	1,598	1,632	556	552
25	2,400	2,365	1,452	1,494	492	533
26-29	9,107	9,705	5,615	5,643	1,687	1,648
30-34	11,954	11,457	5,883	5,623	1,602	1,627
35-39	12,456	12,687	5,871	6,236	1,652	1,523
40-44	13,293	13,623	5,821	6,099	1,785	1,894
45-49	12,348	12,355	4,968	4,986	1,463	1,483
50-54	10,796	10,779	3,582	3,663	975	1,210
55-59	7,886	7,922	2,318	2,309	633	701
60-64	5,432	6,255	1,400	1,705	297	467
65 or Older	11,801	12,242	2,464	2,398	611	616

\*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 2.25B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Percentages, 2003 and 2004**

Age Category	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	50.1	50.3	22.6	22.8	6.8	6.9
12	2.9	2.3	0.9	1.1	0.1	0.1
13	6.1	6.1	2.2	2.9	0.2	0.4
14	13.1	12.3	7.1 <sup>a</sup>	5.6	1.3	1.0
15	20.9	20.4	11.7	12.7	3.1 <sup>a</sup>	2.2
16	28.5	27.7	18.0	18.3	4.3	4.5
17	35.3	37.3	24.5	26.6	6.8	8.1
18	43.7	46.3	31.5	33.1	10.1	11.0
19	52.4	51.5	36.3	37.4	13.8	13.9
20	59.6	56.3	41.4	40.3	16.1	16.3
21	69.7	69.8	47.8	48.2	18.7	19.2
22	69.8	69.1	47.0	46.8	17.4	17.8
23	67.4 <sup>a</sup>	63.8	46.1	43.1	16.5	14.3
24	65.5	63.7	44.0 <sup>a</sup>	40.3	15.3	13.7
25	66.1	65.7	40.0	41.5	13.6	14.8
26-29	61.7	63.2	38.0	36.8	11.4	10.7
30-34	59.2	58.4	29.1	28.6	7.9	8.3
35-39	59.5	60.6	28.1	29.8	7.9	7.3
40-44	58.6	60.1	25.7	26.9	7.9	8.4
45-49	57.7	57.5	23.2	23.2	6.8	6.9
50-54	54.0	55.6	17.9	18.9	4.9	6.2
55-59	52.9	48.3	15.5	14.1	4.2	4.3
60-64	46.2	49.9	11.9	13.6	2.5	3.7
65 or Older	34.4	35.3	7.2	6.9	1.8	1.8

\*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.