

**Table 1.58C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004**

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	641	729	354	353	250	210
<b>AGE</b>						
12-17	76	70	64	59	42	40
18-25	135	141	103	110	70	68
26 or Older	617	700	331	313	237	187
<b>GENDER</b>						
Male	468	493	236	238	155	142
Female	448	474	247	223	175	141
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	606	673	323	329	211	197
White	548	631	291	319	184	184
Black or African American	196	173	103	78	64	60
American Indian or Alaska Native	46	47	19	28	17	10
Native Hawaiian or Other Pacific Islander	16	*	10	*	6	14
Asian	103	125	65	51	54	23
Two or More Races	63	70	19	38	12	33
Hispanic or Latino	214	218	144	99	106	66

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.58D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2003 and 2004**

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.27	0.30	0.15	0.15	0.11	0.09
<b>AGE</b>						
12-17	0.30	0.28	0.26	0.23	0.17	0.16
18-25	0.43	0.44	0.33	0.34	0.22	0.21
26 or Older	0.34	0.38	0.18	0.17	0.13	0.10
<b>GENDER</b>						
Male	0.41	0.42	0.20	0.20	0.13	0.12
Female	0.37	0.38	0.20	0.18	0.14	0.11
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.29	0.32	0.16	0.16	0.10	0.09
White	0.33	0.38	0.18	0.19	0.11	0.11
Black or African American	0.71	0.62	0.38	0.28	0.23	0.21
American Indian or Alaska Native	3.66	3.61	1.51	2.19	1.32	0.76
Native Hawaiian or Other Pacific Islander	3.16	*	2.11	*	1.29	2.26
Asian	1.05	1.26	0.66	0.51	0.55	0.23
Two or More Races	2.46	2.73	0.76	1.50	0.46	1.31
Hispanic or Latino	0.72	0.70	0.48	0.32	0.35	0.21

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.58P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.6481		0.2977		0.2207	
<b>AGE</b>						
12-17	0.9027		0.3390		0.1073	
18-25	0.8095		0.5653		0.8520	
26 or Older	0.5784		0.2500		0.2811	
<b>GENDER</b>						
Male	0.6249		0.5632		0.4969	
Female	0.2294		0.3212		0.2302	
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.8312		0.9331		0.4537	
White	0.9509		0.8575		0.6055	
Black or African American	0.2338		0.1354		0.4489	
American Indian or Alaska Native	0.8681		0.6094		0.2283	
Native Hawaiian or Other Pacific Islander	*		*		0.5765	
Asian	0.7910		0.2650		0.1400	
Two or More Races	0.3536		0.0090		0.0167	
Hispanic or Latino	0.4982		0.0123		0.1518	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.59C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004**

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	76	70	64	59	42	40
<b>GENDER</b>						
Male	49	48	43	39	28	26
Female	53	49	46	45	31	31
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	70	68	60	55	38	35
White	61	58	52	49	33	31
Black or African American	29	25	22	20	15	13
American Indian or Alaska Native	5	5	5	5	3	4
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	12	13	10	10	9	9
Two or More Races	8	11	6	9	5	6
Hispanic or Latino	32	32	27	25	19	20
<b>GENDER/RACE/HISPANIC ORIGIN</b>						
Male, White, Not Hispanic	39	38	33	33	22	20
Female, White, Not Hispanic	42	39	37	35	23	23
Male, Black, Not Hispanic	18	17	14	12	9	9
Female, Black, Not Hispanic	20	19	17	17	13	10
Male, Hispanic	22	23	20	17	12	15
Female, Hispanic	25	23	20	20	13	15

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.59D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2003 and 2004**

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.30	0.28	0.26	0.23	0.17	0.16
<b>GENDER</b>						
Male	0.38	0.37	0.33	0.30	0.22	0.20
Female	0.44	0.40	0.38	0.36	0.25	0.25
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.34	0.32	0.29	0.26	0.18	0.16
White	0.39	0.37	0.34	0.32	0.21	0.20
Black or African American	0.77	0.67	0.60	0.54	0.41	0.34
American Indian or Alaska Native	2.64	3.21	2.61	2.97	1.61	2.68
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.23	1.30	1.00	1.03	0.89	0.86
Two or More Races	1.93	2.59	1.50	2.05	1.18	1.28
Hispanic or Latino	0.80	0.76	0.65	0.60	0.46	0.48
<b>GENDER/RACE/HISPANIC ORIGIN</b>						
Male, White, Not Hispanic	0.48	0.49	0.42	0.42	0.28	0.25
Female, White, Not Hispanic	0.55	0.51	0.49	0.46	0.31	0.30
Male, Black, Not Hispanic	0.98	0.89	0.76	0.61	0.46	0.45
Female, Black, Not Hispanic	1.10	1.01	0.90	0.90	0.69	0.53
Male, Hispanic	1.08	1.07	0.94	0.78	0.59	0.70
Female, Hispanic	1.25	1.10	1.01	0.97	0.67	0.75

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.59P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.9027		0.3390		0.1073	
<b>GENDER</b>						
Male	0.6606		0.0345		0.0758	
Female	0.5546		0.5376		0.6100	
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.6487		0.4545		0.0363	
White	0.3191		0.9368		0.1551	
Black or African American	0.0789		0.0232		0.1055	
American Indian or Alaska Native	0.1588		0.5310		0.7741	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.9428		0.8475		0.7200	
Two or More Races	0.0634		0.3576		0.9836	
Hispanic or Latino	0.4863		0.5453		0.5320	
<b>GENDER/RACE/HISPANIC ORIGIN</b>						
Male, White, Not Hispanic	0.9032		0.4005		0.0488	
Female, White, Not Hispanic	0.1752		0.3661		0.8793	
Male, Black, Not Hispanic	0.1220		0.0750		0.6554	
Female, Black, Not Hispanic	0.3159		0.1469		0.0858	
Male, Hispanic	0.5955		0.1628		0.5922	
Female, Hispanic	0.6689		0.6038		0.7361	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.60C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004**

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	135	141	103	110	70	68
<b>GENDER</b>						
Male	91	100	73	81	54	51
Female	88	85	67	67	40	47
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	121	127	98	97	65	58
White	104	111	85	87	61	55
Black or African American	38	39	26	26	15	17
American Indian or Alaska Native	8	*	6	*	5	3
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	26	24	17	18	12	9
Two or More Races	12	13	10	11	8	7
Hispanic or Latino	56	57	37	41	26	30
<b>EDUCATION</b>						
< High School	58	63	47	48	33	35
High School Graduate	78	76	58	53	39	39
Some College	71	85	56	72	37	41
College Graduate	44	44	31	32	20	19
<b>CURRENT EMPLOYMENT</b>						
Full-Time	88	95	66	66	48	41
Part-Time	58	69	50	54	33	32
Unemployed	36	40	31	33	22	24
Other <sup>1</sup>	55	64	38	50	26	33

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

<sup>1</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.60D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2003 and 2004**

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.43	0.44	0.33	0.34	0.22	0.21
<b>GENDER</b>						
Male	0.57	0.62	0.46	0.50	0.34	0.32
Female	0.56	0.53	0.43	0.42	0.25	0.29
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.46	0.48	0.37	0.37	0.25	0.22
White	0.53	0.56	0.43	0.44	0.31	0.28
Black or African American	0.88	0.89	0.60	0.59	0.36	0.39
American Indian or Alaska Native	4.42	*	3.48	*	3.00	1.93
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.79	1.58	1.15	1.20	0.82	0.59
Two or More Races	3.37	3.68	2.88	3.08	2.07	1.95
Hispanic or Latino	0.99	1.00	0.66	0.72	0.46	0.53
<b>EDUCATION</b>						
< High School	0.84	0.94	0.68	0.71	0.49	0.52
High School Graduate	0.71	0.69	0.53	0.48	0.36	0.35
Some College	0.72	0.83	0.57	0.69	0.37	0.39
College Graduate	1.10	1.06	0.78	0.76	0.51	0.45
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.60	0.63	0.46	0.44	0.33	0.27
Part-Time	0.70	0.85	0.61	0.66	0.41	0.40
Unemployed	1.35	1.53	1.16	1.29	0.83	0.95
Other <sup>1</sup>	0.88	1.00	0.61	0.78	0.42	0.51

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

<sup>1</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.



**Table 1.60P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.8095		0.5653		0.8520	
<b>GENDER</b>						
Male	0.2549		0.3247		0.4397	
Female	0.3525		0.7836		0.2177	
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.5280		0.2954		0.9484	
White	0.9726		0.5071		0.9185	
Black or African American	0.0571		0.0416		0.2090	
American Indian or Alaska Native	*		*		0.0787	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.3914		0.4750		0.1659	
Two or More Races	0.0459		0.4087		0.5380	
Hispanic or Latino	0.4172		0.2990		0.5755	
<b>EDUCATION</b>						
< High School	0.9192		0.9402		0.5791	
High School Graduate	0.8920		0.9701		0.5847	
Some College	0.2442		0.1145		0.2958	
College Graduate	0.3047		0.3803		0.2192	
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.8267		0.9320		0.7523	
Part-Time	0.9420		0.9066		0.5978	
Unemployed	0.8068		0.5381		0.8414	
Other <sup>1</sup>	0.9785		0.1832		0.1314	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

<sup>1</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.61C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004**

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	617	700	331	313	237	187
<b>GENDER</b>						
Male	453	474	219	213	146	128
Female	432	458	234	198	169	125
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	587	645	298	299	201	179
White	529	607	267	292	171	168
Black or African American	185	164	95	70	61	55
American Indian or Alaska Native	44	44	16	25	15	8
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	95	128	61	50	52	19
Two or More Races	62	66	13	34	7	31
Hispanic or Latino	201	208	140	87	99	54
<b>EDUCATION</b>						
< High School	243	238	128	124	98	78
High School Graduate	311	339	166	163	121	102
Some College	312	347	168	156	120	95
College Graduate	359	350	158	160	95	83
<b>CURRENT EMPLOYMENT</b>						
Full-Time	486	510	235	254	137	147
Part-Time	208	213	114	89	92	54
Unemployed	106	125	74	70	60	54
Other <sup>1</sup>	273	283	159	130	129	79

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

<sup>1</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.61D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2003 and 2004**

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.34	0.38	0.18	0.17	0.13	0.10
<b>GENDER</b>						
Male	0.53	0.54	0.25	0.24	0.17	0.15
Female	0.46	0.48	0.25	0.21	0.18	0.13
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.37	0.40	0.19	0.18	0.12	0.11
White	0.40	0.46	0.20	0.22	0.13	0.13
Black or African American	0.95	0.83	0.49	0.35	0.31	0.28
American Indian or Alaska Native	4.90	4.62	1.84	2.61	1.72	0.83
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.29	1.73	0.83	0.67	0.70	0.25
Two or More Races	3.44	3.75	0.70	1.92	0.41	1.74
Hispanic or Latino	1.00	0.99	0.69	0.41	0.49	0.26
<b>EDUCATION</b>						
< High School	0.78	0.80	0.41	0.42	0.32	0.26
High School Graduate	0.55	0.59	0.30	0.29	0.22	0.18
Some College	0.73	0.78	0.39	0.35	0.28	0.21
College Graduate	0.70	0.68	0.31	0.31	0.19	0.16
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.47	0.49	0.23	0.24	0.13	0.14
Part-Time	1.02	1.09	0.56	0.46	0.45	0.28
Unemployed	1.97	2.60	1.38	1.45	1.11	1.12
Other <sup>1</sup>	0.52	0.52	0.30	0.24	0.25	0.14

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

<sup>1</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.61P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.5784		0.2500		0.2811	
<b>GENDER</b>						
Male	0.7590		0.5475		0.9254	
Female	0.2538		0.2650		0.1270	
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.6623		0.7497		0.6886	
White	0.8245		0.9369		0.7746	
Black or African American	0.1687		0.1004		0.4391	
American Indian or Alaska Native	0.9708		0.5885		0.4953	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.6439		0.3832		0.2934	
Two or More Races	0.7727		0.0121		0.0129	
Hispanic or Latino	0.7248		0.0284		0.0569	
<b>EDUCATION</b>						
< High School	0.5932		0.7518		0.6998	
High School Graduate	0.4294		0.4699		0.2331	
Some College	0.4539		0.0438		0.0777	
College Graduate	0.8459		0.8302		0.5196	
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.4274		0.4878		0.2731	
Part-Time	0.3070		0.4214		0.0485	
Unemployed	0.1759		0.7938		0.4735	
Other <sup>1</sup>	0.4150		0.5111		0.0162	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

<sup>1</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.62C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004**

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	635	724	351	341	249	203
<b>GENDER</b>						
Male	466	489	234	231	154	139
Female	443	469	246	213	173	135
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	602	669	320	323	211	191
White	543	630	288	313	184	180
Black or African American	192	171	100	75	62	58
American Indian or Alaska Native	45	48	18	28	16	9
Native Hawaiian or Other Pacific Islander	15	*	9	*	5	14
Asian	102	127	64	53	53	20
Two or More Races	63	70	18	38	11	33
Hispanic or Latino	213	212	144	94	104	62
<b>EDUCATION</b>						
< High School	257	255	141	135	106	86
High School Graduate	321	348	176	174	125	110
Some College	316	368	169	179	125	108
College Graduate	362	354	165	165	98	85
<b>CURRENT EMPLOYMENT</b>						
Full-Time	495	519	243	263	147	153
Part-Time	215	225	122	111	97	62
Unemployed	112	131	77	79	63	59
Other <sup>1</sup>	282	291	166	143	131	87

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

<sup>1</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.62D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2003 and 2004**

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.30	0.34	0.17	0.16	0.12	0.09
<b>GENDER</b>						
Male	0.46	0.47	0.23	0.22	0.15	0.13
Female	0.40	0.42	0.22	0.19	0.16	0.12
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.32	0.36	0.17	0.17	0.11	0.10
White	0.36	0.42	0.19	0.21	0.12	0.12
Black or African American	0.81	0.71	0.42	0.31	0.26	0.24
American Indian or Alaska Native	4.19	4.23	1.65	2.43	1.52	0.76
Native Hawaiian or Other Pacific Islander	3.57	*	2.23	*	1.09	2.50
Asian	1.16	1.42	0.72	0.59	0.60	0.23
Two or More Races	2.91	3.32	0.83	1.80	0.50	1.56
Hispanic or Latino	0.82	0.79	0.56	0.35	0.40	0.23
<b>EDUCATION</b>						
< High School	0.68	0.70	0.37	0.37	0.28	0.23
High School Graduate	0.48	0.51	0.26	0.26	0.19	0.16
Some College	0.60	0.67	0.32	0.33	0.24	0.20
College Graduate	0.66	0.63	0.30	0.30	0.18	0.15
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.42	0.44	0.21	0.22	0.13	0.13
Part-Time	0.75	0.82	0.42	0.40	0.34	0.23
Unemployed	1.39	1.77	0.96	1.07	0.78	0.80
Other <sup>1</sup>	0.48	0.47	0.28	0.23	0.22	0.14

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

<sup>1</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.62P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.6292		0.3762		0.3400	
<b>GENDER</b>						
Male	0.5970		0.8293		0.7256	
Female	0.1982		0.2611		0.2633	
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.7818		0.9542		0.7093	
White	0.8594		0.8584		0.7938	
Black or African American	0.3412		0.3273		0.6897	
American Indian or Alaska Native	0.9895		0.6658		0.1992	
Native Hawaiian or Other Pacific Islander	*		*		0.2344	
Asian	0.7955		0.2770		0.1449	
Two or More Races	0.4815		0.0147		0.0140	
Hispanic or Latino	0.5549		0.0143		0.1070	
<b>EDUCATION</b>						
< High School	0.5854		0.7697		0.8562	
High School Graduate	0.4112		0.4771		0.3298	
Some College	0.6434		0.1803		0.1984	
College Graduate	0.7585		0.9400		0.7001	
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.4769		0.5353		0.3189	
Part-Time	0.3688		0.5364		0.0528	
Unemployed	0.1565		0.5436		0.4194	
Other <sup>1</sup>	0.4497		0.7669		0.0490	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

<sup>1</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.