

**Table 7.70A Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004**

Drug	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>ILLCIT DRUG<sup>2</sup></b>	6,300	6,193	105	118	6,152	6,031
Marijuana and Hashish	4,575	4,525	86	92	4,452	4,397
Cocaine	601	630	8	8	587	617
Crack	116	119	1	*	115	118
Heroin	34	35	1	2	32	31
Hallucinogens	354 <sup>b</sup>	295	6	5	344 <sup>b</sup>	287
LSD	35	29	0	2	35	28
PCP	15	11	2	1	14	9
Ecstasy	223 <sup>b</sup>	176	4	4	217 <sup>b</sup>	171
Inhalants	131	129	1	2	130	126
Nonmedical Use of Psychotherapeutics <sup>3</sup>	2,329	2,330	30	35	2,285	2,280
Pain Relievers	1,630	1,690	21	30	1,596	1,645
OxyContin <sup>®</sup>	--	71	--	1	--	70
Tranquilizers	692	682	5	2	684	676
Stimulants	492	513	7	8	483	503
Methamphetamine	207	218	1	2	205	216
Sedatives	124	96	2	3	123	93
<b>ILLCIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	2,961	2,907	35	41	2,907	2,845

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.<sup>3</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 7.70B Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2002-2003 and 2003-2004**

Drug	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>ILLICIT DRUG<sup>2</sup></b>	10.2	10.0	4.3	4.6	10.4	10.2
Marijuana and Hashish	7.4	7.3	3.5	3.6	7.6	7.5
Cocaine	1.0	1.0	0.3	0.3	1.0	1.0
Crack	0.2	0.2	0.0	*	0.2	0.2
Heroin	0.1	0.1	0.0	0.1	0.1	0.1
Hallucinogens	0.6 <sup>b</sup>	0.5	0.3	0.2	0.6 <sup>b</sup>	0.5
LSD	0.1	0.0	0.0	0.1	0.1	0.0
PCP	0.0	0.0	0.1	0.1	0.0	0.0
Ecstasy	0.4 <sup>b</sup>	0.3	0.1	0.1	0.4 <sup>b</sup>	0.3
Inhalants	0.2	0.2	0.0	0.1	0.2	0.2
Nonmedical Use of Psychotherapeutics <sup>3</sup>	3.8	3.8	1.2	1.4	3.9	3.9
Pain Relievers	2.6	2.7	0.9	1.2	2.7	2.8
OxyContin <sup>®</sup>	--	0.2	--	0.1	--	0.2
Tranquilizers	1.1	1.1	0.2	0.1	1.2	1.1
Stimulants	0.8	0.8	0.3	0.3	0.8	0.9
Methamphetamine	0.3	0.4	0.0	0.1	0.3	0.4
Sedatives	0.2	0.2	0.1	0.1	0.2	0.2
<b>ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	4.8	4.7	1.4	1.6	4.9	4.8

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.<sup>3</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 7.71A Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004**

Demographic Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>TOTAL</b>	6,300	6,193	105	118	6,152	6,031
<b>AGE</b>						
15-17	994	978	12	15	975	953
18-25	2,573	2,540	71	71	2,476	2,446
26-44	2,733	2,674	22	32	2,701	2,631
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	5,632	5,538	90	94	5,502	5,405
White	4,534	4,414	62	66	4,447	4,321
Black or African American	780	773	24	23	746	742
American Indian or Alaska Native	60	66	*	*	59	65
Native Hawaiian or Other Pacific Islander	24	*	*	*	23	*
Asian	131	112	*	*	128	108
Two or More Races	102 <sup>a</sup>	146	*	*	99 <sup>a</sup>	143
Hispanic or Latino	668	654	15	24	651	626
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	54	62	N/A	N/A
Second	N/A	N/A	31	34	N/A	N/A
Third	N/A	N/A	18	21	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 7.71B Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Percentages, Annual Averages Based on 2002-2003 and 2003-2004**

Demographic Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>TOTAL</b>	10.2	10.0	4.3	4.6	10.4	10.2
<b>AGE</b>						
15-17	16.5	16.1	12.8	16.0	16.5	16.0
18-25	16.4	16.0	7.5	7.8	16.9	16.4
26-44	6.8	6.7	1.6	2.1	7.0	6.9
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	10.8	10.6	4.7	4.5	11.0	10.8
White	11.4	11.1	4.4	4.2	11.6	11.4
Black or African American	9.5	9.4	8.0	7.8	9.4	9.4
American Indian or Alaska Native	15.4	17.8	*	*	16.3	18.6
Native Hawaiian or Other Pacific Islander	12.6	*	*	*	12.8	*
Asian	4.3	3.5	*	*	4.4	3.6
Two or More Races	15.4 <sup>a</sup>	20.3	*	*	15.9 <sup>a</sup>	20.8
Hispanic or Latino	7.2	6.9	3.0	5.0	7.4	7.0
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	7.7	8.0	N/A	N/A
Second	N/A	N/A	3.2	3.8	N/A	N/A
Third	N/A	N/A	2.3	2.4	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 7.72A Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004**

Demographic Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>TOTAL</b>	4,575	4,525	86	92	4,452	4,397
<b>AGE</b>						
15-17	724	723	9	11	709	706
18-25	2,086	1,992	60	56	2,004	1,918
26-44	1,765	1,810	17	26	1,739	1,774
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	4,171	4,125	74	75	4,061	4,018
White	3,374	3,301	52	54	3,299	3,225
Black or African American	594	601	18	16	565	578
American Indian or Alaska Native	44	52	*	*	43	52
Native Hawaiian or Other Pacific Islander	10	10	*	*	10	10
Asian	78	69	*	*	74	66
Two or More Races	72	92	*	*	70	89
Hispanic or Latino	404	400	11	18	391	379
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	43	43	N/A	N/A
Second	N/A	N/A	25	29	N/A	N/A
Third	N/A	N/A	16	19	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 7.72B Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Percentages, Annual Averages Based on 2002-2003 and 2003-2004**

Demographic Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>TOTAL</b>	7.4	7.3	3.5	3.6	7.6	7.5
<b>AGE</b>						
15-17	12.0	11.9	9.5	12.1	12.0	11.9
18-25	13.3 <sup>b</sup>	12.5	6.3	6.1	13.7 <sup>b</sup>	12.9
26-44	4.4	4.5	1.2	1.7	4.5	4.7
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	8.0	7.9	3.9	3.6	8.1	8.0
White	8.5	8.3	3.7	3.5	8.6	8.5
Black or African American	7.2	7.3	6.2	5.3	7.1	7.3
American Indian or Alaska Native	11.3	14.1	*	*	12.0	14.7
Native Hawaiian or Other Pacific Islander	5.2	5.5	*	*	5.4	5.6
Asian	2.5	2.2	*	*	2.6	2.2
Two or More Races	11.0	12.8	*	*	11.2	12.9
Hispanic or Latino	4.3	4.2	2.2	3.7	4.5	4.2
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	6.1	5.6	N/A	N/A
Second	N/A	N/A	2.5	3.2	N/A	N/A
Third	N/A	N/A	2.1	2.2	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 7.73A Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004**

Substance	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>TOBACCO PRODUCT<sup>2</sup></b>	19,288	18,893	459	477	18,707	18,289
Cigarettes	18,655	18,237	438	460	18,110	17,664
Smokeless Tobacco	184	159	7	3	178	156
Cigars	1,876	1,934	35	35	1,815	1,870
Pipe Tobacco	160	159	8	8	152	151
<b>ALCOHOL</b>	31,647	31,572	238	285	31,246	31,135
Binge Alcohol Use <sup>3</sup>	13,849	13,897	99	114	13,676	13,708
Heavy Alcohol Use <sup>3</sup>	3,166	3,296	16	13	3,140	3,273

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 7.73B Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2002-2003 and 2003-2004**

Substance	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>TOBACCO PRODUCT<sup>2</sup></b>	31.3 <sup>a</sup>	30.5	18.9	18.7	31.7	31.0
Cigarettes	30.2 <sup>a</sup>	29.5	18.0	18.0	30.7 <sup>a</sup>	30.0
Smokeless Tobacco	0.3	0.3	0.3	0.1	0.3	0.3
Cigars	3.0	3.1	1.4	1.4	3.1	3.2
Pipe Tobacco	0.3	0.3	0.3	0.3	0.3	0.3
<b>ALCOHOL</b>	51.3	51.1	9.8	11.2	53.0	52.8
Binge Alcohol Use <sup>3</sup>	22.4	22.5	4.1	4.5	23.2	23.3
Heavy Alcohol Use <sup>3</sup>	5.1	5.3	0.7	0.5	5.3	5.6

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.



**Table 7.74A Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004**

Demographic Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>TOTAL</b>	18,655	18,237	438	460	18,110	17,664
<b>AGE</b>						
15-17	1,257	1,206	25	24	1,223	1,168
18-25	5,739	5,707	263	256	5,433	5,405
26-44	11,659	11,324	149	180	11,454	11,091
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	16,915	16,458	403	422	16,420	15,938
White	14,035	13,655	356	367	13,604	13,205
Black or African American	1,994	1,993	28	41	1,953	1,940
American Indian or Alaska Native	191	186	*	*	187	182
Native Hawaiian or Other Pacific Islander	69	*	*	*	67	*
Asian	372 <sup>a</sup>	270	*	*	368 <sup>a</sup>	267
Two or More Races	253	303	*	*	241 <sup>a</sup>	294
Hispanic or Latino	1,740	1,779	35	38	1,690	1,726
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	155	174	N/A	N/A
Second	N/A	N/A	150 <sup>a</sup>	120	N/A	N/A
Third	N/A	N/A	130	156	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 7.74B Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Percentages, Annual Averages Based on 2002-2003 and 2003-2004**

Demographic Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>TOTAL</b>	30.2 <sup>a</sup>	29.5	18.0	18.0	30.7 <sup>a</sup>	30.0
<b>AGE</b>						
15-17	20.9 <sup>a</sup>	19.8	25.9	26.0	20.7 <sup>a</sup>	19.6
18-25	36.7	35.9	27.8	28.0	37.2 <sup>a</sup>	36.3
26-44	29.1	28.4	10.8	11.7	29.8	29.1
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	32.3 <sup>a</sup>	31.5	21.0	20.4	32.7 <sup>a</sup>	31.9
White	35.2	34.5	25.0	23.6	35.6	34.9
Black or African American	24.2	24.2	9.3	13.6	24.7	24.6
American Indian or Alaska Native	49.2	49.9	*	*	52.0	51.8
Native Hawaiian or Other Pacific Islander	36.5	*	*	*	36.7	*
Asian	12.2 <sup>b</sup>	8.5	*	*	12.8 <sup>b</sup>	8.9
Two or More Races	38.4	42.2	*	*	38.6	42.6
Hispanic or Latino	18.7	18.7	6.8	8.0	19.3	19.2
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	22.0	22.7	N/A	N/A
Second	N/A	N/A	15.5	13.4	N/A	N/A
Third	N/A	N/A	17.3	18.0	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 7.75A Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004**

Demographic Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>TOTAL</b>	31,647	31,572	238	285	31,246	31,135
<b>AGE</b>						
15-17	1,717	1,729	14	14	1,694	1,700
18-25	8,722	8,894	100	97	8,577	8,743
26-44	21,208	20,948	124	174	20,975	20,692
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	28,113	27,987	194	232	27,783	27,638
White	23,042	23,059	154	184	22,802	22,792
Black or African American	3,384	3,260	19	29	3,336	3,201
American Indian or Alaska Native	194	184	*	*	186	178
Native Hawaiian or Other Pacific Islander	78	*	*	*	77	*
Asian	1,063	1,008	*	*	1,036	996
Two or More Races	352	413	*	*	346 <sup>a</sup>	408
Hispanic or Latino	3,534	3,585	44	53	3,464	3,498
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	138	170	N/A	N/A
Second	N/A	N/A	59	62	N/A	N/A
Third	N/A	N/A	35	43	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 7.75B Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Percentages, Annual Averages Based on 2002-2003 and 2003-2004**

Demographic Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>TOTAL</b>	51.3	51.1	9.8	11.2	53.0	52.8
<b>AGE</b>						
15-17	28.5	28.4	14.5	14.9	28.7	28.5
18-25	55.7	55.9	10.5	10.6	58.7	58.7
26-44	53.0	52.6	8.9	11.3	54.6	54.3
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	53.7	53.5	10.1	11.2	55.3	55.3
White	57.8	58.2	10.8	11.8	59.6	60.2
Black or African American	41.0	39.5	6.4	9.6	42.2	40.6
American Indian or Alaska Native	49.8	49.5	*	*	51.7	50.6
Native Hawaiian or Other Pacific Islander	40.9	*	*	*	42.1	*
Asian	34.9	31.8	*	*	36.0	33.3
Two or More Races	53.3	57.4	*	*	55.6	59.2
Hispanic or Latino	37.9	37.6	8.6	11.0	39.6	39.0
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	19.6	22.2	N/A	N/A
Second	N/A	N/A	6.1	7.0	N/A	N/A
Third	N/A	N/A	4.7	4.9	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 7.76A Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004**

Demographic Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>TOTAL</b>	13,849	13,897	99	114	13,676	13,708
<b>AGE</b>						
15-17	999	1,051	7	8	985	1,031
18-25	4,968	5,097	46	47	4,893	5,018
26-44	7,882	7,749	46	59	7,798	7,659
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	12,194	12,175	84	96	12,050	12,020
White	10,065	10,108	57	63	9,975	10,012
Black or African American	1,513	1,422	9	17	1,481	1,382
American Indian or Alaska Native	134	140	*	*	126	135
Native Hawaiian or Other Pacific Islander	*	27	*	*	*	26
Asian	274	287	*	*	263	277
Two or More Races	156	191	*	*	154	189
Hispanic or Latino	1,655	1,722	15	18	1,626	1,688
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	77	81	N/A	N/A
Second	N/A	N/A	13	17	N/A	N/A
Third	N/A	N/A	5	9	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 7.76B Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Percentages, Annual Averages Based on 2002-2003 and 2003-2004**

Demographic Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>TOTAL</b>	22.4	22.5	4.1	4.5	23.2	23.3
<b>AGE</b>						
15-17	16.6	17.3	7.0	8.8	16.7	17.3
18-25	31.7	32.0	4.8	5.1	33.5	33.7
26-44	19.7	19.4	3.3	3.8	20.3	20.1
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	23.3	23.3	4.4	4.7	24.0	24.1
White	25.3	25.5	4.0	4.1	26.1	26.4
Black or African American	18.3	17.2	3.0	5.7	18.7	17.5
American Indian or Alaska Native	34.3	37.7	*	*	35.1	38.3
Native Hawaiian or Other Pacific Islander	*	15.3	*	*	*	15.4
Asian	9.0	9.1	*	*	9.1	9.3
Two or More Races	23.7	26.6	*	*	24.7	27.4
Hispanic or Latino	17.8	18.1	2.9	3.7	18.6	18.8
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	10.9	10.6	N/A	N/A
Second	N/A	N/A	1.4	1.9	N/A	N/A
Third	N/A	N/A	0.7	1.1	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 7.77A Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004**

Demographic Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>TOTAL</b>	3,166	3,296	16	13	3,140	3,273
<b>AGE</b>						
15-17	218	235	0	*	216	232
18-25	1,380	1,416	7	7	1,365	1,401
26-44	1,568	1,645	9	5	1,559	1,640
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	2,945	3,079	15	12	2,924	3,061
White	2,568	2,714	11	10	2,553	2,698
Black or African American	237	227	1	*	233	226
American Indian or Alaska Native	27	31	*	*	25	29
Native Hawaiian or Other Pacific Islander	12	7	*	*	11	7
Asian	66	59	*	*	66	59
Two or More Races	36	41	*	*	35	41
Hispanic or Latino	221	217	1	1	217	213
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	12	12	N/A	N/A
Second	N/A	N/A	*	*	N/A	N/A
Third	N/A	N/A	1	1	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 7.77B Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Percentages, Annual Averages Based on 2002-2003 and 2003-2004**

Demographic Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
<b>TOTAL</b>	5.1	5.3	0.7	0.5	5.3	5.6
<b>AGE</b>						
15-17	3.6	3.9	0.1	*	3.7	3.9
18-25	8.8	8.9	0.7	0.7	9.3	9.4
26-44	3.9	4.1	0.6	0.3	4.1	4.3
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	5.6	5.9	0.8	0.6	5.8	6.1
White	6.4 <sup>a</sup>	6.8	0.8	0.6	6.7 <sup>a</sup>	7.1
Black or African American	2.9	2.7	0.5	*	2.9	2.9
American Indian or Alaska Native	6.8	8.2	*	*	7.0	8.3
Native Hawaiian or Other Pacific Islander	6.2	4.1	*	*	6.1	3.9
Asian	2.2	1.9	*	*	2.3	2.0
Two or More Races	5.4	5.8	*	*	5.6	6.0
Hispanic or Latino	2.4	2.3	0.2	0.2	2.5	2.4
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	1.6	1.6	N/A	N/A
Second	N/A	N/A	*	*	N/A	N/A
Third	N/A	N/A	0.1	0.1	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

<sup>a</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2003-2004 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.