

Table 5.59A Received Substance Use Treatment at a Specialty Facility in the Past Year, by Detailed Age Category: Numbers in Thousands, 2003 and 2004

| Age Category | SUBSTANCE FOR WHICH TREATMENT AT A SPECIALTY FACILITY WAS RECEIVED IN PAST YEAR | | | | | | | |
|--------------|---|-------|-----------------|-------|--|------|--|-------|
| | Illicit Drug ¹ | | Alcohol | | Both Illicit Drug and Alcohol ¹ | | Illicit Drug or Alcohol ^{1,2} | |
| | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 |
| TOTAL | 1,103 ^a | 1,427 | 1,298 | 1,535 | 595 | 718 | 1,874 ^a | 2,327 |
| 12 | 3 | 1 | 2 | 3 | 2 | 1 | 6 | 3 |
| 13 | 1 | 4 | 3 | 1 | * | * | 3 | 5 |
| 14 | 13 | 17 | 2 ^a | 11 | 1 | 6 | 13 | 23 |
| 15 | 17 | 29 | 18 | 27 | 7 | 18 | 31 | 41 |
| 16 | 43 | 38 | 32 | 34 | 23 | 26 | 58 | 47 |
| 17 | 37 | 45 | 39 | 50 | 21 | 30 | 56 | 66 |
| 18 | 56 | 81 | 51 | 41 | 33 | 27 | 76 | 97 |
| 19 | 42 | 37 | 21 | 27 | 15 | 20 | 54 | 45 |
| 20 | 39 | 28 | 31 | 37 | 18 | 8 | 53 | 65 |
| 21 | 47 | 57 | 30 ^a | 61 | 16 | 28 | 65 | 91 |
| 22 | 34 | 33 | 49 | 50 | 23 | 18 | 63 | 69 |
| 23 | 46 | 36 | 46 | 46 | 28 | 19 | 64 | 68 |
| 24 | 45 | 50 | 44 | 33 | 29 | 20 | 61 | 68 |
| 25 | 32 | 30 | 33 | 29 | 15 | 14 | 51 | 45 |
| 26-29 | 94 | 171 | 112 | 164 | 50 | 80 | 159 | 269 |
| 30-34 | 78 | 114 | 119 | 136 | 39 | 29 | 163 | 221 |
| 35-39 | 199 | 138 | 205 | 185 | 103 | 90 | 303 | 247 |
| 40-44 | 137 | 251 | 209 | 243 | 88 | 138 | 274 | 379 |
| 45-49 | 55 ^b | 173 | 140 | 189 | 26 ^a | 106 | 169 | 255 |
| 50-54 | 76 | 41 | 63 | 122 | 46 | 27 | 105 | 136 |
| 55-59 | 11 | 10 | 20 | 44 | 11 | 10 | 20 | 44 |
| 60-64 | * | 43 | * | 3 | * | 3 | * | 43 |
| 65 or Older | * | * | 28 | * | * | * | 28 | * |

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 5.59B Received Substance Use Treatment at a Specialty Facility in the Past Year, by Detailed Age Category: Percentages, 2003 and 2004

| Age Category | SUBSTANCE FOR WHICH TREATMENT AT A SPECIALTY FACILITY WAS RECEIVED IN PAST YEAR | | | | | | | |
|--------------|---|------|------------------|------|--|------|--|------|
| | Illicit Drug ¹ | | Alcohol | | Both Illicit Drug and Alcohol ¹ | | Illicit Drug or Alcohol ^{1,2} | |
| | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 |
| TOTAL | 0.5 ^a | 0.6 | 0.5 | 0.6 | 0.3 | 0.3 | 0.8 ^a | 1.0 |
| 12 | 0.1 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.1 |
| 13 | 0.0 | 0.1 | 0.1 | 0.0 | * | * | 0.1 | 0.1 |
| 14 | 0.3 | 0.4 | 0.0 ^a | 0.3 | 0.0 | 0.1 | 0.3 | 0.5 |
| 15 | 0.4 | 0.7 | 0.4 | 0.6 | 0.2 | 0.4 | 0.8 | 1.0 |
| 16 | 1.0 | 0.9 | 0.8 | 0.8 | 0.6 | 0.6 | 1.4 | 1.1 |
| 17 | 0.9 | 1.1 | 1.0 | 1.2 | 0.5 | 0.7 | 1.4 | 1.6 |
| 18 | 1.2 | 1.8 | 1.1 | 0.9 | 0.7 | 0.6 | 1.6 | 2.1 |
| 19 | 1.1 | 0.9 | 0.5 | 0.7 | 0.4 | 0.5 | 1.4 | 1.1 |
| 20 | 1.0 | 0.7 | 0.8 | 0.9 | 0.4 | 0.2 | 1.3 | 1.6 |
| 21 | 1.2 | 1.4 | 0.8 | 1.5 | 0.4 | 0.7 | 1.7 | 2.2 |
| 22 | 0.8 | 0.8 | 1.2 | 1.2 | 0.6 | 0.4 | 1.6 | 1.7 |
| 23 | 1.1 | 0.9 | 1.1 | 1.2 | 0.7 | 0.5 | 1.6 | 1.7 |
| 24 | 1.2 | 1.2 | 1.2 | 0.8 | 0.8 | 0.5 | 1.7 | 1.7 |
| 25 | 0.9 | 0.8 | 0.9 | 0.8 | 0.4 | 0.4 | 1.4 | 1.2 |
| 26-29 | 0.6 | 1.1 | 0.8 | 1.1 | 0.3 | 0.5 | 1.1 | 1.8 |
| 30-34 | 0.4 | 0.6 | 0.6 | 0.7 | 0.2 | 0.1 | 0.8 | 1.1 |
| 35-39 | 0.9 | 0.7 | 1.0 | 0.9 | 0.5 | 0.4 | 1.4 | 1.2 |
| 40-44 | 0.6 | 1.1 | 0.9 | 1.1 | 0.4 | 0.6 | 1.2 | 1.7 |
| 45-49 | 0.3 ^b | 0.8 | 0.7 | 0.9 | 0.1 ^a | 0.5 | 0.8 | 1.2 |
| 50-54 | 0.4 | 0.2 | 0.3 | 0.6 | 0.2 | 0.1 | 0.5 | 0.7 |
| 55-59 | 0.1 | 0.1 | 0.1 | 0.3 | 0.1 | 0.1 | 0.1 | 0.3 |
| 60-64 | * | 0.3 | * | 0.0 | * | 0.0 | * | 0.3 |
| 65 or Older | * | * | 0.1 | * | * | * | 0.1 | * |

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 5.60A Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2003 and 2004

| Demographic Characteristic | SUBSTANCE FOR WHICH TREATMENT AT A SPECIALTY FACILITY WAS RECEIVED IN PAST YEAR | | | | | | | |
|---|---|-------|---------|-------|--|------|--|-------|
| | Illicit Drug ¹ | | Alcohol | | Both Illicit Drug and Alcohol ¹ | | Illicit Drug or Alcohol ^{1,2} | |
| | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 |
| TOTAL | 1,103 ^a | 1,427 | 1,298 | 1,535 | 595 | 718 | 1,874 ^a | 2,327 |
| AGE | | | | | | | | |
| 12-17 | 113 | 134 | 95 | 126 | 55 | 81 | 168 | 185 |
| 18-25 | 340 | 352 | 305 | 323 | 178 | 154 | 486 | 548 |
| 26 or Older | 649 ^a | 941 | 898 | 1,086 | 363 | 483 | 1,221 | 1,594 |
| GENDER | | | | | | | | |
| Male | 732 | 914 | 897 | 1,149 | 415 | 542 | 1,263 | 1,571 |
| Female | 371 | 513 | 401 | 386 | 181 | 175 | 612 | 755 |
| HISPANIC ORIGIN AND RACE | | | | | | | | |
| Not Hispanic or Latino | 1,014 | 1,285 | 1,159 | 1,301 | 554 | 639 | 1,683 | 2,013 |
| White | 757 | 845 | 832 | 928 | 350 | 398 | 1,286 | 1,425 |
| Black or African American | 202 | 334 | 251 | 285 | 162 | 190 | 305 | 443 |
| American Indian or Alaska Native | 21 | 7 | 38 | 26 | 18 | 6 | 43 | 27 |
| Native Hawaiian or Other Pacific Islander | * | * | 5 | * | * | * | 5 | * |
| Asian | 9 | 46 | 7 | 12 | 6 | * | 11 | 58 |
| Two or More Races | 24 | 34 | 26 | 31 | 18 | 26 | 32 | 40 |
| Hispanic or Latino | 89 | 142 | 140 | 234 | 41 | 79 | 191 | 314 |

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 5.60B Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2003 and 2004

| Demographic Characteristic | SUBSTANCE FOR WHICH TREATMENT AT A SPECIALTY FACILITY WAS RECEIVED IN PAST YEAR | | | | | | | |
|---|---|------|---------|------|--|------|--|------|
| | Illicit Drug ¹ | | Alcohol | | Both Illicit Drug and Alcohol ¹ | | Illicit Drug or Alcohol ^{1,2} | |
| | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 |
| TOTAL | 0.5 ^a | 0.6 | 0.5 | 0.6 | 0.3 | 0.3 | 0.8 ^a | 1.0 |
| AGE | | | | | | | | |
| 12-17 | 0.5 | 0.5 | 0.4 | 0.5 | 0.2 | 0.3 | 0.7 | 0.7 |
| 18-25 | 1.1 | 1.1 | 1.0 | 1.0 | 0.6 | 0.5 | 1.5 | 1.7 |
| 26 or Older | 0.4 ^a | 0.5 | 0.5 | 0.6 | 0.2 | 0.3 | 0.7 | 0.9 |
| GENDER | | | | | | | | |
| Male | 0.6 | 0.8 | 0.8 | 1.0 | 0.4 | 0.5 | 1.1 | 1.3 |
| Female | 0.3 | 0.4 | 0.3 | 0.3 | 0.1 | 0.1 | 0.5 | 0.6 |
| HISPANIC ORIGIN AND RACE | | | | | | | | |
| Not Hispanic or Latino | 0.5 | 0.6 | 0.6 | 0.6 | 0.3 | 0.3 | 0.8 | 1.0 |
| White | 0.5 | 0.5 | 0.5 | 0.6 | 0.2 | 0.2 | 0.8 | 0.9 |
| Black or African American | 0.7 | 1.2 | 0.9 | 1.0 | 0.6 | 0.7 | 1.1 | 1.6 |
| American Indian or Alaska Native | 1.7 | 0.5 | 3.0 | 2.0 | 1.4 | 0.5 | 3.4 | 2.1 |
| Native Hawaiian or Other Pacific Islander | * | * | 1.1 | * | * | * | 1.1 | * |
| Asian | 0.1 | 0.5 | 0.1 | 0.1 | 0.1 | * | 0.1 | 0.6 |
| Two or More Races | 0.9 | 1.3 | 1.0 | 1.2 | 0.7 | 1.0 | 1.3 | 1.6 |
| Hispanic or Latino | 0.3 | 0.5 | 0.5 | 0.8 | 0.1 | 0.3 | 0.6 | 1.0 |

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 5.61A Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2003 and 2004

| Geographic Characteristic | SUBSTANCE FOR WHICH TREATMENT AT A SPECIALTY FACILITY WAS RECEIVED IN PAST YEAR | | | | | | | |
|----------------------------|---|-------|---------|-------|--|------|--|-------|
| | Illicit Drug ¹ | | Alcohol | | Both Illicit Drug and Alcohol ¹ | | Illicit Drug or Alcohol ^{1,2} | |
| | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 |
| TOTAL | 1,103 ^a | 1,427 | 1,298 | 1,535 | 595 | 718 | 1,874 ^a | 2,327 |
| GEOGRAPHIC DIVISION | | | | | | | | |
| Northeast | 237 | 345 | 247 | 322 | 140 | 173 | 367 | 507 |
| New England | 54 | 96 | 65 | 82 | 26 | 57 | 100 | 129 |
| Middle Atlantic | 183 | 249 | 182 | 239 | 114 | 117 | 268 | 378 |
| Midwest | 250 | 307 | 372 | 363 | 132 | 142 | 511 | 537 |
| East North Central | 162 | 176 | 243 | 246 | 73 | 82 | 347 | 346 |
| West North Central | 88 | 130 | 129 | 117 | 59 | 60 | 165 | 190 |
| South | 401 | 527 | 385 | 487 | 217 | 243 | 583 | 813 |
| South Atlantic | 231 | 344 | 210 | 325 | 126 | 194 | 327 | 505 |
| East South Central | 82 | 62 | 79 | 64 | 52 | 22 | 111 | 105 |
| West South Central | 87 | 122 | 95 | 98 | 39 | 27 | 145 | 204 |
| West | 214 | 248 | 295 | 363 | 106 | 160 | 413 | 470 |
| Mountain | 83 | 68 | 103 | 74 | 30 | 34 | 163 | 118 |
| Pacific | 131 | 180 | 191 | 289 | 76 | 127 | 250 | 352 |
| COUNTY TYPE | | | | | | | | |
| Large Metro | 629 | 740 | 646 | 787 | 333 | 345 | 959 | 1,219 |
| Small Metro | 291 ^a | 447 | 386 | 488 | 125 ^b | 259 | 587 | 695 |
| 250K - 1 Mil. Pop. | 209 | 284 | 263 | 306 | 86 ^a | 152 | 412 | 453 |
| < 250K Pop. | 81 ^a | 163 | 123 | 183 | 39 ^a | 107 | 175 | 243 |
| Nonmetro | 183 | 240 | 266 | 260 | 137 | 114 | 329 | 412 |
| Urbanized | 99 | 115 | 152 | 116 | 81 | 65 | 182 | 168 |
| Less Urbanized | 70 | 122 | 93 | 121 | 43 | 47 | 124 ^a | 218 |
| Completely Rural | 13 | 4 | 22 | 22 | 13 | 2 | 23 | 26 |

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 5.61B Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2003 and 2004

| Geographic Characteristic | SUBSTANCE FOR WHICH TREATMENT AT A SPECIALTY FACILITY WAS RECEIVED IN PAST YEAR | | | | | | | |
|----------------------------|---|------|---------|------|--|------|--|------|
| | Illicit Drug ¹ | | Alcohol | | Both Illicit Drug and Alcohol ¹ | | Illicit Drug or Alcohol ^{1,2} | |
| | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 |
| TOTAL | 0.5 ^a | 0.6 | 0.5 | 0.6 | 0.3 | 0.3 | 0.8 ^a | 1.0 |
| GEOGRAPHIC DIVISION | | | | | | | | |
| Northeast | 0.5 | 0.8 | 0.5 | 0.7 | 0.3 | 0.4 | 0.8 | 1.1 |
| New England | 0.5 | 0.8 | 0.5 | 0.7 | 0.2 | 0.5 | 0.8 | 1.1 |
| Middle Atlantic | 0.5 | 0.7 | 0.5 | 0.7 | 0.3 | 0.3 | 0.8 | 1.1 |
| Midwest | 0.5 | 0.6 | 0.7 | 0.7 | 0.2 | 0.3 | 1.0 | 1.0 |
| East North Central | 0.4 | 0.5 | 0.6 | 0.6 | 0.2 | 0.2 | 0.9 | 0.9 |
| West North Central | 0.5 | 0.8 | 0.8 | 0.7 | 0.4 | 0.4 | 1.0 | 1.2 |
| South | 0.5 | 0.6 | 0.5 | 0.6 | 0.3 | 0.3 | 0.7 | 0.9 |
| South Atlantic | 0.5 | 0.8 | 0.5 | 0.7 | 0.3 | 0.4 | 0.7 | 1.1 |
| East South Central | 0.6 | 0.4 | 0.6 | 0.4 | 0.4 | 0.2 | 0.8 | 0.7 |
| West South Central | 0.3 | 0.5 | 0.4 | 0.4 | 0.2 | 0.1 | 0.6 | 0.8 |
| West | 0.4 | 0.5 | 0.5 | 0.7 | 0.2 | 0.3 | 0.8 | 0.9 |
| Mountain | 0.5 | 0.4 | 0.7 | 0.5 | 0.2 | 0.2 | 1.0 | 0.7 |
| Pacific | 0.3 | 0.5 | 0.5 | 0.7 | 0.2 | 0.3 | 0.7 | 0.9 |
| COUNTY TYPE | | | | | | | | |
| Large Metro | 0.5 | 0.6 | 0.5 | 0.6 | 0.3 | 0.3 | 0.8 | 0.9 |
| Small Metro | 0.4 ^a | 0.6 | 0.5 | 0.7 | 0.2 ^b | 0.4 | 0.8 | 1.0 |
| 250K - 1 Mil. Pop. | 0.4 | 0.6 | 0.6 | 0.6 | 0.2 ^a | 0.3 | 0.9 | 1.0 |
| < 250K Pop. | 0.3 ^a | 0.7 | 0.5 | 0.7 | 0.2 ^a | 0.4 | 0.7 | 1.0 |
| Nonmetro | 0.5 | 0.6 | 0.7 | 0.6 | 0.3 | 0.3 | 0.8 | 1.0 |
| Urbanized | 0.6 | 0.7 | 0.8 | 0.7 | 0.5 | 0.4 | 1.0 | 1.0 |
| Less Urbanized | 0.4 | 0.6 | 0.5 | 0.6 | 0.2 | 0.2 | 0.7 | 1.1 |
| Completely Rural | 0.3 | 0.1 | 0.5 | 0.6 | 0.3 | 0.0 | 0.6 | 0.7 |

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.