

Table 6.27C Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Psychological Distress Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2003 and 2004

Location of Treatment/Counseling ¹	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2003	2004	2003	2004	2003	2004	2003	2004
Outpatient Mental Health Clinic or Center	129	265	27	51	100	198	71	*
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	199	296	36	62	142	196	*	*
Doctor's Office - Not Part of a Clinic	146	214	21	37	94	126	*	*
Outpatient Medical Clinic	75	178	14	25	53	83	*	*
Partial Day Hospital or Day Treatment Program	42	37	8	18	36	32	20	*
School or University Setting/Clinic/Center ²	19	13	5	12	18	3	*	*
Some Other Place	38	57	9	7	33	42	*	*

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.27D Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Psychological Distress Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2003 and 2004

Location of Treatment/Counseling ¹	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2003	2004	2003	2004	2003	2004	2003	2004
Outpatient Mental Health Clinic or Center	1.86	3.54	2.12	3.96	2.38	4.45	4.61	*
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	2.30	3.30	2.29	3.96	2.74	4.01	*	*
Doctor's Office - Not Part of a Clinic	2.07	2.79	1.80	3.00	2.36	3.39	*	*
Outpatient Medical Clinic	1.23	2.64	1.26	2.12	1.49	2.32	*	*
Partial Day Hospital or Day Treatment Program	0.70	0.57	0.79	1.57	1.01	0.88	1.45	*
School or University Setting/Clinic/Center ²	0.32	0.20	0.53	1.13	0.51	0.09	*	*
Some Other Place	0.64	0.89	0.82	0.67	0.93	1.17	*	*

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.27P Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Psychological Distress Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2004 Versus 2003

Location of Treatment/Counseling ¹	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2003	2004	2003	2004	2003	2004	2003	2004
Outpatient Mental Health Clinic or Center	0.8733		0.4543		0.6150		*	
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	0.4749		0.4760		0.7828		*	
Doctor's Office - Not Part of a Clinic	0.7660		0.6717		0.5531		*	
Outpatient Medical Clinic	0.3015		0.6751		0.8276		*	
Partial Day Hospital or Day Treatment Program	0.2740		0.4480		0.4454		*	
School or University Setting/Clinic/Center ²	0.9015		0.1313		0.2486		*	
Some Other Place	0.9687		0.2616		0.7930		*	

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).