

**Table 1.108A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	22,798	22,745	2,255	2,187	638	611
<b>GEOGRAPHIC DIVISION</b>						
Northeast	4,694	4,802	489	453	138	107
New England	1,306	1,236	115	135	40	24
Middle Atlantic	3,388	3,566	374	318	98	83
Midwest	4,839	5,080	491	452	122	122
East North Central	3,572	3,574	352	332	72	93
West North Central	1,266	1,506	139	120	49	29
South	7,855 <sup>a</sup>	6,989	733	728	237	212
South Atlantic	4,509 <sup>a</sup>	3,931	408	424	154	150
East South Central	1,194	1,049	104	88	19	27
West South Central	2,152	2,010	221	216	64	36
West	5,411	5,873	543	553	141	171
Mountain	1,717	1,764	164	157	29	43
Pacific	3,694	4,109	379	396	111	128
<b>COUNTY TYPE</b>						
Large Metro	12,580	12,763	1,242	1,232	396	358
Small Metro	6,780	6,830	685	659	152	187
250K – 1 Mil. Pop.	4,536	4,521	478	464	114	132
< 250K Pop.	2,245	2,309	207	195	37	55
Nonmetro	3,438	3,153	328	297	89	66
Urbanized	1,731	1,446	140	132	33	29
Less Urbanized	1,441	1,406	170 <sup>a</sup>	116	53 <sup>a</sup>	28
Completely Rural	266	300	18	49	3	9

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 1.108B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	9.5	9.4	0.9	0.9	0.3	0.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	10.3	10.5	1.1	1.0	0.3	0.2
New England	10.9	10.3	1.0	1.1	0.3	0.2
Middle Atlantic	10.1	10.6	1.1	0.9	0.3	0.2
Midwest	8.9	9.3	0.9	0.8	0.2	0.2
East North Central	9.4	9.4	0.9	0.9	0.2	0.2
West North Central	7.8	9.2	0.9	0.7	0.3	0.2
South	9.1 <sup>b</sup>	8.0	0.9	0.8	0.3	0.2
South Atlantic	10.0 <sup>a</sup>	8.5	0.9	0.9	0.3	0.3
East South Central	8.3	7.2	0.7	0.6	0.1	0.2
West South Central	8.1	7.5	0.8	0.8	0.2	0.1
West	9.9	10.6	1.0	1.0	0.3	0.3
Mountain	10.7	10.7	1.0	1.0	0.2	0.3
Pacific	9.5	10.5	1.0	1.0	0.3	0.3
<b>COUNTY TYPE</b>						
Large Metro	9.8	9.7	1.0	0.9	0.3	0.3
Small Metro	9.5	9.5	1.0	0.9	0.2	0.3
250K – 1 Mil. Pop.	9.6	9.5	1.0	1.0	0.2	0.3
< 250K Pop.	9.2	9.7	0.8	0.8	0.2	0.2
Nonmetro	8.5	7.8	0.8	0.7	0.2	0.2
Urbanized	10.4	9.4	0.8	0.9	0.2	0.2
Less Urbanized	7.1	6.9	0.8 <sup>b</sup>	0.6	0.3 <sup>a</sup>	0.1
Completely Rural	7.4	6.3	0.5	1.0	0.1	0.2

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 1.109A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	2,762	2,658	1,168	1,144	312	313
<b>GEOGRAPHIC DIVISION</b>						
Northeast	496	446	227	207	69	45
New England	139 <sup>a</sup>	102	60	50	16	9
Middle Atlantic	357	344	167	158	53	36
Midwest	611	597	260	245	64	81
East North Central	433	412	186	164	45	56
West North Central	178	185	75	82	19	25
South	944	934	373	394	97	99
South Atlantic	492	470	192	207	50	53
East South Central	176	172	70	64	11	24
West South Central	275	293	111	123	36	22
West	711	681	308	297	81	88
Mountain	194	242	87	85	22	28
Pacific	517	439	221	212	60	60
<b>COUNTY TYPE</b>						
Large Metro	1,407	1,392	598	583	155	158
Small Metro	846	793	363	361	97	101
250K – 1 Mil. Pop.	549	531	245	259	68	78
< 250K Pop.	298	262	118	102	29	23
Nonmetro	509	474	208	200	60	54
Urbanized	194	196	71	88	21	26
Less Urbanized	277	221	122 <sup>a</sup>	80	36	19
Completely Rural	38	57	15	32	2	9

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 1.109B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	11.0	10.5	4.6	4.5	1.2	1.2
<b>GEOGRAPHIC DIVISION</b>						
Northeast	10.9	9.8	5.0	4.6	1.5	1.0
New England	11.9 <sup>a</sup>	8.7	5.1	4.2	1.4	0.8
Middle Atlantic	10.6	10.2	5.0	4.7	1.6	1.1
Midwest	10.8	10.5	4.6	4.3	1.1	1.4
East North Central	10.8	10.3	4.6	4.1	1.1	1.4
West North Central	10.7	11.2	4.5	4.9	1.1	1.5
South	10.5	10.3	4.1	4.3	1.1	1.1
South Atlantic	10.7	10.1	4.2	4.5	1.1	1.2
East South Central	12.2	11.9	4.8	4.4	0.8	1.6
West South Central	9.3	9.8	3.7	4.1	1.2	0.7
West	11.8	11.2	5.1	4.9	1.4	1.4
Mountain	11.1	13.8	5.0	4.9	1.2	1.6
Pacific	12.2	10.2	5.2	4.9	1.4	1.4
<b>COUNTY TYPE</b>						
Large Metro	10.5	10.2	4.5	4.3	1.2	1.2
Small Metro	11.1	10.6	4.7	4.8	1.3	1.4
250K – 1 Mil. Pop.	10.8	10.4	4.8	5.1	1.3	1.5
< 250K Pop.	11.7	11.1	4.6	4.3	1.1	1.0
Nonmetro	12.1	11.2	4.9	4.7	1.4	1.3
Urbanized	11.4	12.4	4.2	5.6	1.3	1.6
Less Urbanized	12.9 <sup>a</sup>	10.3	5.7 <sup>a</sup>	3.7	1.7	0.9
Completely Rural	10.1	11.4	3.9	6.3	0.5	1.8

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 1.110A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	4,516	4,333	674	685	144	146
<b>GEOGRAPHIC DIVISION</b>						
Northeast	921	871	185	171	41	34
New England	250	247	45	47	19	14
Middle Atlantic	672	624	140	124	22	19
Midwest	1,051	1,014	138	163	29	26
East North Central	751	694	104	128	21	23
West North Central	299	320	33	35	8	3
South	1,468	1,338	195	171	51	37
South Atlantic	779	702	111	93	31	27
East South Central	262	228	23	24	8	3
West South Central	427	408	61	53	13	7
West	1,075	1,109	157	180	23	50
Mountain	362	372	46	63	5	15
Pacific	713	737	111	117	18	35
<b>COUNTY TYPE</b>						
Large Metro	2,256	2,266	374	386	84	79
Small Metro	1,560	1,435	217	222	36	55
250K – 1 Mil. Pop.	996	941	145	152	28	43
< 250K Pop.	564	494	72	70	8	12
Nonmetro	700	631	83	77	24	12
Urbanized	330	293	34	37	6	3
Less Urbanized	318	283	46	34	16	9
Completely Rural	52	55	3	6	1	*

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 1.110B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	14.0	13.3	2.1	2.1	0.4	0.5
<b>GEOGRAPHIC DIVISION</b>						
Northeast	16.2	15.3	3.3	3.0	0.7	0.6
New England	16.8	16.5	3.0	3.1	1.3	1.0
Middle Atlantic	16.0	14.8	3.3	2.9	0.5	0.5
Midwest	14.1	13.6	1.8	2.2	0.4	0.4
East North Central	14.7	13.5	2.0	2.5	0.4	0.5
West North Central	12.9	13.8	1.4	1.5	0.3	0.1
South	12.7	11.4	1.7	1.5	0.4	0.3
South Atlantic	13.6	12.0	1.9	1.6	0.5	0.5
East South Central	13.4	11.7	1.2	1.3	0.4	0.2
West South Central	11.0	10.4	1.6	1.3	0.3	0.2
West	14.3	14.5	2.1	2.4	0.3	0.7
Mountain	15.8	15.8	2.0	2.7	0.2	0.6
Pacific	13.6	14.0	2.1	2.2	0.3	0.7
<b>COUNTY TYPE</b>						
Large Metro	13.5	13.2	2.2	2.2	0.5	0.5
Small Metro	14.7	13.9	2.0	2.2	0.3	0.5
250K – 1 Mil. Pop.	14.5	13.9	2.1	2.2	0.4	0.6
< 250K Pop.	15.2	13.9	1.9	2.0	0.2	0.3
Nonmetro	14.2	12.7	1.7	1.6	0.5	0.2
Urbanized	14.3	13.9	1.5	1.7	0.3	0.1
Less Urbanized	13.7	12.0	2.0	1.5	0.7	0.4
Completely Rural	16.0	11.2	0.9	1.2	0.4	*

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 1.111A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	15,521	15,754	413	358	181	152
<b>GEOGRAPHIC DIVISION</b>						
Northeast	3,276	3,485	77	75	28	28
New England	917	886	10	39	5	1
Middle Atlantic	2,359	2,598	66	36	23	27
Midwest	3,177	3,469	93	44	28	15
East North Central	2,388	2,468	62	41	6	14
West North Central	789	1,001	31	3	22	1
South	5,443 <sup>a</sup>	4,717	165	164	89	76
South Atlantic	3,238	2,759	105	124	73	70
East South Central	755	649	11	*	*	*
West South Central	1,450	1,309	49	40	15	6
West	3,625	4,083	78	76	36	33
Mountain	1,161	1,150	31	8	3	*
Pacific	2,464	2,933	47	67	33	33
<b>COUNTY TYPE</b>						
Large Metro	8,918	9,105	271	263	157	121
Small Metro	4,374	4,601	105	76	19	31
250K – 1 Mil. Pop.	2,991	3,048	88	53	18	11
< 250K Pop.	1,383	1,553	17	23	*	20
Nonmetro	2,229	2,049	38	20	6	*
Urbanized	1,207	957	35	7	6	*
Less Urbanized	846	903	2	1	*	*
Completely Rural	176	188	*	11	*	0

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 1.111B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	8.5	8.5	0.2	0.2	0.1	0.1
<b>GEOGRAPHIC DIVISION</b>						
Northeast	9.3	9.9	0.2	0.2	0.1	0.1
New England	9.9	9.5	0.1	0.4	0.1	0.0
Middle Atlantic	9.1	10.0	0.3	0.1	0.1	0.1
Midwest	7.7	8.4	0.2	0.1	0.1	0.0
East North Central	8.3	8.5	0.2	0.1	0.0	0.0
West North Central	6.4	8.0	0.3	0.0	0.2	0.0
South	8.3 <sup>a</sup>	7.1	0.3	0.2	0.1	0.1
South Atlantic	9.3	7.7	0.3	0.3	0.2	0.2
East South Central	6.9	5.8	0.1	*	*	*
West South Central	7.4	6.5	0.2	0.2	0.1	0.0
West	8.8	9.8	0.2	0.2	0.1	0.1
Mountain	9.7	9.3	0.3	0.1	0.0	*
Pacific	8.4	10.0	0.2	0.2	0.1	0.1
<b>COUNTY TYPE</b>						
Large Metro	9.1	9.1	0.3	0.3	0.2	0.1
Small Metro	8.2	8.5	0.2	0.1	0.0	0.1
250K – 1 Mil. Pop.	8.5	8.5	0.2	0.1	0.1	0.0
< 250K Pop.	7.6	8.6	0.1	0.1	*	0.1
Nonmetro	7.1	6.5	0.1	0.1	0.0	*
Urbanized	9.5	8.2	0.3	0.1	0.0	*
Less Urbanized	5.3	5.7	0.0	0.0	*	*
Completely Rural	6.1	5.0	*	0.3	*	0.0

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.



**Table 1.112A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	20,036	20,087	1,087	1,043	325	298
<b>GEOGRAPHIC DIVISION</b>						
Northeast	4,198	4,356	262	246	69	62
New England	1,167	1,134	55	86	25	15
Middle Atlantic	3,031	3,222	207	160	45	47
Midwest	4,227	4,484	231	207	57	41
East North Central	3,139	3,162	166	168	27	37
West North Central	1,088	1,321	64	38	30	4
South	6,911 <sup>a</sup>	6,055	360	334	140	113
South Atlantic	4,016	3,461	215	217	104	96
East South Central	1,018	877	34	24	8	3
West South Central	1,877	1,717	110	93	28	13
West	4,700	5,192	235	256	59	83
Mountain	1,523	1,522	77	72	8	15
Pacific	3,177	3,670	158	184	52	68
<b>COUNTY TYPE</b>						
Large Metro	11,173	11,371	645	649	241	200
Small Metro	5,934	6,036	322	298	55	86
250K – 1 Mil. Pop.	3,987	3,989	232	205	46	53
< 250K Pop.	1,947	2,047	89	93	8	32
Nonmetro	2,929	2,679	121	97	30	12
Urbanized	1,537	1,250	70	44	12	3
Less Urbanized	1,164	1,186	48	35	16	9
Completely Rural	228	243	3	17	1	*

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 1.112B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	9.3	9.2	0.5	0.5	0.2	0.1
<b>GEOGRAPHIC DIVISION</b>						
Northeast	10.2	10.6	0.6	0.6	0.2	0.2
New England	10.8	10.5	0.5	0.8	0.2	0.1
Middle Atlantic	10.0	10.6	0.7	0.5	0.1	0.2
Midwest	8.7	9.2	0.5	0.4	0.1	0.1
East North Central	9.3	9.3	0.5	0.5	0.1	0.1
West North Central	7.5	8.9	0.4	0.3	0.2	0.0
South	9.0 <sup>b</sup>	7.7	0.5	0.4	0.2	0.1
South Atlantic	9.9 <sup>a</sup>	8.3	0.5	0.5	0.3	0.2
East South Central	7.9	6.7	0.3	0.2	0.1	0.0
West South Central	8.0	7.2	0.5	0.4	0.1	0.1
West	9.7	10.5	0.5	0.5	0.1	0.2
Mountain	10.7	10.4	0.5	0.5	0.1	0.1
Pacific	9.2	10.6	0.5	0.5	0.1	0.2
<b>COUNTY TYPE</b>						
Large Metro	9.7	9.7	0.6	0.6	0.2	0.2
Small Metro	9.3	9.4	0.5	0.5	0.1	0.1
250K – 1 Mil. Pop.	9.5	9.4	0.6	0.5	0.1	0.1
< 250K Pop.	8.9	9.5	0.4	0.4	0.0	0.1
Nonmetro	8.1	7.4	0.3	0.3	0.1	0.0
Urbanized	10.2	9.1	0.5	0.3	0.1	0.0
Less Urbanized	6.4	6.5	0.3	0.2	0.1	0.1
Completely Rural	7.1	5.7	0.1	0.4	0.0	*

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.