

Table 1.113A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	48,013	48,709	14,643	15,172	6,007	6,405
GEOGRAPHIC DIVISION						
Northeast	8,565	8,870	2,415	2,515	992	1,124
New England	2,323	2,379	657	750	261	328
Middle Atlantic	6,241	6,491	1,758	1,766	731	796
Midwest	10,235	10,265	3,097	3,193	1,195	1,277
East North Central	7,381	7,168	2,312	2,210	904	919
West North Central	2,854	3,097	785	982	291	359
South	17,397	16,755	5,285	5,644	2,377	2,447
South Atlantic	9,221	8,770	2,608	2,801	1,147	1,199
East South Central	2,954	2,853	1,066	1,023	482	450
West South Central	5,223	5,132	1,612	1,819	748	798
West	11,816	12,820	3,845	3,820	1,444	1,556
Mountain	3,731	4,055	1,210	1,242	507	494
Pacific	8,085	8,765	2,636	2,577	937	1,062
COUNTY TYPE						
Large Metro	25,719	26,142	7,441	7,900	2,886 ^a	3,350
Small Metro	14,948	15,019	4,786	4,810	2,148	2,013
250K – 1 Mil. Pop.	10,001	10,185	3,187	3,315	1,409	1,406
< 250K Pop.	4,947	4,834	1,599	1,495	739	607
Nonmetro	7,345	7,548	2,415	2,463	973	1,042
Urbanized	3,223	3,131	1,055	1,067	431	443
Less Urbanized	3,515	3,655	1,181	1,149	463	484
Completely Rural	607	762	179	247	78	115

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.113B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	20.0	20.0	6.1	6.2	2.5	2.6
GEOGRAPHIC DIVISION						
Northeast	18.8	19.4	5.3	5.5	2.2	2.5
New England	19.4	19.9	5.5	6.3	2.2	2.7
Middle Atlantic	18.6	19.3	5.2	5.2	2.2	2.4
Midwest	18.9	18.8	5.7	5.9	2.2	2.3
East North Central	19.5	18.8	6.1	5.8	2.4	2.4
West North Central	17.5	18.9	4.8	6.0	1.8	2.2
South	20.2	19.1	6.1	6.4	2.8	2.8
South Atlantic	20.4	19.0	5.8	6.1	2.5	2.6
East South Central	20.5	19.6	7.4	7.0	3.3	3.1
West South Central	19.7	19.0	6.1	6.7	2.8	3.0
West	21.6	23.1	7.0	6.9	2.6	2.8
Mountain	23.3	24.7	7.6	7.6	3.2	3.0
Pacific	20.9	22.5	6.8	6.6	2.4	2.7
COUNTY TYPE						
Large Metro	20.0	19.9	5.8	6.0	2.2	2.6
Small Metro	20.9	21.0	6.7	6.7	3.0	2.8
250K – 1 Mil. Pop.	21.2	21.3	6.8	6.9	3.0	2.9
< 250K Pop.	20.2	20.2	6.5	6.3	3.0	2.5
Nonmetro	18.1	18.6	6.0	6.1	2.4	2.6
Urbanized	19.3	20.4	6.3	6.9	2.6	2.9
Less Urbanized	17.3	17.9	5.8	5.6	2.3	2.4
Completely Rural	17.0	16.0	5.0	5.2	2.2	2.4

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.114A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	3,394 ^b	3,014	2,225	2,105	914	839
GEOGRAPHIC DIVISION						
Northeast	539 ^a	441	366	319	153	129
New England	155	127	114	95	49	39
Middle Atlantic	384 ^a	314	251	224	104	90
Midwest	752 ^a	671	491	455	195	193
East North Central	552 ^b	465	355	314	143	131
West North Central	200	206	136	141	52	62
South	1,227	1,160	793	835	329	350
South Atlantic	595	533	391	379	151	165
East South Central	220	217	145	166	54	77
West South Central	412	410	257	290	123	108
West	876 ^a	742	575	495	237 ^a	167
Mountain	243	253	150	170	49	54
Pacific	633 ^b	489	425 ^a	325	188 ^b	113
COUNTY TYPE						
Large Metro	1,625	1,498	1,077	1,040	456	401
Small Metro	1,110 ^b	919	727	659	290	269
250K – 1 Mil. Pop.	713	618	467	449	189	189
< 250K Pop.	397 ^a	301	260	210	100	79
Nonmetro	659	597	420	406	168	169
Urbanized	264	234	179	159	71	72
Less Urbanized	342	300	214	203	83	81
Completely Rural	54	63	27	44	14	17

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.114B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	13.5 ^b	11.9	8.8	8.3	3.6	3.3
GEOGRAPHIC DIVISION						
Northeast	11.9 ^a	9.7	8.1	7.0	3.4	2.8
New England	13.2	10.8	9.8	8.1	4.2	3.3
Middle Atlantic	11.4 ^a	9.3	7.5	6.7	3.1	2.7
Midwest	13.2 ^a	11.8	8.6	8.0	3.4	3.4
East North Central	13.8 ^b	11.6	8.9	7.8	3.6	3.3
West North Central	12.0	12.5	8.1	8.6	3.1	3.7
South	13.6	12.8	8.8	9.2	3.7	3.9
South Atlantic	13.0	11.5	8.5	8.2	3.3	3.6
East South Central	15.2	15.0	10.0	11.4	3.8	5.3
West South Central	13.9	13.8	8.6	9.7	4.1	3.6
West	14.6 ^b	12.2	9.6	8.1	4.0 ^b	2.7
Mountain	13.9	14.4	8.6	9.7	2.8	3.1
Pacific	14.9 ^b	11.3	10.0 ^a	7.5	4.4 ^b	2.6
COUNTY TYPE						
Large Metro	12.2 ^a	11.0	8.1	7.6	3.4	2.9
Small Metro	14.5 ^b	12.3	9.5	8.8	3.8	3.6
250K – 1 Mil. Pop.	14.0 ^a	12.1	9.2	8.8	3.7	3.7
< 250K Pop.	15.6 ^a	12.7	10.2	8.9	4.0	3.4
Nonmetro	15.6	14.1	9.9	9.6	4.0	4.0
Urbanized	15.5	14.8	10.6	10.1	4.2	4.6
Less Urbanized	15.9	14.0	10.0	9.5	3.9	3.8
Completely Rural	14.2	12.4	7.2	8.7	3.6	3.3

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.115A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	9,392 ^a	9,858	4,751	4,887	1,959	2,034
GEOGRAPHIC DIVISION						
Northeast	1,652	1,685	889	843	362	361
New England	487	495	270	284	114	128
Middle Atlantic	1,165	1,190	619	559	247	233
Midwest	2,119	2,144	1,060	1,029	418	390
East North Central	1,504	1,510	782	726	319	291
West North Central	614	634	279	303	100	99
South	3,328 ^a	3,570	1,719	1,851	760	826
South Atlantic	1,590	1,657	780	828	346	337
East South Central	611	636	323	324	135	149
West South Central	1,126 ^a	1,276	615	699	278	339
West	2,293	2,459	1,083	1,165	420	457
Mountain	711 ^a	811	351	370	129	149
Pacific	1,582	1,648	732	794	290	308
COUNTY TYPE						
Large Metro	4,619 ^a	5,039	2,248	2,485	893	1,009
Small Metro	3,272	3,273	1,733	1,658	732	720
250K – 1 Mil. Pop.	2,037	2,160	1,106	1,100	455	481
< 250K Pop.	1,235	1,113	627	559	277	239
Nonmetro	1,502	1,547	770	744	334	306
Urbanized	724	704	362	343	150	151
Less Urbanized	666	680	346	312	153	124
Completely Rural	111	162	63	89	*	31

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.115B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	29.2	30.3	14.8	15.0	6.1	6.3
GEOGRAPHIC DIVISION						
Northeast	29.1	29.5	15.6	14.8	6.4	6.3
New England	32.8	33.2	18.1	19.0	7.7	8.5
Middle Atlantic	27.8	28.2	14.7	13.3	5.9	5.5
Midwest	28.5	28.8	14.2	13.8	5.6	5.2
East North Central	29.3	29.5	15.2	14.2	6.2	5.7
West North Central	26.5	27.2	12.0	13.0	4.3	4.3
South	28.8	30.5	14.9	15.8	6.6	7.1
South Atlantic	27.9	28.4	13.7	14.2	6.1	5.8
East South Central	31.3	32.6	16.6	16.6	6.9	7.7
West South Central	29.0	32.5	15.9	17.8	7.2	8.6
West	30.5	32.3	14.4	15.3	5.6	6.0
Mountain	31.1	34.6	15.4	15.8	5.7	6.4
Pacific	30.2	31.2	14.0	15.1	5.5	5.8
COUNTY TYPE						
Large Metro	27.7	29.3	13.5	14.4	5.4	5.9
Small Metro	30.9	31.7	16.3	16.1	6.9	7.0
250K – 1 Mil. Pop.	29.6	31.9	16.1	16.2	6.6	7.1
< 250K Pop.	33.2	31.4	16.9	15.8	7.5	6.7
Nonmetro	30.4	31.2	15.6	15.0	6.8	6.2
Urbanized	31.4	33.4	15.7	16.3	6.5	7.2
Less Urbanized	28.8	28.9	14.9	13.3	6.6	5.2
Completely Rural	34.3	33.0	19.3	18.0	*	6.3

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.116A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	35,227	35,838	7,667	8,180	3,134	3,532
GEOGRAPHIC DIVISION						
Northeast	6,373	6,744	1,160	1,354	477	634
New England	1,681	1,757	272	371	98	161
Middle Atlantic	4,692	4,987	888	982	380	473
Midwest	7,365	7,450	1,546	1,708	581	694
East North Central	5,325	5,193	1,175	1,171	442	496
West North Central	2,040	2,257	371	538	139	198
South	12,842	12,025	2,774	2,958	1,288	1,272
South Atlantic	7,035	6,579	1,437	1,595	649	697
East South Central	2,122	2,000	597	534	293	224
West South Central	3,685	3,445	739	830	346	351
West	8,646	9,619	2,187	2,160	787	932
Mountain	2,777	2,991	708	702	328	290
Pacific	5,869	6,628	1,479	1,458	459	642
COUNTY TYPE						
Large Metro	19,476	19,606	4,115	4,374	1,537	1,941
Small Metro	10,566	10,827	2,326	2,493	1,127	1,024
250K – 1 Mil. Pop.	7,251	7,407	1,614	1,766	765	735
< 250K Pop.	3,315	3,420	713	726	362	289
Nonmetro	5,185	5,405	1,225	1,313	470	567
Urbanized	2,235	2,193	514	565	211	220
Less Urbanized	2,507	2,674	622	634	226	280
Completely Rural	443	538	89	114	33	67

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.116B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	19.2	19.3	4.2	4.4	1.7	1.9
GEOGRAPHIC DIVISION						
Northeast	18.1	19.1	3.3	3.8	1.4	1.8
New England	18.1	18.9	2.9	4.0	1.1	1.7
Middle Atlantic	18.1	19.1	3.4	3.8	1.5	1.8
Midwest	17.9	18.0	3.8	4.1	1.4	1.7
East North Central	18.5	17.9	4.1	4.0	1.5	1.7
West North Central	16.6	18.1	3.0	4.3	1.1	1.6
South	19.6	18.0	4.2	4.4	2.0	1.9
South Atlantic	20.2	18.5	4.1	4.5	1.9	2.0
East South Central	19.3	18.0	5.4	4.8	2.7	2.0
West South Central	18.7	17.2	3.8	4.1	1.8	1.7
West	21.0	23.0	5.3	5.2	1.9	2.2
Mountain	23.2	24.3	5.9	5.7	2.7	2.4
Pacific	20.1	22.5	5.1	5.0	1.6	2.2
COUNTY TYPE						
Large Metro	19.8	19.6	4.2	4.4	1.6	1.9
Small Metro	19.8	20.1	4.4	4.6	2.1	1.9
250K – 1 Mil. Pop.	20.6	20.7	4.6	4.9	2.2	2.1
< 250K Pop.	18.2	19.0	3.9	4.0	2.0	1.6
Nonmetro	16.5	17.2	3.9	4.2	1.5	1.8
Urbanized	17.6	18.8	4.0	4.8	1.7	1.9
Less Urbanized	15.8	16.8	3.9	4.0	1.4	1.8
Completely Rural	15.4	14.3	3.1	3.0	1.2	1.8

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.117A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	44,619	45,695	12,418	13,068	5,093	5,567
GEOGRAPHIC DIVISION						
Northeast	8,026	8,429	2,049	2,196	839	996
New England	2,168	2,252	542	655	212	289
Middle Atlantic	5,857	6,177	1,507	1,541	627	707
Midwest	9,484	9,594	2,606	2,737	1,000	1,084
East North Central	6,829	6,703	1,957	1,896	761	787
West North Central	2,654	2,891	649	841	239	297
South	16,170	15,595	4,492	4,809	2,048	2,098
South Atlantic	8,626	8,237	2,217	2,422	996	1,034
East South Central	2,733	2,636	920	858	428	373
West South Central	4,811	4,722	1,354	1,529	624	690
West	10,940 ^a	12,078	3,271	3,325	1,207	1,389
Mountain	3,488	3,802	1,060	1,073	457	440
Pacific	7,451	8,276	2,211	2,252	750	949
COUNTY TYPE						
Large Metro	24,094	24,644	6,364	6,859	2,430 ^a	2,950
Small Metro	13,838	14,100	4,059	4,151	1,859	1,744
250K – 1 Mil. Pop.	9,288	9,567	2,720	2,866	1,220	1,217
< 250K Pop.	4,550	4,533	1,340	1,285	639	527
Nonmetro	6,687	6,951	1,995	2,057	805	873
Urbanized	2,959	2,897	876	908	360	371
Less Urbanized	3,174	3,355	967	946	379	404
Completely Rural	554	700	152	203	65	98

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.117B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	20.7	21.0	5.8	6.0	2.4	2.6
GEOGRAPHIC DIVISION						
Northeast	19.6	20.5	5.0	5.3	2.0	2.4
New England	20.1	20.9	5.0	6.1	2.0	2.7
Middle Atlantic	19.4	20.4	5.0	5.1	2.1	2.3
Midwest	19.5	19.6	5.4	5.6	2.1	2.2
East North Central	20.1	19.7	5.8	5.6	2.2	2.3
West North Central	18.2	19.6	4.4	5.7	1.6	2.0
South	21.0	19.9	5.8	6.1	2.7	2.7
South Atlantic	21.2	19.9	5.5	5.8	2.5	2.5
East South Central	21.1	20.1	7.1	6.6	3.3	2.9
West South Central	20.4	19.7	5.7	6.4	2.6	2.9
West	22.5	24.5	6.7	6.7	2.5	2.8
Mountain	24.5	25.9	7.4	7.3	3.2	3.0
Pacific	21.6	23.8	6.4	6.5	2.2	2.7
COUNTY TYPE						
Large Metro	21.0	21.0	5.5	5.8	2.1 ^a	2.5
Small Metro	21.6	22.0	6.3	6.5	2.9	2.7
250K – 1 Mil. Pop.	22.1	22.5	6.5	6.7	2.9	2.9
< 250K Pop.	20.8	21.0	6.1	6.0	2.9	2.4
Nonmetro	18.4	19.1	5.5	5.7	2.2	2.4
Urbanized	19.7	21.0	5.8	6.6	2.4	2.7
Less Urbanized	17.5	18.4	5.3	5.2	2.1	2.2
Completely Rural	17.4	16.4	4.8	4.8	2.0	2.3

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.