

Table 1.19A Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	110,057	112,085	34,807	35,041	19,071	19,720
12	443	470	265	292	112	102
13	810	715	509	483	204	209
14	1,104	981	777 ^a	669	395 ^b	289
15	1,489 ^a	1,323	1,054	954	547	482
16	1,750	1,644	1,277	1,267	638	682
17	1,970	1,888	1,419	1,376	778	747
18	2,456	2,372	1,782	1,667	974	934
19	2,229	2,446	1,520	1,665	897	970
20	2,354	2,455	1,520	1,584	851	990
21	2,541	2,400	1,492	1,440	885	832
22	2,529	2,498	1,412	1,379	826	813
23	2,335	2,468	1,111 ^a	1,303	605 ^b	756
24	2,421	2,260	1,117	1,053	655	616
25	2,178	2,323	959	1,026	546	631
26-29	9,210	9,309	3,603	3,541	2,028	2,000
30-34	10,693	10,456	3,080	3,397	1,837	1,851
35-39	12,434	11,605	2,953	2,703	1,500	1,521
40-44	14,723	14,478	3,272	3,042	1,705	1,695
45-49	13,295	13,919	2,543	2,588	1,465	1,458
50-54	10,917	10,961	1,743	1,632	935	1,028
55-59	6,264 ^a	7,566	830	959	427	591
60-64	3,032	3,727	248	418	140	239
65 or Older	2,879 ^a	3,819	321	602	122	282

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

^aDifference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.19B Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	45.8	46.1	14.5	14.4	7.9	8.1
12	11.2	11.7	6.7	7.3	2.8	2.5
13	18.4	16.9	11.6	11.4	4.6	4.9
14	25.2 ^a	22.6	17.8 ^a	15.4	9.0 ^b	6.7
15	34.7 ^b	30.3	24.6 ^a	21.9	12.7	11.1
16	42.5 ^b	38.1	31.0	29.4	15.5	15.8
17	48.4	45.9	34.9	33.5	19.1	18.2
18	53.4	51.9	38.8	36.5	21.2	20.5
19	56.6	57.0	38.6	38.8	22.8	22.6
20	59.0	59.8	38.1	38.6	21.3	24.1
21	62.3	59.7	36.6	35.8	21.7	20.7
22	62.9	61.6	35.1	34.0	20.5	20.0
23	59.5 ^a	63.5	28.3 ^b	33.5	15.4 ^b	19.5
24	59.8	60.8	27.6	28.3	16.2	16.6
25	60.5	60.4	26.7	26.7	15.2	16.4
26-29	60.0	59.9	23.5	22.8	13.2	12.9
30-34	54.5	54.1	15.7	17.6	9.4	9.6
35-39	59.4	58.2	14.1	13.6	7.2	7.6
40-44	64.9 ^a	61.9	14.4	13.0	7.5	7.2
45-49	61.8	63.5	11.8	11.8	6.8	6.6
50-54	56.3	55.6	9.0	8.3	4.8	5.2
55-59	38.2 ^a	44.1	5.1	5.6	2.6	3.4
60-64	24.2	28.2	2.0	3.2	1.1	1.8
65 or Older	8.3 ^a	10.9	0.9	1.7	0.4	0.8

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.20A Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	96,772	97,545	25,451	25,375	14,576	14,626
12	64	53	42	40	15	10
13	241	226	188	170	81	65
14	560 ^a	444	453 ^b	349	217	165
15	964	886	740	701	412	345
16	1,358	1,255	1,032	984	525	520
17	1,602	1,538	1,201	1,138	659	625
18	2,047	2,022	1,433	1,409	790	797
19	1,975	2,129	1,294	1,372	748	805
20	2,116	2,128	1,315	1,328	744	854
21	2,303	2,135	1,207	1,167	736	662
22	2,270	2,226	1,156	1,142	692	691
23	2,065	2,225	881 ^a	1,042	494	594
24	2,216	2,072	918	839	544	487
25	1,990	2,094	758	813	448	499
26-29	8,093	8,218	2,764	2,724	1,651	1,539
30-34	9,573	9,154	2,114	2,449	1,265	1,463
35-39	11,274	10,388	2,058	1,687	1,083	972
40-44	13,650	13,259	2,152	2,008	1,297	1,103
45-49	12,325	12,943	1,769	1,760	1,093	1,043
50-54	10,194	9,996	1,299	1,109	737	736
55-59	5,456 ^a	6,607	515	653	273	425
60-64	2,468	3,026	68	233	26	113
65 or Older	1,969	2,521	94	257	45	114

*Low precision; no estimate reported.

^aDifference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.20B Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	40.2	40.1	10.6	10.4	6.1	6.0
12	1.6	1.3	1.1	1.0	0.4	0.2
13	5.5	5.4	4.3	4.0	1.8	1.5
14	12.8 ^b	10.2	10.4 ^b	8.0	5.0	3.8
15	22.4	20.3	17.2	16.1	9.6	7.9
16	33.0 ^b	29.1	25.1	22.8	12.8	12.0
17	39.4	37.4	29.5	27.7	16.2	15.2
18	44.5	44.3	31.2	30.8	17.2	17.4
19	50.2	49.6	32.9	31.9	19.0	18.7
20	53.0	51.8	33.0	32.3	18.6	20.8
21	56.5	53.2	29.6	29.0	18.0	16.5
22	56.5	54.9	28.8	28.2	17.2	17.0
23	52.6 ^b	57.3	22.4 ^b	26.8	12.6 ^a	15.3
24	54.8	55.8	22.7	22.6	13.4	13.1
25	55.3	54.5	21.1	21.2	12.4	13.0
26-29	52.7	52.9	18.0	17.5	10.8	9.9
30-34	48.8	47.4	10.8	12.7	6.4	7.6
35-39	53.8	52.1	9.8	8.5	5.2	4.9
40-44	60.2 ^a	56.7	9.5	8.6	5.7	4.7
45-49	57.3	59.0	8.2	8.0	5.1	4.8
50-54	52.6	50.7	6.7	5.6	3.8	3.7
55-59	33.3 ^a	38.5	3.1	3.8	1.7	2.5
60-64	19.7	22.9	0.5	1.8	0.2	0.9
65 or Older	5.7	7.2	0.3	0.7	0.1	0.3

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.21A Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	34,153	33,673	5,658	5,523	2,021 ^a	2,397
12	4	4	3	4	*	0
13	17	11	13	10	5	2
14	54	39	24	29	7	16
15	103	75	68	60	26	22
16	177	177	130	130	26	39
17	256	276	173	191	63	63
18	409	411	258	253	80	91
19	465	510	260	281	87	98
20	548	576	317	303	92	108
21	705	653	364	293	125	111
22	763	691	316	335	97	132
23	673	707	227	284	81	99
24	707	667	224	240	74	90
25	624	690	167 ^b	259	55 ^a	103
26-29	2,519	2,530	780	602	230	289
30-34	3,178	2,996	612	483	252	158
35-39	4,989 ^b	4,068	486	415	205	210
40-44	6,709	6,404	659	506	235	289
45-49	5,487 ^a	6,212	319	457	157	256
50-54	3,809	3,398	163	225	56	140
55-59	1,268 ^a	1,794	43	101	34	37
60-64	317	495	29	48	22	*
65 or Older	374	290	23	15	12	*

*Low precision; no estimate reported.

^aDifference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.21B Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	14.2	13.8	2.4	2.3	0.8	1.0
12	0.1	0.1	0.1	0.1	*	0.0
13	0.4	0.3	0.3	0.2	0.1	0.0
14	1.2	0.9	0.6	0.7	0.2	0.4
15	2.4	1.7	1.6	1.4	0.6	0.5
16	4.3	4.1	3.1	3.0	0.6	0.9
17	6.3	6.7	4.3	4.6	1.5	1.5
18	8.9	9.0	5.6	5.5	1.7	2.0
19	11.8	11.9	6.6	6.5	2.2	2.3
20	13.7	14.0	8.0	7.4	2.3	2.6
21	17.3	16.3	8.9	7.3	3.1	2.8
22	19.0	17.0	7.9	8.3	2.4	3.3
23	17.1	18.2	5.8	7.3	2.1	2.5
24	17.5	18.0	5.5	6.4	1.8	2.4
25	17.4	18.0	4.6 ^a	6.7	1.5 ^a	2.7
26-29	16.4	16.3	5.1	3.9	1.5	1.9
30-34	16.2	15.5	3.1	2.5	1.3	0.8
35-39	23.8 ^a	20.4	2.3	2.1	1.0	1.1
40-44	29.6	27.4	2.9	2.2	1.0	1.2
45-49	25.5 ^a	28.3	1.5	2.1	0.7	1.2
50-54	19.7	17.2	0.8	1.1	0.3	0.7
55-59	7.7	10.5	0.3	0.6	0.2	0.2
60-64	2.5	3.7	0.2	0.4	0.2	*
65 or Older	1.1	0.8	0.1	0.0	0.0	*

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.22A Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	7,840	7,928	1,304	1,381	467 ^a	682
12	1	*	1	*	*	*
13	3	3	3	2	3	1
14	14	8	8	6	4	3
15	24 ^a	8	11	5	2	1
16	34	31	21	17	2	8
17	48	57	19	24	6	11
18	86	106	45	46	6	19
19	117	124	41	67	12	19
20	108	133	20	33	11	12
21	159	114	58 ^a	27	23 ^a	7
22	189 ^a	133	29	45	5 ^a	23
23	156	197	31	49	11	10
24	163	170	25	26	6	8
25	157	168	13 ^a	33	6	6
26-29	681	681	159	139	45	78
30-34	918	883	129	111	66	50
35-39	1,389	1,138	192	192	56	91
40-44	1,514	1,627	227	229	90	134
45-49	1,163	1,332	111	174	65	101
50-54	544	670	117	60	34	32
55-59	219	267	14	35	14	22
60-64	44	62	7	*	*	*
65 or Older	108	15	23	15	*	*

*Low precision; no estimate reported.

^aDifference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.22B Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	3.3	3.3	0.5	0.6	0.2	0.3
12	0.0	*	0.0	*	*	*
13	0.1	0.1	0.1	0.0	0.1	0.0
14	0.3	0.2	0.2	0.1	0.1	0.1
15	0.6 ^a	0.2	0.3	0.1	0.0	0.0
16	0.8	0.7	0.5	0.4	0.0	0.2
17	1.2	1.4	0.5	0.6	0.1	0.3
18	1.9	2.3	1.0	1.0	0.1	0.4
19	3.0	2.9	1.0	1.6	0.3	0.4
20	2.7	3.2	0.5	0.8	0.3	0.3
21	3.9	2.8	1.4 ^a	0.7	0.6 ^a	0.2
22	4.7 ^a	3.3	0.7	1.1	0.1 ^a	0.6
23	4.0	5.1	0.8	1.3	0.3	0.3
24	4.0	4.6	0.6	0.7	0.1	0.2
25	4.4	4.4	0.4 ^a	0.9	0.2	0.1
26-29	4.4	4.4	1.0	0.9	0.3	0.5
30-34	4.7	4.6	0.7	0.6	0.3	0.3
35-39	6.6	5.7	0.9	1.0	0.3	0.5
40-44	6.7	7.0	1.0	1.0	0.4	0.6
45-49	5.4	6.1	0.5	0.8	0.3	0.5
50-54	2.8	3.4	0.6	0.3	0.2	0.2
55-59	1.3	1.6	0.1	0.2	0.1	0.1
60-64	0.4	0.5	0.1	*	*	*
65 or Older	0.3	0.0	0.1	0.0	*	*

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.23A Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	34,333	33,728	3,878	3,809	929	1,088
12	24	22	14	8	8	3
13	78	67	59	41	18	19
14	127 ^a	79	86	60	22	14
15	205	171	142	110	26	30
16	284	274	204	195	59	63
17	441	384	248	251	77	77
18	579	497	320	308	81	83
19	622 ^a	751	292	355	71	78
20	753	822	299	320	75	103
21	964	903	308	290	74	91
22	1,054 ^a	900	251	251	53	41
23	970	1,020	187	222	53	53
24	1,004	959	161	164	43	23
25	910	968	126	141	26	31
26-29	3,695	3,647	491	425	122	146
30-34	3,601	3,592	296	253	46	37
35-39	3,936	3,643	178	110	42	56
40-44	4,799	4,289	130	113	26	15
45-49	4,598	4,625	40	34	6	15
50-54	3,990	3,475	5 ^a	112	*	76
55-59	1,180 ^b	1,885	18	*	*	*
60-64	262 ^a	592	*	46	*	35
65 or Older	257	162	24	*	*	*

*Low precision; no estimate reported.

^aDifference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.23B Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	14.3	13.9	1.6	1.6	0.4	0.4
12	0.6	0.5	0.3	0.2	0.2	0.1
13	1.8	1.6	1.3	1.0	0.4	0.4
14	2.9 ^a	1.8	2.0	1.4	0.5	0.3
15	4.8	3.9	3.3	2.5	0.6	0.7
16	6.9	6.3	4.9	4.5	1.4	1.5
17	10.9	9.3	6.1	6.1	1.9	1.9
18	12.6	10.9	7.0	6.7	1.8	1.8
19	15.8	17.5	7.4	8.3	1.8	1.8
20	18.9	20.0	7.5	7.8	1.9	2.5
21	23.6	22.5	7.5	7.2	1.8	2.3
22	26.2 ^b	22.2	6.2	6.2	1.3	1.0
23	24.7	26.3	4.8	5.7	1.4	1.4
24	24.8	25.8	4.0	4.4	1.1	0.6
25	25.3	25.2	3.5	3.7	0.7	0.8
26-29	24.1	23.5	3.2	2.7	0.8	0.9
30-34	18.3	18.6	1.5	1.3	0.2	0.2
35-39	18.8	18.3	0.8	0.6	0.2	0.3
40-44	21.2 ^a	18.3	0.6	0.5	0.1	0.1
45-49	21.4	21.1	0.2	0.2	0.0	0.1
50-54	20.6	17.6	0.0 ^a	0.6	*	0.4
55-59	7.2 ^b	11.0	0.1	*	*	*
60-64	2.1 ^a	4.5	*	0.3	*	0.3
65 or Older	0.7	0.5	0.1	*	*	*

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.24A Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	22,798	22,745	2,255	2,187	638	611
12	291	305	145	164	53	50
13	461	404	195	205	47	71
14	531	508	246	224	64	51
15	554	496	241	221	73	56
16	481	470	172	184	38	52
17	444	475	170	145	37	33
18	503	446	161	151	45	25
19	476	539	116	140	20	35
20	513	515	99	104	15	21
21	616	516	104	88	10	16
22	628	585	71	84	19	20
23	601	619	48	55	20	8
24	625	557	48	35	9	9
25	553	556	28	28	7	13
26-29	2,146	2,328	67	88	16	11
30-34	2,432	2,593	107	70	54	41
35-39	2,717	2,436	65	51	38	39
40-44	2,974	2,883	81	75	20	15
45-49	2,377	2,573	28	30	14	2
50-54	1,681	1,544	25	*	*	*
55-59	680	762	*	*	*	*
60-64	241	382	40	*	40	*
65 or Older	273	253	*	43	*	43

*Low precision; no estimate reported.

^aDifference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.24B Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	9.5	9.4	0.9	0.9	0.3	0.3
12	7.3	7.6	3.6	4.1	1.3	1.2
13	10.5	9.6	4.4	4.9	1.1	1.7
14	12.1	11.7	5.6	5.2	1.5	1.2
15	12.9	11.4	5.6	5.1	1.7	1.3
16	11.7	10.9	4.2	4.3	0.9	1.2
17	10.9	11.6	4.2	3.5	0.9	0.8
18	10.9	9.8	3.5	3.3	1.0	0.6
19	12.1	12.6	2.9	3.3	0.5	0.8
20	12.9	12.5	2.5	2.5	0.4	0.5
21	15.1	12.9	2.5	2.2	0.2	0.4
22	15.6	14.4	1.8	2.1	0.5	0.5
23	15.3	15.9	1.2	1.4	0.5	0.2
24	15.4	15.0	1.2	0.9	0.2	0.2
25	15.4	14.5	0.8	0.7	0.2	0.3
26-29	14.0	15.0	0.4	0.6	0.1	0.1
30-34	12.4	13.4	0.5	0.4	0.3	0.2
35-39	13.0	12.2	0.3	0.3	0.2	0.2
40-44	13.1	12.3	0.4	0.3	0.1	0.1
45-49	11.1	11.7	0.1	0.1	0.1	0.0
50-54	8.7	7.8	0.1	*	*	*
55-59	4.1	4.4	*	*	*	*
60-64	1.9	2.9	0.3	*	0.3	*
65 or Older	0.8	0.7	*	0.1	*	0.1

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.25A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	48,013	48,709	14,643	15,172	6,007	6,405
12	189	173	115	119	46	56
13	366 ^a	292	251	193	98	83
14	461 ^b	348	321	260	169 ^b	99
15	672 ^b	543	438	384	186	148
16	806	741	526	528	179	207
17	899	918	574	620	236	246
18	1,217	1,068	784	670	319	258
19	1,098	1,222	672	780	300	338
20	1,152 ^a	1,357	622	714	248	273
21	1,306	1,289	689	632	309	269
22	1,339	1,272	629	643	249	254
23	1,179	1,334	517	557	198	255
24	1,133	1,152	467	452	188	197
25	969 ^b	1,164	371	439	149	189
26-29	4,096	4,297	1,546	1,462	627	627
30-34	4,265	4,359	1,417	1,467	629	585
35-39	5,145	4,723	1,186	1,291	498	539
40-44	6,300	5,816	1,269	1,383	480	571
45-49	5,967	6,159	1,106	1,010	447	471
50-54	5,013	4,525	461	662	183	355
55-59	2,261 ^a	2,982	337	397	154	167
60-64	979	1,426	141	206	51	91
65 or Older	1,202	1,549	205	301	65	125

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

^aDifference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.25B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	20.0	20.0	6.1	6.2	2.5	2.6
12	4.8	4.3	2.9	3.0	1.2	1.4
13	8.3	6.9	5.7	4.6	2.2	2.0
14	10.5 ^b	8.0	7.3	6.0	3.9 ^b	2.3
15	15.7 ^b	12.5	10.2	8.8	4.3	3.4
16	19.6 ^a	17.2	12.8	12.3	4.3	4.8
17	22.1	22.3	14.1	15.1	5.8	6.0
18	26.5 ^a	23.4	17.1 ^a	14.7	6.9	5.7
19	27.9	28.5	17.1	18.2	7.6	7.9
20	28.9 ^a	33.0	15.6	17.4	6.2	6.7
21	32.0	32.1	16.9	15.7	7.6	6.7
22	33.3	31.4	15.7	15.8	6.2	6.3
23	30.0 ^b	34.3	13.2	14.3	5.0	6.6
24	28.0	31.0	11.5	12.2	4.6	5.3
25	26.9 ^a	30.3	10.3	11.4	4.1	4.9
26-29	26.7	27.7	10.1	9.4	4.1	4.0
30-34	21.7	22.6	7.2	7.6	3.2	3.0
35-39	24.6	23.7	5.7	6.5	2.4	2.7
40-44	27.8 ^a	24.9	5.6	5.9	2.1	2.4
45-49	27.7	28.1	5.1	4.6	2.1	2.1
50-54	25.9	23.0	2.4	3.4	0.9	1.8
55-59	13.8 ^a	17.4	2.1	2.3	0.9	1.0
60-64	7.8	10.8	1.1	1.6	0.4	0.7
65 or Older	3.5	4.4	0.6	0.9	0.2	0.4

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.26A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	31,768	32,692	11,256	11,815	4,404	4,658
12	163	148	98	107	39	54
13	291	245	203	159	84	66
14	398 ^b	298	282	221	153 ^b	80
15	547 ^b	432	349	309	153	125
16	698	616	442	436	142	165
17	783	777	490	508	179	196
18	1,083 ^a	927	657	579	244	217
19	925 ^a	1,075	546	646	219	247
20	990 ^a	1,149	499	569	194	194
21	1,114	1,083	564	527	236	200
22	1,094	1,069	490	529	196	181
23	972	1,087	412	464	154	187
24	909	951	396	365	154	155
25	725 ^b	937	269 ^a	351	107	153
26-29	3,095	3,335	1,173	1,138	401	472
30-34	3,134	3,287	1,073	1,113	440	451
35-39	3,135	3,180	850	993	322	363
40-44	3,663	3,422	932	956	351	355
45-49	3,318	3,389	771	609	331	280
50-54	2,455	2,529	314	550	149	306
55-59	1,160	1,509	277	326	64	87
60-64	433	662	46	144	26	51
65 or Older	683	585	122	214	65	72

*Low precision; no estimate reported.

^aDifference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.26B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	13.2	13.4	4.7	4.9	1.8	1.9
12	4.1	3.7	2.5	2.7	1.0	1.3
13	6.6	5.8	4.6	3.8	1.9	1.6
14	9.1 ^b	6.9	6.5	5.1	3.5 ^b	1.8
15	12.7 ^b	9.9	8.1	7.1	3.6	2.9
16	17.0 ^a	14.3	10.7	10.1	3.5	3.8
17	19.2	18.9	12.1	12.4	4.4	4.8
18	23.6 ^a	20.3	14.3	12.7	5.3	4.7
19	23.5	25.0	13.9	15.0	5.6	5.8
20	24.8 ^a	28.0	12.5	13.9	4.9	4.7
21	27.3	27.0	13.8	13.1	5.8	5.0
22	27.2	26.4	12.2	13.1	4.9	4.5
23	24.7 ^a	28.0	10.5	11.9	3.9	4.8
24	22.5 ^a	25.6	9.8	9.8	3.8	4.2
25	20.1 ^b	24.4	7.5	9.1	3.0	4.0
26-29	20.2	21.5	7.6	7.3	2.6	3.0
30-34	16.0	17.0	5.5	5.8	2.2	2.3
35-39	15.0	15.9	4.1	5.0	1.5	1.8
40-44	16.2	14.6	4.1	4.1	1.5	1.5
45-49	15.4	15.5	3.6	2.8	1.5	1.3
50-54	12.7	12.8	1.6	2.8	0.8	1.6
55-59	7.1	8.8	1.7	1.9	0.4	0.5
60-64	3.5	5.0	0.4	1.1	0.2	0.4
65 or Older	2.0	1.7	0.4	0.6	0.2	0.2

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.27A Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	70,657	71,822	19,658	20,109	8,247	8,963
12	411	443	236	267	101	96
13	711 ^a	604	417	374	146	159
14	841	741	514	437	239 ^b	155
15	1,058 ^b	882	641	555	261	217
16	1,106	1,037	689	703	250	298
17	1,229	1,237	772	793	329	327
18	1,571	1,432	1,015	898	412	360
19	1,405 ^a	1,595	840	971	392	428
20	1,486	1,659	818	914	335	402
21	1,721	1,620	926	883	391	395
22	1,774	1,668	824	833	353	369
23	1,599	1,724	666	760	260 ^a	353
24	1,628	1,529	613	615	277	265
25	1,445	1,572	499	594	198 ^a	278
26-29	6,055	6,274	2,038	1,891	835	886
30-34	6,856	6,858	1,912	1,851	870	711
35-39	8,257	7,592	1,675	1,632	694	772
40-44	9,771	9,394	1,815	1,858	726	836
45-49	8,601	9,286	1,329	1,331	565	667
50-54	7,032	6,586	610	817	233	468
55-59	3,238 ^a	4,202	362	499	188	204
60-64	1,244 ^a	1,878	209	275	113	149
65 or Older	1,618	2,008	239	360	77	168

*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

^aDifference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.27B Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	29.4	29.5	8.2	8.3	3.4	3.7
12	10.3	11.1	5.9	6.7	2.6	2.4
13	16.2	14.3	9.5	8.9	3.3	3.8
14	19.2	17.1	11.8	10.1	5.5 ^b	3.6
15	24.6 ^b	20.2	14.9 ^a	12.7	6.1	5.0
16	26.9 ^a	24.0	16.7	16.3	6.1	6.9
17	30.2	30.1	19.0	19.3	8.1	8.0
18	34.2	31.4	22.1	19.7	9.0	7.9
19	35.7	37.2	21.3	22.6	10.0	10.0
20	37.2	40.4	20.5	22.2	8.4	9.8
21	42.2	40.3	22.7	22.0	9.6	9.8
22	44.1	41.1	20.5	20.5	8.8	9.1
23	40.7 ^a	44.4	17.0	19.6	6.6 ^b	9.1
24	40.2	41.2	15.2	16.5	6.8	7.1
25	40.2	40.9	13.9	15.4	5.5	7.2
26-29	39.5	40.4	13.3	12.2	5.4	5.7
30-34	34.9	35.5	9.7	9.6	4.4	3.7
35-39	39.4	38.1	8.0	8.2	3.3	3.9
40-44	43.1	40.2	8.0	7.9	3.2	3.6
45-49	40.0	42.3	6.2	6.1	2.6	3.0
50-54	36.3	33.4	3.1	4.1	1.2	2.4
55-59	19.7 ^a	24.5	2.2	2.9	1.1	1.2
60-64	9.9 ^a	14.2	1.7	2.1	0.9	1.1
65 or Older	4.7	5.7	0.7	1.0	0.2	0.5

*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.