

Table 1.98A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	7,840	7,928	1,304	1,381	467 ^a	682
GEOGRAPHIC DIVISION						
Northeast	1,316	1,355	195	274	88	177
New England	352	421	48	60	14	37
Middle Atlantic	964	934	147	214	74	139
Midwest	1,688	1,511	425	302	155	146
East North Central	1,196	958	329 ^a	199	129	68
West North Central	492	553	96	103	27	79
South	2,939	2,821	490	531	206	234
South Atlantic	1,723	1,556	318	295	144	118
East South Central	510	460	105	114	32	46
West South Central	707	805	67	123	30	70
West	1,896	2,241	194	274	17 ^b	125
Mountain	608	636	50 ^a	151	11	76
Pacific	1,288	1,605	144	123	6 ^a	49
COUNTY TYPE						
Large Metro	4,153	4,454	657	765	245	392
Small Metro	2,438	2,293	425	454	147	245
250K – 1 Mil. Pop.	1,649	1,453	268	317	86 ^a	188
< 250K Pop.	789	840	157	137	61	58
Nonmetro	1,249	1,181	222	162	75	44
Urbanized	655 ^a	436	99	57	36	15
Less Urbanized	535	629	109	98	35	28
Completely Rural	59	115	13	6	4	1

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.98B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	3.3	3.3	0.5	0.6	0.2	0.3
GEOGRAPHIC DIVISION						
Northeast	2.9	3.0	0.4	0.6	0.2	0.4
New England	2.9	3.5	0.4	0.5	0.1	0.3
Middle Atlantic	2.9	2.8	0.4	0.6	0.2	0.4
Midwest	3.1	2.8	0.8	0.6	0.3	0.3
East North Central	3.2	2.5	0.9 ^a	0.5	0.3	0.2
West North Central	3.0	3.4	0.6	0.6	0.2	0.5
South	3.4	3.2	0.6	0.6	0.2	0.3
South Atlantic	3.8	3.4	0.7	0.6	0.3	0.3
East South Central	3.5	3.2	0.7	0.8	0.2	0.3
West South Central	2.7	3.0	0.3	0.5	0.1	0.3
West	3.5	4.0	0.4	0.5	0.0 ^b	0.2
Mountain	3.8	3.9	0.3 ^a	0.9	0.1	0.5
Pacific	3.3	4.1	0.4	0.3	0.0 ^a	0.1
COUNTY TYPE						
Large Metro	3.2	3.4	0.5	0.6	0.2	0.3
Small Metro	3.4	3.2	0.6	0.6	0.2	0.3
250K – 1 Mil. Pop.	3.5	3.0	0.6	0.7	0.2 ^a	0.4
< 250K Pop.	3.2	3.5	0.6	0.6	0.2	0.2
Nonmetro	3.1	2.9	0.5	0.4	0.2	0.1
Urbanized	3.9	2.8	0.6	0.4	0.2	0.1
Less Urbanized	2.6	3.1	0.5	0.5	0.2	0.1
Completely Rural	1.7	2.4	0.4	0.1	0.1	0.0

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.99A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	125	108	63	54	16	25
GEOGRAPHIC DIVISION						
Northeast	22	17	13	8	4	2
New England	10	5	8	2	0	2
Middle Atlantic	13	11	6	6	3	1
Midwest	23	20	15	9	3	2
East North Central	19	9	13	6	2	1
West North Central	4	11	2	4	1	1
South	40	41	21	17	6	7
South Atlantic	14	17	4	8	2	4
East South Central	9	14	2	5	*	1
West South Central	17	10	14	5	4	2
West	40	30	14	19	4	13
Mountain	17	8	7	3	2	2
Pacific	23	22	6	16	2	11
COUNTY TYPE						
Large Metro	48	46	23	27	4	13
Small Metro	54	37	26	15	7	8
250K – 1 Mil. Pop.	36	26	17	9	4	3
< 250K Pop.	18	11	9	6	3	5
Nonmetro	23	25	14	11	5	5
Urbanized	9	12	4	6	*	4
Less Urbanized	13	11	9	2	5	1
Completely Rural	2	2	1	2	*	*

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.99B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.5	0.4	0.3	0.2	0.1	0.1
GEOGRAPHIC DIVISION						
Northeast	0.5	0.4	0.3	0.2	0.1	0.1
New England	0.8	0.4	0.6	0.2	0.0	0.1
Middle Atlantic	0.4	0.3	0.2	0.2	0.1	0.0
Midwest	0.4	0.3	0.3	0.2	0.0	0.0
East North Central	0.5	0.2	0.3	0.1	0.1	0.0
West North Central	0.2	0.6	0.1	0.2	0.0	0.0
South	0.4	0.5	0.2	0.2	0.1	0.1
South Atlantic	0.3	0.4	0.1	0.2	0.1	0.1
East South Central	0.6	1.0	0.2	0.3	*	0.1
West South Central	0.6	0.3	0.5	0.2	0.1	0.1
West	0.7	0.5	0.2	0.3	0.1	0.2
Mountain	1.0	0.5	0.4	0.2	0.1	0.1
Pacific	0.5	0.5	0.2	0.4	0.0	0.3
COUNTY TYPE						
Large Metro	0.4	0.3	0.2	0.2	0.0	0.1
Small Metro	0.7	0.5	0.3	0.2	0.1	0.1
250K – 1 Mil. Pop.	0.7	0.5	0.3	0.2	0.1	0.1
< 250K Pop.	0.7	0.5	0.4	0.3	0.1	0.2
Nonmetro	0.6	0.6	0.3	0.3	0.1	0.1
Urbanized	0.5	0.7	0.3	0.4	*	0.2
Less Urbanized	0.6	0.5	0.4	0.1	0.2	0.1
Completely Rural	0.5	0.5	0.3	0.5	*	*

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.100A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	1,135	1,145	263	328	80	104
GEOGRAPHIC DIVISION						
Northeast	188	159	55	60	21	16
New England	60	62	29	18	13	3
Middle Atlantic	128	98	25	41	8	13
Midwest	275	275	73	76	26	20
East North Central	158	164	48	57	14	19
West North Central	117	111	25	19	11	1
South	360	395	85	116	26	55
South Atlantic	175	164	45	61	13	24
East South Central	88	99	21	25	5	16
West South Central	97	132	18	30	7	15
West	313	316	50	76	8	13
Mountain	134	143	21	33	5	9
Pacific	179	173	29	43	4	4
COUNTY TYPE						
Large Metro	465	501	85 ^b	151	24 ^a	53
Small Metro	457	396	116	121	39	41
250K – 1 Mil. Pop.	268	238	68	63	24	27
< 250K Pop.	189	157	49	57	15	14
Nonmetro	213	248	62	56	18	11
Urbanized	86	96	27	23	8	5
Less Urbanized	107	127	32	30	10	6
Completely Rural	20	25	*	3	0	0

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.100B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	3.5	3.5	0.8	1.0	0.3	0.3
GEOGRAPHIC DIVISION						
Northeast	3.3	2.8	1.0	1.0	0.4	0.3
New England	4.0	4.1	2.0	1.2	0.9	0.2
Middle Atlantic	3.0	2.3	0.6	1.0	0.2	0.3
Midwest	3.7	3.7	1.0	1.0	0.3	0.3
East North Central	3.1	3.2	0.9	1.1	0.3	0.4
West North Central	5.0	4.8	1.1	0.8	0.5	0.1
South	3.1	3.4	0.7	1.0	0.2	0.5
South Atlantic	3.1	2.8	0.8	1.0	0.2	0.4
East South Central	4.5	5.1	1.1	1.3	0.3	0.8
West South Central	2.5	3.4	0.5	0.8	0.2	0.4
West	4.2	4.1	0.7	1.0	0.1	0.2
Mountain	5.9	6.1	0.9	1.4	0.2	0.4
Pacific	3.4	3.3	0.6	0.8	0.1	0.1
COUNTY TYPE						
Large Metro	2.8	2.9	0.5 ^b	0.9	0.1 ^a	0.3
Small Metro	4.3	3.8	1.1	1.2	0.4	0.4
250K – 1 Mil. Pop.	3.9	3.5	1.0	0.9	0.3	0.4
< 250K Pop.	5.1	4.4	1.3	1.6	0.4	0.4
Nonmetro	4.3	5.0	1.2	1.1	0.4	0.2
Urbanized	3.7	4.6	1.2	1.1	0.3	0.2
Less Urbanized	4.6	5.4	1.4	1.3	0.4	0.3
Completely Rural	6.1	5.1	*	0.6	0.0	0.1

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.101A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	6,580	6,675	978	1,000	370	553
GEOGRAPHIC DIVISION						
Northeast	1,107	1,179	127	206	63 ^a	158
New England	283	355	11	40	1	32
Middle Atlantic	824	825	116	166	63	126
Midwest	1,391	1,216	337	217	127	124
East North Central	1,020	784	268 ^a	136	113	47
West North Central	371	432	68	81	15	77
South	2,540	2,385	385	398	175	172
South Atlantic	1,533	1,375	269	226	129	90
East South Central	413	347	81	83	27	29
West South Central	593	664	34	88	19	53
West	1,543	1,894	130	179	5 ^a	98
Mountain	457	485	22 ^a	115	5	65
Pacific	1,086	1,410	109	64	*	34
COUNTY TYPE						
Large Metro	3,640	3,907	549	587	217	327
Small Metro	1,927	1,860	283	318	101	197
250K – 1 Mil. Pop.	1,345	1,188	183	245	58 ^a	158
< 250K Pop.	582	672	100	74	44	39
Nonmetro	1,013	908	146	95	53	28
Urbanized	560 ^a	328	67	28	29	7
Less Urbanized	415	492	68	66	20	21
Completely Rural	38	88	10	1	4	1

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.101B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	3.6	3.6	0.5	0.5	0.2	0.3
GEOGRAPHIC DIVISION						
Northeast	3.1	3.3	0.4	0.6	0.2 ^a	0.4
New England	3.0	3.8	0.1	0.4	0.0	0.3
Middle Atlantic	3.2	3.2	0.4	0.6	0.2	0.5
Midwest	3.4	2.9	0.8	0.5	0.3	0.3
East North Central	3.5 ^a	2.7	0.9 ^a	0.5	0.4	0.2
West North Central	3.0	3.5	0.6	0.6	0.1	0.6
South	3.9	3.6	0.6	0.6	0.3	0.3
South Atlantic	4.4	3.9	0.8	0.6	0.4	0.3
East South Central	3.8	3.1	0.7	0.7	0.2	0.3
West South Central	3.0	3.3	0.2	0.4	0.1	0.3
West	3.8	4.5	0.3	0.4	0.0 ^a	0.2
Mountain	3.8	3.9	0.2 ^a	0.9	0.0	0.5
Pacific	3.7	4.8	0.4	0.2	*	0.1
COUNTY TYPE						
Large Metro	3.7	3.9	0.6	0.6	0.2	0.3
Small Metro	3.6	3.5	0.5	0.6	0.2	0.4
250K – 1 Mil. Pop.	3.8	3.3	0.5	0.7	0.2 ^a	0.4
< 250K Pop.	3.2	3.7	0.5	0.4	0.2	0.2
Nonmetro	3.2	2.9	0.5	0.3	0.2	0.1
Urbanized	4.4 ^a	2.8	0.5	0.2	0.2	0.1
Less Urbanized	2.6	3.1	0.4	0.4	0.1	0.1
Completely Rural	1.3	2.3	0.4	0.0	0.1	0.0

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.102A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	7,715	7,820	1,241	1,327	451	657
GEOGRAPHIC DIVISION						
Northeast	1,294	1,338	182	266	84	174
New England	342	416	40	58	13	36
Middle Atlantic	952	922	141	208	71	138
Midwest	1,665	1,491	410	293	153	144
East North Central	1,177	948	316	193	127	66
West North Central	488	543	94	99	26	78
South	2,899	2,780	469	514	200	227
South Atlantic	1,708	1,539	314	287	142	114
East South Central	501	446	103	109	32	45
West South Central	690	796	52 ^a	118	26	68
West	1,856	2,210	181	255	13 ^a	112
Mountain	591	628	43 ^a	147	9	74
Pacific	1,265	1,582	138	108	4	37
COUNTY TYPE						
Large Metro	4,105	4,408	634	738	241	380
Small Metro	2,384	2,256	399	439	140	238
250K – 1 Mil. Pop.	1,613	1,426	251	308	81 ^a	185
< 250K Pop.	771	829	148	131	58	53
Nonmetro	1,225	1,156	208	151	70	40
Urbanized	646 ^a	424	95	51	36	12
Less Urbanized	522	619	101	96	30	27
Completely Rural	57	113	12	4	4	1

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.102B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	3.6	3.6	0.6	0.6	0.2	0.3
GEOGRAPHIC DIVISION						
Northeast	3.2	3.3	0.4	0.6	0.2	0.4
New England	3.2	3.9	0.4	0.5	0.1	0.3
Middle Atlantic	3.2	3.0	0.5	0.7	0.2	0.5
Midwest	3.4	3.1	0.8	0.6	0.3	0.3
East North Central	3.5	2.8	0.9	0.6	0.4	0.2
West North Central	3.3	3.7	0.6	0.7	0.2	0.5
South	3.8	3.5	0.6	0.7	0.3	0.3
South Atlantic	4.2	3.7	0.8	0.7	0.3	0.3
East South Central	3.9	3.4	0.8	0.8	0.3	0.3
West South Central	2.9	3.3	0.2 ^a	0.5	0.1	0.3
West	3.8	4.5	0.4	0.5	0.0 ^a	0.2
Mountain	4.1	4.3	0.3 ^a	1.0	0.1	0.5
Pacific	3.7	4.6	0.4	0.3	0.0	0.1
COUNTY TYPE						
Large Metro	3.6	3.8	0.6	0.6	0.2	0.3
Small Metro	3.7	3.5	0.6	0.7	0.2	0.4
250K – 1 Mil. Pop.	3.8	3.3	0.6	0.7	0.2 ^a	0.4
< 250K Pop.	3.5	3.9	0.7	0.6	0.3	0.2
Nonmetro	3.4	3.2	0.6	0.4	0.2	0.1
Urbanized	4.3	3.1	0.6	0.4	0.2	0.1
Less Urbanized	2.9	3.4	0.6	0.5	0.2	0.1
Completely Rural	1.8	2.7	0.4	0.1	0.1	0.0

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.