

Table 1.113C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	729	712	353	342	210	229
GEOGRAPHIC DIVISION						
Northeast	358	323	152	146	81	103
New England	171	124	67	65	42	41
Middle Atlantic	316	299	136	131	69	94
Midwest	263	247	129	131	87	84
East North Central	220	199	107	107	73	69
West North Central	144	151	74	78	49	48
South	440	429	198	220	127	142
South Atlantic	336	332	157	168	100	104
East South Central	155	173	65	89	46	57
West South Central	238	216	103	109	63	76
West	376	364	214	187	117	117
Mountain	152	172	100	89	65	51
Pacific	344	323	189	164	97	106
COUNTY TYPE						
Large Metro	667	625	282	271	151	180
Small Metro	453	482	188	197	122	116
250K – 1 Mil. Pop.	376	398	155	169	105	99
< 250K Pop.	266	258	114	106	72	64
Nonmetro	264	278	130	131	75	86
Urbanized	181	165	93	93	53	61
Less Urbanized	194	215	97	93	55	61
Completely Rural	98	105	40	40	20	26

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.113D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.30	0.29	0.15	0.14	0.09	0.09
GEOGRAPHIC DIVISION						
Northeast	0.79	0.71	0.33	0.32	0.18	0.23
New England	1.43	1.04	0.56	0.55	0.35	0.34
Middle Atlantic	0.94	0.89	0.41	0.39	0.21	0.28
Midwest	0.48	0.45	0.24	0.24	0.16	0.15
East North Central	0.58	0.52	0.28	0.28	0.19	0.18
West North Central	0.88	0.92	0.45	0.48	0.30	0.30
South	0.51	0.49	0.23	0.25	0.15	0.16
South Atlantic	0.74	0.72	0.35	0.37	0.22	0.23
East South Central	1.08	1.19	0.45	0.61	0.32	0.39
West South Central	0.90	0.80	0.39	0.40	0.24	0.28
West	0.69	0.66	0.39	0.34	0.21	0.21
Mountain	0.95	1.04	0.62	0.54	0.41	0.31
Pacific	0.89	0.83	0.49	0.42	0.25	0.27
COUNTY TYPE						
Large Metro	0.45	0.41	0.21	0.20	0.11	0.13
Small Metro	0.49	0.54	0.24	0.25	0.17	0.16
250K – 1 Mil. Pop.	0.60	0.66	0.31	0.33	0.22	0.20
< 250K Pop.	0.82	0.86	0.40	0.40	0.29	0.26
Nonmetro	0.58	0.54	0.31	0.31	0.18	0.20
Urbanized	0.92	0.83	0.55	0.54	0.32	0.36
Less Urbanized	0.79	0.80	0.44	0.43	0.26	0.29
Completely Rural	2.00	1.59	0.96	0.77	0.51	0.47

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.113P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.8785		0.4599		0.2904	
GEOGRAPHIC DIVISION						
Northeast	0.5623		0.6571		0.3216	
New England	0.8071		0.3267		0.2580	
Middle Atlantic	0.5998		0.9932		0.5892	
Midwest	0.9349		0.6735		0.5329	
East North Central	0.4090		0.4599		0.9147	
West North Central	0.2991		0.0776		0.3454	
South	0.1310		0.3670		0.8755	
South Atlantic	0.1823		0.5457		0.8444	
East South Central	0.5782		0.6336		0.6170	
West South Central	0.5994		0.2246		0.6933	
West	0.1147		0.7758		0.5858	
Mountain	0.3471		0.9919		0.7471	
Pacific	0.2002		0.7452		0.4205	
COUNTY TYPE						
Large Metro	0.8728		0.4317		0.0808	
Small Metro	0.8825		0.9190		0.4159	
250K – 1 Mil. Pop.	0.8692		0.6729		0.8919	
< 250K Pop.	0.9943		0.6219		0.2139	
Nonmetro	0.5247		0.7853		0.5307	
Urbanized	0.3939		0.4207		0.5448	
Less Urbanized	0.5998		0.7606		0.8114	
Completely Rural	0.6888		0.8932		0.7511	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.114C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	70	72	59	61	40	37
GEOGRAPHIC DIVISION						
Northeast	32	27	26	22	18	14
New England	17	13	12	11	10	8
Middle Atlantic	27	23	23	19	15	12
Midwest	29	28	25	24	16	16
East North Central	21	22	20	19	13	12
West North Central	19	16	15	13	8	10
South	44	44	37	38	25	25
South Atlantic	31	31	27	26	18	20
East South Central	20	18	14	17	9	11
West South Central	24	25	21	22	15	12
West	34	40	29	35	20	20
Mountain	15	20	10	17	7	7
Pacific	31	35	27	31	19	18
COUNTY TYPE						
Large Metro	61	59	48	49	32	28
Small Metro	45	42	35	35	22	21
250K – 1 Mil. Pop.	35	35	25	29	16	18
< 250K Pop.	33	24	25	20	16	11
Nonmetro	37	34	26	28	17	17
Urbanized	21	22	16	18	9	12
Less Urbanized	29	25	21	19	13	11
Completely Rural	10	12	6	9	5	5

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.114D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.28	0.28	0.23	0.24	0.16	0.14
GEOGRAPHIC DIVISION						
Northeast	0.70	0.59	0.58	0.48	0.39	0.31
New England	1.46	1.14	1.04	0.95	0.83	0.66
Middle Atlantic	0.79	0.69	0.69	0.57	0.44	0.35
Midwest	0.50	0.49	0.44	0.42	0.28	0.29
East North Central	0.53	0.56	0.49	0.48	0.33	0.31
West North Central	1.14	0.96	0.92	0.81	0.50	0.63
South	0.49	0.49	0.41	0.42	0.28	0.28
South Atlantic	0.68	0.66	0.58	0.57	0.40	0.42
East South Central	1.40	1.27	0.94	1.16	0.62	0.75
West South Central	0.81	0.84	0.70	0.73	0.49	0.41
West	0.57	0.67	0.48	0.58	0.33	0.32
Mountain	0.83	1.12	0.58	0.95	0.39	0.42
Pacific	0.73	0.82	0.63	0.72	0.44	0.42
COUNTY TYPE						
Large Metro	0.41	0.39	0.34	0.34	0.24	0.20
Small Metro	0.50	0.51	0.42	0.43	0.28	0.27
250K – 1 Mil. Pop.	0.58	0.62	0.48	0.52	0.30	0.33
< 250K Pop.	0.99	0.90	0.78	0.75	0.56	0.44
Nonmetro	0.68	0.66	0.51	0.57	0.36	0.38
Urbanized	1.01	1.08	0.74	0.97	0.47	0.71
Less Urbanized	1.05	0.95	0.81	0.76	0.57	0.49
Completely Rural	2.02	1.98	1.33	1.61	1.15	1.01

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.114P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.0001		0.1208		0.1403	
GEOGRAPHIC DIVISION						
Northeast	0.0169		0.1672		0.2916	
New England	0.1898		0.2344		0.4412	
Middle Atlantic	0.0463		0.3589		0.4505	
Midwest	0.0478		0.3166		0.9437	
East North Central	0.0047		0.1286		0.5140	
West North Central	0.7334		0.7166		0.4441	
South	0.2288		0.4913		0.6086	
South Atlantic	0.1162		0.6564		0.6620	
East South Central	0.9071		0.3371		0.1165	
West South Central	0.9409		0.2782		0.4168	
West	0.0064		0.0558		0.0096	
Mountain	0.7342		0.3373		0.6745	
Pacific	0.0012		0.0103		0.0031	
COUNTY TYPE						
Large Metro	0.0310		0.3346		0.1144	
Small Metro	0.0022		0.2574		0.6235	
250K – 1 Mil. Pop.	0.0297		0.6090		0.9974	
< 250K Pop.	0.0321		0.2270		0.4111	
Nonmetro	0.1203		0.6548		0.9592	
Urbanized	0.6371		0.6819		0.6636	
Less Urbanized	0.1827		0.6672		0.8657	
Completely Rural	0.5399		0.4489		0.8652	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.115C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	141	142	110	108	68	72
GEOGRAPHIC DIVISION						
Northeast	62	57	50	43	27	28
New England	28	26	26	23	15	15
Middle Atlantic	56	50	43	34	23	22
Midwest	58	63	45	44	26	26
East North Central	47	53	35	36	22	22
West North Central	35	34	28	26	15	14
South	84	88	63	68	40	46
South Atlantic	52	67	39	48	28	32
East South Central	38	31	21	24	14	18
West South Central	54	48	44	40	25	27
West	74	77	60	60	41	38
Mountain	32	35	25	29	15	19
Pacific	67	69	54	52	38	33
COUNTY TYPE						
Large Metro	134	149	84	95	48	53
Small Metro	120	113	87	77	50	47
250K – 1 Mil. Pop.	95	94	68	64	37	39
< 250K Pop.	82	64	58	42	34	27
Nonmetro	72	71	43	45	27	28
Urbanized	55	54	33	34	19	19
Less Urbanized	44	44	30	27	19	17
Completely Rural	24	22	16	15	*	8

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.115D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.44	0.44	0.34	0.33	0.21	0.22
GEOGRAPHIC DIVISION						
Northeast	1.10	1.00	0.88	0.75	0.48	0.48
New England	1.85	1.71	1.73	1.52	0.99	0.98
Middle Atlantic	1.33	1.19	1.02	0.81	0.55	0.53
Midwest	0.78	0.84	0.60	0.60	0.35	0.35
East North Central	0.91	1.03	0.69	0.70	0.42	0.42
West North Central	1.49	1.47	1.21	1.11	0.63	0.61
South	0.73	0.75	0.54	0.58	0.35	0.39
South Atlantic	0.91	1.14	0.69	0.83	0.49	0.54
East South Central	1.94	1.57	1.08	1.23	0.74	0.95
West South Central	1.40	1.21	1.13	1.01	0.63	0.68
West	0.99	1.01	0.80	0.78	0.54	0.50
Mountain	1.40	1.47	1.11	1.24	0.65	0.81
Pacific	1.28	1.30	1.03	0.99	0.73	0.63
COUNTY TYPE						
Large Metro	0.61	0.61	0.45	0.46	0.27	0.28
Small Metro	0.78	0.77	0.66	0.61	0.40	0.42
250K – 1 Mil. Pop.	0.98	0.95	0.83	0.75	0.46	0.52
< 250K Pop.	1.33	1.33	1.14	1.02	0.77	0.72
Nonmetro	1.06	1.04	0.72	0.79	0.54	0.53
Urbanized	1.70	1.77	1.15	1.27	0.78	0.83
Less Urbanized	1.38	1.39	1.08	1.05	0.78	0.68
Completely Rural	4.38	2.52	3.46	2.29	*	1.57

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.115P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.0585		0.5494		0.5633	
GEOGRAPHIC DIVISION						
Northeast	0.7649		0.4507		0.9572	
New England	0.8734		0.7083		0.5399	
Middle Atlantic	0.7968		0.2545		0.6450	
Midwest	0.7891		0.6084		0.4371	
East North Central	0.9224		0.2731		0.3755	
West North Central	0.7389		0.5461		0.9485	
South	0.1142		0.2499		0.3641	
South Atlantic	0.7035		0.6283		0.6963	
East South Central	0.6015		0.9764		0.5328	
West South Central	0.0586		0.1960		0.1125	
West	0.1999		0.4231		0.5629	
Mountain	0.0875		0.8031		0.4937	
Pacific	0.5615		0.4439		0.7568	
COUNTY TYPE						
Large Metro	0.0762		0.1481		0.2091	
Small Metro	0.4317		0.7641		0.8982	
250K – 1 Mil. Pop.	0.0919		0.8806		0.4746	
< 250K Pop.	0.3332		0.4753		0.4994	
Nonmetro	0.5781		0.5934		0.4333	
Urbanized	0.4194		0.7379		0.5516	
Less Urbanized	0.9674		0.2657		0.1849	
Completely Rural	0.7968		0.7495		*	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.116C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	700	686	313	310	187	215
GEOGRAPHIC DIVISION						
Northeast	343	313	126	135	67	99
New England	162	123	48	56	31	39
Middle Atlantic	302	287	116	123	59	91
Midwest	248	236	116	119	81	79
East North Central	212	190	97	95	65	65
West North Central	129	144	64	72	48	45
South	419	407	182	194	119	127
South Atlantic	326	318	149	156	89	96
East South Central	155	163	63	77	50	51
West South Central	215	201	83	90	62	66
West	368	348	189	166	100	109
Mountain	153	168	84	79	58	47
Pacific	335	307	170	146	81	98
COUNTY TYPE						
Large Metro	619	589	243	248	131	169
Small Metro	409	450	156	174	108	107
250K – 1 Mil. Pop.	345	375	136	152	95	92
< 250K Pop.	221	230	82	84	57	54
Nonmetro	241	253	121	109	67	77
Urbanized	158	149	82	79	46	52
Less Urbanized	177	193	83	79	47	58
Completely Rural	85	94	35	31	14	23

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.116D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.38	0.37	0.17	0.17	0.10	0.12
GEOGRAPHIC DIVISION						
Northeast	0.97	0.88	0.36	0.38	0.19	0.28
New England	1.75	1.32	0.51	0.60	0.34	0.42
Middle Atlantic	1.16	1.10	0.45	0.47	0.23	0.35
Midwest	0.60	0.57	0.28	0.29	0.20	0.19
East North Central	0.74	0.66	0.34	0.33	0.23	0.22
West North Central	1.05	1.15	0.52	0.58	0.39	0.36
South	0.64	0.61	0.28	0.29	0.18	0.19
South Atlantic	0.93	0.89	0.43	0.44	0.25	0.27
East South Central	1.41	1.46	0.57	0.69	0.45	0.46
West South Central	1.09	1.00	0.42	0.45	0.31	0.33
West	0.90	0.83	0.46	0.40	0.24	0.26
Mountain	1.28	1.36	0.70	0.64	0.48	0.38
Pacific	1.15	1.04	0.58	0.50	0.28	0.33
COUNTY TYPE						
Large Metro	0.56	0.52	0.24	0.24	0.13	0.17
Small Metro	0.63	0.69	0.28	0.31	0.20	0.19
250K – 1 Mil. Pop.	0.78	0.84	0.37	0.41	0.27	0.26
< 250K Pop.	1.00	1.08	0.43	0.44	0.32	0.29
Nonmetro	0.71	0.68	0.38	0.34	0.21	0.24
Urbanized	1.10	1.05	0.65	0.64	0.37	0.42
Less Urbanized	1.00	0.98	0.51	0.48	0.29	0.35
Completely Rural	2.39	1.95	1.16	0.79	0.48	0.57

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.116P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.8604		0.3455		0.2107	
GEOGRAPHIC DIVISION						
Northeast	0.4479		0.3046		0.1921	
New England	0.7176		0.1828		0.2035	
Middle Atlantic	0.5041		0.5893		0.3949	
Midwest	0.9356		0.3653		0.3375	
East North Central	0.5688		0.9324		0.5767	
West North Central	0.3235		0.0937		0.3853	
South	0.0732		0.6206		0.8167	
South Atlantic	0.1902		0.5584		0.7990	
East South Central	0.5076		0.4746		0.3114	
West South Central	0.3043		0.5309		0.9888	
West	0.1000		0.8118		0.3729	
Mountain	0.5861		0.8072		0.5289	
Pacific	0.1208		0.8813		0.1636	
COUNTY TYPE						
Large Metro	0.7543		0.5938		0.0761	
Small Metro	0.7287		0.5093		0.4594	
250K – 1 Mil. Pop.	0.9594		0.5313		0.7414	
< 250K Pop.	0.5865		0.8427		0.3762	
Nonmetro	0.4670		0.5752		0.3313	
Urbanized	0.4523		0.3886		0.6975	
Less Urbanized	0.4845		0.9344		0.4680	
Completely Rural	0.7016		0.9562		0.4023	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.117C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	724	714	341	341	203	229
GEOGRAPHIC DIVISION						
Northeast	355	320	142	144	75	103
New England	173	122	62	64	38	42
Middle Atlantic	311	296	128	129	65	95
Midwest	258	246	125	130	86	82
East North Central	218	200	104	106	71	68
West North Central	139	148	70	77	49	46
South	437	426	195	216	124	140
South Atlantic	337	329	156	167	95	102
East South Central	151	169	62	88	49	57
West South Central	235	215	99	105	63	76
West	374	364	207	185	113	117
Mountain	154	173	97	86	65	50
Pacific	342	322	183	163	93	106
COUNTY TYPE						
Large Metro	655	620	272	270	145	180
Small Metro	439	477	178	192	118	115
250K – 1 Mil. Pop.	365	393	150	164	102	98
< 250K Pop.	251	252	104	102	67	62
Nonmetro	256	268	126	123	73	83
Urbanized	178	158	90	88	53	58
Less Urbanized	182	208	90	88	51	59
Completely Rural	95	101	39	38	20	25

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.117D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.34	0.33	0.16	0.16	0.09	0.11
GEOGRAPHIC DIVISION						
Northeast	0.87	0.78	0.35	0.35	0.18	0.25
New England	1.61	1.13	0.57	0.60	0.35	0.39
Middle Atlantic	1.03	0.98	0.42	0.43	0.21	0.31
Midwest	0.53	0.50	0.26	0.27	0.18	0.17
East North Central	0.64	0.59	0.31	0.31	0.21	0.20
West North Central	0.95	1.00	0.48	0.52	0.33	0.31
South	0.57	0.54	0.25	0.28	0.16	0.18
South Atlantic	0.83	0.79	0.38	0.40	0.23	0.25
East South Central	1.17	1.29	0.48	0.67	0.38	0.44
West South Central	1.00	0.90	0.42	0.44	0.27	0.32
West	0.77	0.74	0.43	0.37	0.23	0.24
Mountain	1.08	1.18	0.68	0.58	0.45	0.34
Pacific	0.99	0.93	0.53	0.47	0.27	0.30
COUNTY TYPE						
Large Metro	0.49	0.46	0.23	0.22	0.12	0.15
Small Metro	0.55	0.60	0.26	0.28	0.19	0.17
250K – 1 Mil. Pop.	0.67	0.73	0.33	0.36	0.24	0.23
< 250K Pop.	0.89	0.96	0.43	0.43	0.31	0.28
Nonmetro	0.64	0.60	0.35	0.33	0.20	0.22
Urbanized	1.03	0.93	0.60	0.59	0.36	0.39
Less Urbanized	0.87	0.89	0.47	0.46	0.27	0.31
Completely Rural	2.25	1.79	1.07	0.84	0.56	0.54

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.117P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.5944		0.3012		0.1807	
GEOGRAPHIC DIVISION						
Northeast	0.4289		0.4857		0.2285	
New England	0.7055		0.2110		0.1745	
Middle Atlantic	0.4875		0.8717		0.4992	
Midwest	0.8975		0.5293		0.5112	
East North Central	0.5909		0.6386		0.8168	
West North Central	0.3115		0.0778		0.4103	
South	0.1590		0.4234		0.9464	
South Atlantic	0.2305		0.4929		0.9033	
East South Central	0.5765		0.5005		0.4360	
West South Central	0.5968		0.2982		0.5766	
West	0.0635		0.9832		0.3163	
Mountain	0.3790		0.8806		0.7057	
Pacific	0.1056		0.9244		0.1710	
COUNTY TYPE						
Large Metro	0.9578		0.3330		0.0390	
Small Metro	0.6604		0.7365		0.4642	
250K – 1 Mil. Pop.	0.7054		0.5956		0.8931	
< 250K Pop.	0.8214		0.8159		0.2614	
Nonmetro	0.4052		0.7160		0.5281	
Urbanized	0.3638		0.3744		0.5932	
Less Urbanized	0.4792		0.8212		0.7732	
Completely Rural	0.7436		1.0000		0.7205	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.