

Table 1.118C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	564	562	300	304	170	194
GEOGRAPHIC DIVISION						
Northeast	252	252	133	135	62	90
New England	116	106	45	59	24	37
Middle Atlantic	227	230	125	122	57	82
Midwest	241	199	123	118	77	71
East North Central	199	171	104	97	64	61
West North Central	136	105	66	69	44	36
South	322	347	167	193	107	121
South Atlantic	248	270	134	150	77	89
East South Central	113	137	58	73	46	47
West South Central	171	177	81	98	59	66
West	306	294	171	163	86	92
Mountain	119	131	86	77	43	42
Pacific	282	264	148	144	74	82
COUNTY TYPE						
Large Metro	500	467	240	248	126	152
Small Metro	318	350	158	169	97	99
250K – 1 Mil. Pop.	259	296	130	150	79	89
< 250K Pop.	203	181	97	79	59	47
Nonmetro	219	221	115	110	63	66
Urbanized	156	140	77	81	38	43
Less Urbanized	150	163	87	72	52	43
Completely Rural	74	70	38	30	18	23

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.118D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.23	0.23	0.12	0.12	0.07	0.08
GEOGRAPHIC DIVISION						
Northeast	0.55	0.55	0.29	0.30	0.14	0.20
New England	0.97	0.89	0.38	0.49	0.20	0.31
Middle Atlantic	0.68	0.68	0.37	0.36	0.17	0.24
Midwest	0.45	0.37	0.23	0.22	0.14	0.13
East North Central	0.53	0.45	0.27	0.25	0.17	0.16
West North Central	0.84	0.64	0.41	0.42	0.27	0.22
South	0.37	0.40	0.19	0.22	0.12	0.14
South Atlantic	0.55	0.59	0.30	0.32	0.17	0.19
East South Central	0.78	0.94	0.40	0.50	0.32	0.32
West South Central	0.64	0.66	0.30	0.36	0.22	0.24
West	0.56	0.53	0.31	0.29	0.16	0.17
Mountain	0.74	0.80	0.54	0.47	0.27	0.26
Pacific	0.73	0.68	0.38	0.37	0.19	0.21
COUNTY TYPE						
Large Metro	0.34	0.33	0.18	0.18	0.09	0.11
Small Metro	0.36	0.41	0.20	0.22	0.13	0.13
250K – 1 Mil. Pop.	0.42	0.53	0.25	0.29	0.16	0.18
< 250K Pop.	0.64	0.64	0.34	0.31	0.24	0.19
Nonmetro	0.49	0.45	0.28	0.26	0.16	0.16
Urbanized	0.82	0.72	0.45	0.46	0.23	0.26
Less Urbanized	0.65	0.64	0.41	0.34	0.25	0.21
Completely Rural	1.75	1.11	0.96	0.56	0.48	0.43

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.118P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.4798		0.3133		0.4302	
GEOGRAPHIC DIVISION						
Northeast	0.4405		0.9960		0.5723	
New England	0.6412		0.1694		0.1112	
Middle Atlantic	0.2806		0.5550		0.9380	
Midwest	0.8443		0.4755		0.6362	
East North Central	0.7378		0.8545		0.9859	
West North Central	0.8819		0.1219		0.3963	
South	0.8750		0.2416		0.6738	
South Atlantic	0.8722		0.5872		0.7090	
East South Central	0.7794		0.8069		0.8701	
West South Central	0.8860		0.0926		0.6873	
West	0.4875		0.9708		0.8628	
Mountain	0.4989		0.8190		0.7994	
Pacific	0.6536		0.8710		0.9618	
COUNTY TYPE						
Large Metro	0.5118		0.2129		0.1876	
Small Metro	0.8381		0.7298		0.8247	
250K – 1 Mil. Pop.	0.3925		0.2472		0.4222	
< 250K Pop.	0.3693		0.2071		0.0944	
Nonmetro	0.7429		0.7458		0.8724	
Urbanized	0.8079		0.6418		0.8819	
Less Urbanized	0.8081		0.4987		0.6635	
Completely Rural	0.5962		0.8265		0.7612	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 1.119C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	67	65	56	56	36	33
GEOGRAPHIC DIVISION						
Northeast	29	22	26	20	17	12
New England	16	11	11	10	9	5
Middle Atlantic	24	19	24	17	14	11
Midwest	28	26	24	22	14	15
East North Central	22	21	19	18	13	12
West North Central	18	15	15	12	6	9
South	42	42	34	35	23	23
South Atlantic	30	29	25	25	17	17
East South Central	19	18	13	16	8	10
West South Central	22	23	19	19	13	12
West	33	36	28	32	18	17
Mountain	15	17	11	16	7	7
Pacific	30	31	25	28	17	16
COUNTY TYPE						
Large Metro	57	51	45	44	28	25
Small Metro	43	38	34	31	20	17
250K – 1 Mil. Pop.	33	33	24	27	15	15
< 250K Pop.	31	21	23	15	13	9
Nonmetro	34	31	25	26	17	16
Urbanized	19	20	15	17	9	12
Less Urbanized	26	22	20	18	13	10
Completely Rural	9	11	5	9	4	5

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 1.119D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
Standard Errors of Percentages, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.27	0.26	0.22	0.22	0.14	0.13
GEOGRAPHIC DIVISION						
Northeast	0.64	0.49	0.58	0.44	0.37	0.26
New England	1.39	0.98	0.93	0.84	0.78	0.40
Middle Atlantic	0.72	0.57	0.71	0.51	0.42	0.32
Midwest	0.49	0.46	0.42	0.39	0.25	0.27
East North Central	0.54	0.52	0.46	0.46	0.33	0.29
West North Central	1.08	0.89	0.88	0.72	0.36	0.57
South	0.47	0.46	0.37	0.39	0.25	0.25
South Atlantic	0.66	0.63	0.54	0.53	0.36	0.37
East South Central	1.34	1.24	0.88	1.08	0.57	0.67
West South Central	0.75	0.78	0.63	0.65	0.44	0.40
West	0.56	0.59	0.46	0.52	0.30	0.29
Mountain	0.84	0.99	0.64	0.90	0.38	0.41
Pacific	0.70	0.72	0.59	0.64	0.39	0.37
COUNTY TYPE						
Large Metro	0.38	0.34	0.32	0.30	0.21	0.18
Small Metro	0.49	0.47	0.40	0.38	0.24	0.22
250K – 1 Mil. Pop.	0.57	0.59	0.45	0.49	0.28	0.27
< 250K Pop.	0.97	0.80	0.75	0.59	0.48	0.35
Nonmetro	0.63	0.62	0.49	0.55	0.37	0.36
Urbanized	0.92	1.04	0.71	0.94	0.52	0.69
Less Urbanized	0.95	0.86	0.77	0.72	0.57	0.44
Completely Rural	1.93	1.90	1.17	1.49	1.09	0.99

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.119P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2005 Versus 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.0001		0.0895		0.1652	
GEOGRAPHIC DIVISION						
Northeast	0.0046		0.0946		0.1539	
New England	0.0889		0.0673		0.0556	
Middle Atlantic	0.0232		0.3427		0.5859	
Midwest	0.0787		0.5283		0.7086	
East North Central	0.0265		0.3421		0.7131	
West North Central	0.9774		0.8222		0.2056	
South	0.2265		0.7146		0.7324	
South Atlantic	0.1979		0.8156		0.6775	
East South Central	0.4808		0.8329		0.3010	
West South Central	0.9685		0.4111		0.5020	
West	0.0061		0.0669		0.0208	
Mountain	0.9166		0.2382		0.2530	
Pacific	0.0023		0.0079		0.0027	
COUNTY TYPE						
Large Metro	0.0148		0.1741		0.2985	
Small Metro	0.0048		0.3248		0.3889	
250K – 1 Mil. Pop.	0.0540		0.9339		0.7333	
< 250K Pop.	0.0368		0.1032		0.3163	
Nonmetro	0.1285		0.7642		0.7315	
Urbanized	0.5117		0.7131		0.7223	
Less Urbanized	0.1489		0.7075		0.4581	
Completely Rural	0.9648		0.2697		0.9854	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 1.120C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	129	131	94	98	60	61
GEOGRAPHIC DIVISION						
Northeast	55	52	40	38	21	23
New England	23	23	19	21	10	11
Middle Atlantic	50	46	35	31	18	20
Midwest	57	58	40	41	24	22
East North Central	48	49	33	32	21	19
West North Central	32	33	22	24	12	12
South	75	81	53	63	35	40
South Atlantic	49	62	33	44	25	28
East South Central	32	29	21	23	14	17
West South Central	46	44	35	37	20	22
West	71	73	53	54	36	32
Mountain	29	34	21	27	12	18
Pacific	64	65	48	46	34	26
COUNTY TYPE						
Large Metro	119	129	73	84	42	44
Small Metro	104	102	72	67	40	38
250K – 1 Mil. Pop.	84	83	58	55	31	33
< 250K Pop.	66	60	45	37	26	20
Nonmetro	66	63	38	40	26	24
Urbanized	47	47	28	29	18	16
Less Urbanized	43	39	26	26	16	16
Completely Rural	21	20	15	13	*	5

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 1.120D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
Standard Errors of Percentages, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.40	0.40	0.29	0.30	0.19	0.19
GEOGRAPHIC DIVISION						
Northeast	0.96	0.91	0.70	0.66	0.37	0.41
New England	1.54	1.54	1.25	1.38	0.70	0.76
Middle Atlantic	1.18	1.09	0.83	0.72	0.44	0.47
Midwest	0.77	0.78	0.54	0.54	0.32	0.30
East North Central	0.93	0.95	0.65	0.63	0.40	0.36
West North Central	1.38	1.42	0.96	1.03	0.53	0.53
South	0.65	0.69	0.46	0.53	0.31	0.34
South Atlantic	0.86	1.07	0.59	0.76	0.44	0.47
East South Central	1.66	1.47	1.09	1.21	0.74	0.87
West South Central	1.20	1.12	0.90	0.94	0.51	0.57
West	0.94	0.96	0.70	0.70	0.47	0.42
Mountain	1.27	1.43	0.92	1.16	0.50	0.76
Pacific	1.23	1.23	0.92	0.87	0.64	0.50
COUNTY TYPE						
Large Metro	0.57	0.56	0.39	0.42	0.24	0.24
Small Metro	0.68	0.73	0.55	0.55	0.34	0.35
250K – 1 Mil. Pop.	0.87	0.89	0.70	0.67	0.40	0.45
< 250K Pop.	1.10	1.32	0.93	0.92	0.63	0.56
Nonmetro	1.00	0.99	0.66	0.72	0.50	0.47
Urbanized	1.51	1.58	0.98	1.13	0.73	0.69
Less Urbanized	1.34	1.40	0.98	1.00	0.68	0.65
Completely Rural	4.23	2.43	3.43	2.25	*	1.00

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.120P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2005 Versus 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.0332		0.2361		0.8442	
GEOGRAPHIC DIVISION						
Northeast	0.9502		0.3210		0.5341	
New England	0.8859		0.5230		0.9820	
Middle Atlantic	1.0000		0.1199		0.4636	
Midwest	0.6792		0.9346		0.8040	
East North Central	0.8917		0.4077		0.3198	
West North Central	0.5867		0.1861		0.2601	
South	0.0249		0.0524		0.3617	
South Atlantic	0.5284		0.6222		0.5368	
East South Central	0.3969		0.1982		0.3016	
West South Central	0.0110		0.0710		0.1006	
West	0.2982		0.5002		0.9272	
Mountain	0.2069		0.5144		0.1973	
Pacific	0.5914		0.6747		0.4598	
COUNTY TYPE						
Large Metro	0.0806		0.1512		0.5222	
Small Metro	0.2487		0.6362		0.6798	
250K – 1 Mil. Pop.	0.1211		0.5953		0.2703	
< 250K Pop.	0.8719		0.9424		0.4309	
Nonmetro	0.5005		0.8132		0.2834	
Urbanized	0.1927		0.8412		0.9469	
Less Urbanized	0.7872		0.6825		0.4085	
Completely Rural	0.8762		0.6251		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.121C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	529	527	265	274	152	179
GEOGRAPHIC DIVISION						
Northeast	250	243	112	126	51	87
New England	113	103	35	53	15	37
Middle Atlantic	225	221	106	115	49	78
Midwest	218	188	107	107	70	66
East North Central	183	160	91	86	56	57
West North Central	120	98	57	64	42	32
South	297	324	154	168	104	109
South Atlantic	231	256	123	136	68	83
East South Central	106	127	60	62	51	40
West South Central	155	164	69	79	59	59
West	285	274	151	144	69	83
Mountain	115	118	76	67	39	39
Pacific	261	247	130	128	57	73
COUNTY TYPE						
Large Metro	450	431	203	222	112	141
Small Metro	285	320	132	149	84	90
250K – 1 Mil. Pop.	233	275	114	135	68	80
< 250K Pop.	167	158	70	62	49	41
Nonmetro	197	199	106	91	57	58
Urbanized	134	123	69	69	34	36
Less Urbanized	132	147	74	59	44	39
Completely Rural	68	58	34	23	12	22

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.121D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.29	0.28	0.14	0.15	0.08	0.10
GEOGRAPHIC DIVISION						
Northeast	0.71	0.69	0.32	0.36	0.15	0.25
New England	1.22	1.10	0.37	0.57	0.16	0.40
Middle Atlantic	0.87	0.85	0.41	0.44	0.19	0.30
Midwest	0.53	0.45	0.26	0.26	0.17	0.16
East North Central	0.63	0.55	0.32	0.30	0.19	0.20
West North Central	0.98	0.79	0.46	0.52	0.34	0.26
South	0.45	0.49	0.23	0.25	0.16	0.16
South Atlantic	0.66	0.72	0.35	0.38	0.20	0.23
East South Central	0.96	1.14	0.55	0.55	0.46	0.36
West South Central	0.79	0.82	0.35	0.39	0.30	0.29
West	0.69	0.66	0.37	0.35	0.17	0.20
Mountain	0.96	0.96	0.63	0.54	0.33	0.31
Pacific	0.89	0.84	0.45	0.44	0.19	0.25
COUNTY TYPE						
Large Metro	0.42	0.40	0.20	0.22	0.11	0.14
Small Metro	0.45	0.52	0.23	0.26	0.16	0.16
250K – 1 Mil. Pop.	0.56	0.67	0.30	0.36	0.19	0.22
< 250K Pop.	0.78	0.78	0.37	0.34	0.27	0.22
Nonmetro	0.59	0.55	0.33	0.28	0.18	0.18
Urbanized	0.95	0.89	0.54	0.55	0.27	0.31
Less Urbanized	0.77	0.79	0.47	0.36	0.28	0.25
Completely Rural	2.15	1.25	1.13	0.57	0.41	0.53

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.121P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.4574		0.2749		0.2684	
GEOGRAPHIC DIVISION						
Northeast	0.2846		0.5238		0.2715	
New England	0.7697		0.0805		0.0256	
Middle Atlantic	0.1857		0.9828		0.8141	
Midwest	0.7307		0.3488		0.5884	
East North Central	0.5396		0.7484		0.6447	
West North Central	0.7516		0.2348		0.7725	
South	0.5747		0.5720		0.9411	
South Atlantic	0.8709		0.6222		0.5924	
East South Central	0.6977		0.4540		0.4809	
West South Central	0.5740		0.3247		1.0000	
West	0.4102		0.8602		0.4320	
Mountain	0.6883		0.4903		0.6809	
Pacific	0.4720		0.5638		0.2516	
COUNTY TYPE						
Large Metro	0.5868		0.2525		0.1631	
Small Metro	0.6974		0.5557		0.8968	
250K – 1 Mil. Pop.	0.4235		0.2330		0.5449	
< 250K Pop.	0.5888		0.3172		0.2130	
Nonmetro	0.7033		0.8229		0.7190	
Urbanized	0.9839		0.6041		0.9186	
Less Urbanized	0.5948		0.5724		0.9965	
Completely Rural	0.6417		0.6716		0.2998	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.122C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	559	560	289	300	164	191
GEOGRAPHIC DIVISION						
Northeast	251	250	125	133	56	90
New England	119	105	43	58	20	38
Middle Atlantic	224	228	117	120	52	81
Midwest	233	199	117	117	76	69
East North Central	192	171	100	96	63	60
West North Central	132	104	61	68	44	35
South	320	342	164	188	105	118
South Atlantic	247	267	133	146	73	88
East South Central	109	132	58	71	48	45
West South Central	170	175	77	95	59	65
West	305	295	166	162	83	91
Mountain	121	129	85	75	43	42
Pacific	280	265	143	143	71	81
COUNTY TYPE						
Large Metro	490	463	229	244	122	150
Small Metro	306	344	151	162	94	98
250K – 1 Mil. Pop.	250	291	128	145	77	88
< 250K Pop.	190	177	88	76	55	46
Nonmetro	214	212	112	102	61	63
Urbanized	153	132	74	76	37	41
Less Urbanized	143	157	82	66	48	42
Completely Rural	72	65	37	28	18	22

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.122D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.26	0.26	0.13	0.14	0.08	0.09
GEOGRAPHIC DIVISION						
Northeast	0.61	0.61	0.30	0.32	0.14	0.22
New England	1.11	0.97	0.40	0.54	0.19	0.35
Middle Atlantic	0.74	0.75	0.39	0.40	0.17	0.27
Midwest	0.48	0.41	0.24	0.24	0.16	0.14
East North Central	0.57	0.50	0.29	0.28	0.19	0.18
West North Central	0.90	0.71	0.42	0.46	0.30	0.24
South	0.41	0.44	0.21	0.24	0.14	0.15
South Atlantic	0.61	0.64	0.33	0.35	0.18	0.21
East South Central	0.84	1.01	0.45	0.54	0.37	0.35
West South Central	0.72	0.73	0.33	0.40	0.25	0.27
West	0.63	0.60	0.34	0.33	0.17	0.18
Mountain	0.85	0.88	0.60	0.51	0.30	0.28
Pacific	0.81	0.76	0.42	0.41	0.21	0.23
COUNTY TYPE						
Large Metro	0.38	0.36	0.19	0.20	0.10	0.13
Small Metro	0.39	0.46	0.22	0.24	0.14	0.15
250K – 1 Mil. Pop.	0.47	0.58	0.28	0.32	0.18	0.21
< 250K Pop.	0.70	0.71	0.36	0.34	0.25	0.21
Nonmetro	0.54	0.49	0.31	0.27	0.17	0.17
Urbanized	0.91	0.79	0.49	0.50	0.26	0.29
Less Urbanized	0.71	0.71	0.44	0.35	0.26	0.22
Completely Rural	1.94	1.23	1.06	0.61	0.52	0.48

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.122P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.2370		0.1727		0.2787	
GEOGRAPHIC DIVISION						
Northeast	0.2849		0.7665		0.3881	
New England	0.8036		0.0751		0.0356	
Middle Atlantic	0.1910		0.6542		0.9839	
Midwest	0.6779		0.3865		0.6798	
East North Central	0.5490		0.9948		0.9247	
West North Central	0.8787		0.1142		0.5340	
South	0.9929		0.2564		0.7150	
South Atlantic	0.9876		0.5508		0.7637	
East South Central	0.8562		0.7722		0.7212	
West South Central	0.8917		0.1166		0.5859	
West	0.3117		0.7113		0.5164	
Mountain	0.4959		0.6686		0.9433	
Pacific	0.4268		0.5124		0.4900	
COUNTY TYPE						
Large Metro	0.3462		0.1265		0.1226	
Small Metro	0.5690		0.5633		0.9577	
250K – 1 Mil. Pop.	0.2673		0.2303		0.3822	
< 250K Pop.	0.5327		0.3570		0.1291	
Nonmetro	0.5863		0.7842		0.9360	
Urbanized	0.7393		0.5887		0.9565	
Less Urbanized	0.6418		0.5350		0.7869	
Completely Rural	0.5970		0.6710		0.7615	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.