

Table 1.128C Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2004 and 2005

Drug or Method of Administration	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
HALLUCINOGENS	618	608	45	44	130	132	583	573
PCP (Angel Dust, Phencyclidine)	298	289	18	19	42	41	292	284
LSD (Acid)	506	521	24	23	98	96	494	499
Peyote	299	260	11	10	33	37	295	254
Mescaline	340	334	10	9	41	45	335	329
Psilocybin (Mushrooms)	478	440	31	30	108	106	447	419
"Ecstasy" (MDMA)	329	335	31	26	108	113	286	294
INHALANTS	480	460	67	67	111	102	448	436
Amyl Nitrite, "Poppers," Locker Room Odorizers, or "Rush"	303	305	28	29	41	46	297	299
Correction Fluid, Degreaser, or Cleaning Fluid	131	118	37	31	35	34	117	107
Gasoline or Lighter Fluid	174	178	41	41	45	39	161	165
Glue, Shoe Polish, or Toluene	206	171	40	43	40	40	196	157
Halothane, Ether, or Other Anesthetics	103	121	15	13	28	29	98	118
Lacquer Thinner or Other Paint Solvents	127	110	32	27	31	31	112	100
Lighter Gases (Butane, Propane)	79	78	23	23	25	26	70	70
Nitrous Oxide or "Whippets"	336	328	29	27	93	89	311	308
Spray Paints	116	105	35	38	31	32	104	93
Other Aerosol Sprays	137	113	32	32	39	42	127	96
NEEDLE USE^{1,2}	200	228	13	10	32	34	196	225
Heroin Needle Use ^{1,2}	139	150	7	6	18	23	136	149
Cocaine Needle Use ^{1,2}	148	151	4	3	19	20	146	149
Stimulant Needle Use ^{1,2}	145	140	6	5	20	22	145	138
Methamphetamine Needle Use ^{1,2}	131	123	6	4	17	19	131	122
HEROIN	193	226	12	10	36	36	188	223
Smoke Heroin ²	106	121	9	6	23	20	102	118
Sniff or Snort Heroin ²	164	184	10	6	29	28	161	181

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.128D Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Standard Errors of Percentages, 2004 and 2005

Drug or Method of Administration	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
HALLUCINOGENS	0.26	0.25	0.18	0.17	0.40	0.41	0.32	0.31
PCP (Angel Dust, Phencyclidine)	0.12	0.12	0.07	0.08	0.13	0.13	0.16	0.15
LSD (Acid)	0.21	0.21	0.09	0.09	0.31	0.30	0.27	0.27
Peyote	0.12	0.11	0.04	0.04	0.10	0.11	0.16	0.14
Mescaline	0.14	0.14	0.04	0.04	0.13	0.14	0.18	0.18
Psilocybin (Mushrooms)	0.20	0.18	0.12	0.12	0.34	0.33	0.24	0.23
"Ecstasy" (MDMA)	0.14	0.14	0.12	0.10	0.34	0.35	0.16	0.16
INHALANTS	0.20	0.19	0.27	0.26	0.34	0.31	0.24	0.24
Amyl Nitrite, "Poppers," Locker Room Odorizers, or "Rush"	0.13	0.13	0.11	0.11	0.13	0.14	0.16	0.16
Correction Fluid, Degreaser, or Cleaning Fluid	0.05	0.05	0.15	0.12	0.11	0.11	0.06	0.06
Gasoline or Lighter Fluid	0.07	0.07	0.16	0.16	0.14	0.12	0.09	0.09
Glue, Shoe Polish, or Toluene	0.09	0.07	0.16	0.17	0.12	0.12	0.11	0.08
Halothane, Ether, or Other Anesthetics	0.04	0.05	0.06	0.05	0.09	0.09	0.05	0.06
Lacquer Thinner or Other Paint Solvents	0.05	0.05	0.13	0.11	0.10	0.10	0.06	0.05
Lighter Gases (Butane, Propane)	0.03	0.03	0.09	0.09	0.08	0.08	0.04	0.04
Nitrous Oxide or "Whippets"	0.14	0.13	0.12	0.11	0.29	0.27	0.17	0.17
Spray Paints	0.05	0.04	0.14	0.15	0.10	0.10	0.06	0.05
Other Aerosol Sprays	0.06	0.05	0.13	0.13	0.12	0.13	0.07	0.05
NEEDLE USE^{1,2}	0.08	0.09	0.05	0.04	0.10	0.10	0.11	0.12
Heroin Needle Use ^{1,2}	0.06	0.06	0.03	0.03	0.06	0.07	0.07	0.08
Cocaine Needle Use ^{1,2}	0.06	0.06	0.02	0.01	0.06	0.06	0.08	0.08
Stimulant Needle Use ^{1,2}	0.06	0.06	0.03	0.02	0.06	0.07	0.08	0.07
Methamphetamine Needle Use ^{1,2}	0.05	0.05	0.02	0.02	0.05	0.06	0.07	0.07
HEROIN	0.08	0.09	0.05	0.04	0.11	0.11	0.10	0.12
Smoke Heroin ²	0.04	0.05	0.03	0.02	0.07	0.06	0.06	0.06
Sniff or Snort Heroin ²	0.07	0.08	0.04	0.02	0.09	0.09	0.09	0.10

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.128P Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2005 Versus 2004

Drug or Method of Administration	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
HALLUCINOGENS	0.2560		0.0073		0.6030		0.3720	
PCP (Angel Dust, Phencyclidine)	0.5731		0.8147		0.6436		0.5941	
LSD (Acid)	0.0928		0.4217		0.0002		0.3252	
Peyote	0.1449		0.2651		0.9078		0.1432	
Mescaline	0.8365		0.4467		0.8530		0.8573	
Psilocybin (Mushrooms)	0.3463		0.1363		0.4767		0.2814	
"Ecstasy" (MDMA)	0.6124		0.0020		0.9221		0.3478	
INHALANTS	0.6435		0.2099		0.1390		0.9481	
Amyl Nitrite, "Poppers," Locker Room Odorizers, or "Rush"	0.9871		0.9673		0.7418		0.9596	
Correction Fluid, Degreaser, or Cleaning Fluid	0.7209		0.0970		0.7533		0.8190	
Gasoline or Lighter Fluid	0.8695		0.3644		0.0502		0.5656	
Glue, Shoe Polish, or Toluene	0.4920		0.7281		0.1541		0.6365	
Halothane, Ether, or Other Anesthetics	0.1087		0.4894		0.8925		0.0844	
Lacquer Thinner or Other Paint Solvents	0.2891		0.3935		0.3616		0.5053	
Lighter Gases (Butane, Propane)	0.4620		0.9286		0.1431		0.7524	
Nitrous Oxide or "Whippets"	0.4422		0.8771		0.1558		0.2088	
Spray Paints	0.5820		0.6691		0.8666		0.4862	
Other Aerosol Sprays	0.1440		0.8836		0.3917		0.0608	
NEEDLE USE^{1,2}	0.6633		0.1885		0.6431		0.6646	
Heroin Needle Use ^{1,2}	0.4596		0.6794		0.0463		0.6373	
Cocaine Needle Use ^{1,2}	0.8989		0.9328		0.2972		0.7839	
Stimulant Needle Use ^{1,2}	0.2912		0.1504		0.4692		0.2632	
Methamphetamine Needle Use ^{1,2}	0.4536		0.2406		0.6640		0.4386	
HEROIN	0.2364		0.2310		0.4483		0.1623	
Smoke Heroin ²	0.8654		0.2419		0.1550		0.8546	
Sniff or Snort Heroin ²	0.3997		0.0801		0.9011		0.3600	

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.129C Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2004 and 2005

Pain Reliever	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Darvocet [®] , Darvon [®] , or Tylenol [®] with Codeine	441	441	54	47	98	103	418	413
Percocet [®] , Percodan [®] , or Tylox [®]	342	357	30	30	85	88	321	340
Vicodin [®] , Lortab [®] , or Lorcet [®]	399	388	46	44	109	119	362	350
Codeine	237	229	33	30	70	72	222	217
Demerol [®]	145	142	16	14	38	44	138	134
Dilaudid [®]	94	106	4	4	14	21	94	105
Fioricet [®]	66	61	6	4	11	19	65	58
Fiorinal [®]	57	58	4	7	8	15	56	54
Hydrocodone	205	224	27	27	78	81	181	203
Methadone	93	117	14	16	33	37	86	108
Morphine	110	142	18	18	45	49	100	128
OxyContin [®]	134	142	24	21	54	64	120	120
Phenaphen [®] with Codeine	61	82	8	8	20	26	57	77
Propoxyphene	56	41	7	5	15	16	52	37
SK-65 [®]	12	20	4	2	5	6	10	19
Stadol [®]	37	22	2	*	10	8	36	20
Talacen [®]	34	19	4	4	7	6	33	17
Talwin [®]	70	56	5	4	6	7	70	55
Talwin [®] NX	33	14	5	3	6	9	32	11
Tramadol	48	40	6	7	15	19	45	33
Ultram [®]	94	77	9	8	25	30	90	69
SELECTED GROUPS OF DRUGS								
Propoxyphene or Codeine Products ^{1,2}	457	461	56	50	103	107	432	434
Oxycodone Products ^{1,3}	357	361	35	34	91	97	329	342
Hydrocodone Products ^{1,4}	408	400	49	47	111	123	368	364
Tramadol Products ^{1,5}	100	84	10	9	28	34	97	74

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

² Includes Darvocet[®], Darvon[®] or Tylenol[®] with Codeine, codeine, Phenaphen[®] with Codeine, propoxyphene, and SK-65[®].

³ Includes Percocet[®], Percodan[®] or Tylox[®], and OxyContin[®].

⁴ Includes Vicodin[®], Lortab[®], or Lorcet[®], and hydrocodone.

⁵ Includes tramadol and Ultram[®].

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.129D Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Standard Errors of Percentages, 2004 and 2005

Pain Reliever	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Darvocet [®] , Darvon [®] , or Tylenol [®] with Codeine	0.18	0.18	0.21	0.19	0.30	0.32	0.23	0.22
Percocet [®] , Percodan [®] , or Tylox [®]	0.14	0.15	0.12	0.12	0.26	0.27	0.18	0.18
Vicodin [®] , Lortab [®] , or Lorcet [®]	0.17	0.16	0.18	0.17	0.34	0.37	0.20	0.19
Codeine	0.10	0.09	0.13	0.12	0.22	0.22	0.12	0.12
Demerol [®]	0.06	0.06	0.06	0.05	0.12	0.13	0.08	0.07
Dilaudid [®]	0.04	0.04	0.01	0.02	0.04	0.07	0.05	0.06
Fioricet [®]	0.03	0.03	0.02	0.02	0.04	0.06	0.04	0.03
Fiorinal [®]	0.02	0.02	0.02	0.03	0.02	0.05	0.03	0.03
Hydrocodone	0.09	0.09	0.11	0.11	0.24	0.25	0.10	0.11
Methadone	0.04	0.05	0.05	0.06	0.10	0.11	0.05	0.06
Morphine	0.05	0.06	0.07	0.07	0.14	0.15	0.05	0.07
OxyContin [®]	0.06	0.06	0.09	0.08	0.17	0.20	0.07	0.06
Phenaphen [®] with Codeine	0.03	0.03	0.03	0.03	0.06	0.08	0.03	0.04
Propoxyphene	0.02	0.02	0.03	0.02	0.05	0.05	0.03	0.02
SK-65 [®]	0.01	0.01	0.01	0.01	0.02	0.02	0.01	0.01
Stadol [®]	0.02	0.01	0.01	*	0.03	0.02	0.02	0.01
Talacen [®]	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.01
Talwin [®]	0.03	0.02	0.02	0.02	0.02	0.02	0.04	0.03
Talwin [®] NX	0.01	0.01	0.02	0.01	0.02	0.03	0.02	0.01
Tramadol	0.02	0.02	0.02	0.03	0.05	0.06	0.02	0.02
Ultram [®]	0.04	0.03	0.04	0.03	0.08	0.09	0.05	0.04
SELECTED GROUPS OF DRUGS								
Propoxyphene or Codeine Products ^{1,2}	0.19	0.19	0.22	0.20	0.32	0.33	0.24	0.23
Oxycodone Products ^{1,3}	0.15	0.15	0.14	0.13	0.28	0.30	0.18	0.18
Hydrocodone Products ^{1,4}	0.17	0.16	0.19	0.19	0.34	0.38	0.20	0.20
Tramadol Products ^{1,5}	0.04	0.03	0.04	0.03	0.09	0.10	0.05	0.04

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

² Includes Darvocet[®], Darvon[®] or Tylenol[®] with Codeine, codeine, Phenaphen[®] with Codeine, propoxyphene, and SK-65[®].

³ Includes Percocet[®], Percodan[®] or Tylox[®], and OxyContin[®].

⁴ Includes Vicodin[®], Lortab[®], or Lorcet[®], and hydrocodone.

⁵ Includes tramadol and Ultram[®].

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.129P Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Pain Reliever	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Darvocet [®] , Darvon [®] , or Tylenol [®] with Codeine	0.3579		0.0003		0.5735		0.5082	
Percocet [®] , Percodan [®] , or Tylox [®]	0.8241		0.0277		0.1824		0.7023	
Vicodin [®] , Lortab [®] , or Lorcet [®]	0.2451		0.0764		0.0139		0.4624	
Codeine	0.3207		0.1009		0.3869		0.2754	
Demerol [®]	0.6325		0.3164		0.0115		0.8841	
Dilaudid [®]	0.4411		0.3767		0.0010		0.8307	
Fioricet [®]	0.5883		0.1110		0.0110		0.9594	
Fiorinal [®]	0.6894		0.2138		0.0308		0.8655	
Hydrocodone	0.0020		0.8740		0.0111		0.0150	
Methadone	0.0718		0.7425		0.0521		0.2256	
Morphine	0.1131		0.1051		0.0618		0.2112	
OxyContin [®]	0.0564		0.3443		0.0069		0.2927	
Phenaphen [®] with Codeine	0.1897		0.7063		0.0544		0.4375	
Propoxyphene	0.2087		0.9838		0.5609		0.1159	
SK-65 [®]	0.8056		0.3387		0.5924		0.5155	
Stadol [®]	0.1634		*		0.0953		0.3788	
Talacen [®]	0.2561		0.8196		0.4282		0.3092	
Talwin [®]	0.9074		0.2809		0.3542		0.9246	
Talwin [®] NX	0.1998		0.5395		0.3258		0.1167	
Tramadol	0.9043		0.2428		0.0271		0.2051	
Ultram [®]	0.1820		0.2673		0.0911		0.0596	
SELECTED GROUPS OF DRUGS								
Propoxyphene or Codeine Products ^{1,2}	0.5818		0.0001		0.1482		0.6515	
Oxycodone Products ^{1,3}	0.9536		0.0167		0.0719		0.7517	
Hydrocodone Products ^{1,4}	0.1018		0.1063		0.0037		0.2690	
Tramadol Products ^{1,5}	0.2262		0.3458		0.0310		0.0497	

*Low precision; no estimate reported.

¹ Includes other-specific drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

² Includes Darvocet[®], Darvon[®] or Tylenol[®] with Codeine, codeine, Phenaphen[®] with Codeine, propoxyphene, and SK-65[®].

³ Includes Percocet[®], Percodan[®] or Tylox[®], and OxyContin[®].

⁴ Includes Vicodin[®], Lortab[®], or Lorcet[®], and hydrocodone.

⁵ Includes tramadol and Ultram[®].

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.130C Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2004 and 2005

Tranquilizer	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Klonopin® or Clonazepam	137	147	20	17	53	59	122	129
Xanax®, Alprazolam, Ativan®, or Lorazepam	303	324	29	31	77	90	289	301
Valium® or Diazepam	437	420	25	24	77	81	427	402
Atarax®	40	55	5	5	8	10	39	54
BuSpar®	59	61	9	7	25	22	51	56
Equanil®	21	51	3	4	6	9	20	50
Flexeril®	134	127	7	13	35	33	131	120
Librium®	136	106	7	4	12	15	135	104
Limbitrol®	25	26	4	4	8	8	23	24
Meprobamate	38	44	6	3	6	8	37	43
Miltown®	31	34	6	4	3	8	30	32
Rohypnol®	46	50	7	7	18	18	40	46
Serax®	34	37	4	3	9	9	32	36
Soma®	151	133	15	15	48	55	140	120
Tranxene®	37	26	4	3	8	7	36	25
Vistaril®	54	46	8	6	10	13	52	44
SELECTED GROUPS OF DRUGS								
Benzodiazepines ^{1,2}	474	466	36	35	93	105	455	445
Meprobamate Products ^{1,3}	51	61	8	6	9	11	49	60
Muscle Relaxants ^{1,4}	186	176	15	19	55	58	177	164

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

² Includes Klonopin® or clonazepam, Xanax®, alprazolam, Ativan®, or lorazepam, Valium® or diazepam, Librium®, Limbitrol®, Rohypnol®, Serax®, and Tranxene®.

³ Includes Equanil®, meprobamate, and Miltown®.

⁴ Includes Flexeril® and Soma®.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.130D Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Standard Errors of Percentages, 2004 and 2005

Tranquilizer	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Klonopin® or Clonazepam	0.06	0.06	0.08	0.07	0.16	0.18	0.07	0.07
Xanax®, Alprazolam, Ativan®, or Lorazepam	0.13	0.13	0.12	0.12	0.24	0.28	0.16	0.16
Valium® or Diazepam	0.18	0.17	0.10	0.09	0.24	0.25	0.23	0.22
Atarax®	0.02	0.02	0.02	0.02	0.02	0.03	0.02	0.03
BuSpar®	0.02	0.03	0.03	0.03	0.08	0.07	0.03	0.03
Equanil®	0.01	0.02	0.01	0.01	0.02	0.03	0.01	0.03
Flexeril®	0.06	0.05	0.03	0.05	0.11	0.10	0.07	0.06
Librium®	0.06	0.04	0.03	0.02	0.04	0.05	0.07	0.06
Limbitrol®	0.01	0.01	0.02	0.02	0.02	0.03	0.01	0.01
Meprobamate	0.02	0.02	0.02	0.01	0.02	0.02	0.02	0.02
Miltown®	0.01	0.01	0.02	0.01	0.01	0.02	0.02	0.02
Rohypnol®	0.02	0.02	0.03	0.03	0.06	0.06	0.02	0.02
Serax®	0.01	0.02	0.02	0.01	0.03	0.03	0.02	0.02
Soma®	0.06	0.05	0.06	0.06	0.15	0.17	0.08	0.06
Tranxene®	0.02	0.01	0.01	0.01	0.03	0.02	0.02	0.01
Vistaril®	0.02	0.02	0.03	0.03	0.03	0.04	0.03	0.02
SELECTED GROUPS OF DRUGS								
Benzodiazepines ^{1,2}	0.20	0.19	0.14	0.14	0.29	0.32	0.25	0.24
Meprobamate Products ^{1,3}	0.02	0.03	0.03	0.02	0.03	0.04	0.03	0.03
Muscle Relaxants ^{1,4}	0.08	0.07	0.06	0.07	0.17	0.18	0.10	0.09

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

² Includes Klonopin® or clonazepam, Xanax®, alprazolam, Ativan®, or lorazepam, Valium® or diazepam, Librium®, Limbitrol®, Rohypnol®, Serax®, and Tranxene®.

³ Includes Equanil®, meprobamate, and Miltown®.

⁴ Includes Flexeril® and Soma®.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.130P Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Tranquilizer	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Klonopin® or Clonazepam	0.0806		0.4733		0.0195		0.2946	
Xanax®, Alprazolam, Ativan®, or Lorazepam	0.0676		0.8750		0.0000		0.4730	
Valium® or Diazepam	0.8155		0.3876		0.0853		0.9623	
Atarax®	0.7502		0.5336		0.7262		0.7439	
BuSpar®	0.3173		0.2680		0.2422		0.6626	
Equanil®	0.1304		0.5126		0.6581		0.1288	
Flexeril®	0.5912		0.0447		0.7779		0.4230	
Librium®	0.3134		0.0247		0.1657		0.2838	
Limbitrol®	0.9141		0.7587		0.7138		0.7671	
Meprobamate	0.9947		0.3254		0.5765		0.9783	
Miltown®	0.3663		0.2071		0.1881		0.3712	
Rohypnol®	0.5251		0.9502		0.7783		0.5699	
Serax®	0.9297		0.7779		0.6687		0.9524	
Soma®	0.5525		0.3182		0.3224		0.3579	
Tranxene®	0.3832		0.4955		0.5545		0.4925	
Vistaril®	0.6197		0.8497		0.8488		0.5934	
SELECTED GROUPS OF DRUGS								
Benzodiazepines ^{1,2}	0.2135		0.2848		0.0012		0.5087	
Meprobamate Products ^{1,3}	0.4888		0.2136		0.5072		0.4536	
Muscle Relaxants ^{1,4}	0.4906		0.9117		0.8042		0.4110	

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

² Includes Klonopin® or clonazepam, Xanax®, alprazolam, Ativan®, or lorazepam, Valium® or diazepam, Librium®, Limbitrol®, Rohypnol®, Serax®, and Tranxene®.

³ Includes Equanil®, meprobamate, and Miltown®.

⁴ Includes Flexeril® and Soma®.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.131C Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2004 and 2005

Stimulant	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Methamphetamine, ¹ Desoxyn [®] , or Methedrine	352	345	24	26	61	68	349	334
Prescription Diet Pills ²	321	306	21	21	37	41	317	301
Ritalin [®] or Methylphenidate	191	194	27	28	71	69	167	173
Cylert [®]	47	33	8	4	8	11	45	31
Dexedrine [®]	179	190	10	8	29	26	176	189
Dextroamphetamine	132	49	7	8	13	19	131	44
Didrex [®]	39	44	5	4	12	8	37	43
Eskatrol [®]	78	26	5	8	5	3	78	25
Ionamin [®]	73	57	4	2	4	9	72	56
Mazanor [®]	26	10	6	4	3	5	25	8
Obedrin-LA [®]	*	21	*	2	*	*	*	21
Plegine [®]	7	16	4	2	2	1	6	16
Preludin [®]	88	59	5	3	8	11	87	58
Sanorex [®]	21	18	6	9	8	11	18	12
Tenuate [®]	94	29	4	3	5	4	94	28
SELECTED GROUPS OF DRUGS								
Amphetamine, Dextroamphetamine, and Phentermine Products ^{3,4}	347	324	25	26	53	58	336	318
Mazindol Products ^{3,5}	33	19	8	10	8	12	31	12

*Low precision; no estimate reported.

¹ Also known as crank, crystal, ice, or speed.

² Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine[®], Biphetamine[®], Fastin[®], or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

³ Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

⁴ Includes prescription diet pills, Dexedrine[®], dextroamphetamine, and Ionamin[®].

⁵ Includes Mazanor[®] and Sanorex[®].

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.131D Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Standard Errors of Percentages, 2004 and 2005

Stimulant	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Methamphetamine, ¹ Desoxyn [®] , or Methedrine	0.15	0.14	0.09	0.10	0.19	0.21	0.19	0.18
Prescription Diet Pills ²	0.13	0.13	0.08	0.08	0.12	0.13	0.17	0.16
Ritalin [®] or Methylphenidate	0.08	0.08	0.11	0.11	0.22	0.21	0.09	0.09
Cylert [®]	0.02	0.01	0.03	0.02	0.03	0.03	0.02	0.02
Dexedrine [®]	0.07	0.08	0.04	0.03	0.09	0.08	0.10	0.10
Dextroamphetamine	0.06	0.02	0.03	0.03	0.04	0.06	0.07	0.02
Didrex [®]	0.02	0.02	0.02	0.02	0.04	0.03	0.02	0.02
Eskatrol [®]	0.03	0.01	0.02	0.03	0.02	0.01	0.04	0.01
Ionamin [®]	0.03	0.02	0.02	0.01	0.01	0.03	0.04	0.03
Mazanor [®]	0.01	0.00	0.02	0.02	0.01	0.01	0.01	0.00
Obedrin-LA [®]	*	0.01	*	0.01	*	*	*	0.01
Plegine [®]	0.00	0.01	0.01	0.01	0.01	0.00	0.00	0.01
Preludin [®]	0.04	0.02	0.02	0.01	0.02	0.03	0.05	0.03
Sanorex [®]	0.01	0.01	0.02	0.04	0.03	0.03	0.01	0.01
Tenuate [®]	0.04	0.01	0.02	0.01	0.01	0.01	0.05	0.02
SELECTED GROUPS OF DRUGS								
Amphetamine, Dextroamphetamine, and Phentermine Products ^{3,4}	0.14	0.13	0.10	0.10	0.17	0.18	0.18	0.17
Mazindol Products ^{3,5}	0.01	0.01	0.03	0.04	0.03	0.04	0.02	0.01

*Low precision; no estimate reported.

¹ Also known as crank, crystal, ice, or speed.

² Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine[®], Biphetamine[®], Fastin[®], or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

³ Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

⁴ Includes prescription diet pills, Dexedrine[®], dextroamphetamine, and Ionamin[®].

⁵ Includes Mazanor[®] and Sanorex[®].

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.131P Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Stimulant	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Methamphetamine, ¹ Desoxyn [®] , or Methedrine	0.0025		0.8985		0.8168		0.0024	
Prescription Diet Pills ²	0.1105		0.3588		0.3318		0.0907	
Ritalin [®] or Methylphenidate	0.2534		0.7579		0.1924		0.4091	
Cylert [®]	0.0921		0.2105		0.5452		0.0847	
Dexedrine [®]	0.4493		0.0292		0.1345		0.6594	
Dextroamphetamine	0.0313		0.8697		0.0033		0.0071	
Didrex [®]	0.9283		0.4806		0.5881		0.8995	
Eskatrol [®]	0.0156		0.8795		0.1396		0.0206	
Ionamin [®]	0.0660		0.0565		0.4712		0.0664	
Mazanor [®]	0.2094		0.5007		0.8188		0.2307	
Obedrin-LA [®]	*		*		*		*	
Plegine [®]	0.6317		0.0215		0.9085		0.2951	
Preludin [®]	0.1416		0.3545		0.2355		0.1118	
Sanorex [®]	0.3805		0.7506		0.7772		0.2580	
Tenuate [®]	0.1915		0.2862		0.2258		0.2375	
SELECTED GROUPS OF DRUGS								
Amphetamine, Dextroamphetamine, and Phentermine Products ^{3,4}	0.1554		0.3358		0.0779		0.0910	
Mazindol Products ^{3,5}	0.1571		0.7960		0.4417		0.0639	

*Low precision; no estimate reported.

¹ Also known as crank, crystal, ice, or speed.

² Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine[®], Biphetamine[®], Fastin[®], or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

³ Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

⁴ Includes prescription diet pills, Dexedrine[®], dextroamphetamine, and Ionamin[®].

⁵ Includes Mazanor[®] and Sanorex[®].

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.132C Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2004 and 2005

Sedative	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Methaqualone, Sopor [®] , or Quaalude [®]	349	284	10	12	28	24	347	282
Barbiturates ¹	174	184	8	7	15	17	173	182
Restoril [®] or Temazepam	84	122	9	11	15	20	81	120
Amytal [®]	59	60	7	3	6	9	58	59
Butisol [®]	18	37	6	3	7	9	16	35
Chloral Hydrate	50	73	6	8	4	11	49	72
Dalmane [®]	75	86	8	7	5	10	75	85
Halcion [®]	84	83	10	4	11	12	82	82
Phenobarbital	132	107	8	8	12	14	131	106
Placidyl [®]	106	97	3	1	5	8	106	96
Tuinal [®]	122	122	4	3	5	6	121	121
SELECTED GROUPS OF DRUGS								
Temazepam, Flurazepam, or Triazolam ^{2,3}	121	144	14	14	19	24	118	142
Any Barbiturates ^{2,4}	207	205	11	11	19	23	206	202

*Low precision; no estimate reported.

¹ Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal[®], pentobarbital, Seconal[®], secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

² Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

³ Includes Restoril[®], temazepam, Dalmane[®], or Halcion[®].

⁴ Includes barbiturates, Amytal[®], Butisol[®], phenobarbital, and Tuinal[®].

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.132D Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Standard Errors of Percentages, 2004 and 2005

Sedative	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Methaqualone, Sopor [®] , or Quaalude [®]	0.15	0.12	0.04	0.05	0.09	0.07	0.19	0.15
Barbiturates ¹	0.07	0.08	0.03	0.03	0.05	0.05	0.09	0.10
Restoril [®] or Temazepam	0.03	0.05	0.04	0.04	0.05	0.06	0.04	0.06
Amytal [®]	0.02	0.02	0.03	0.01	0.02	0.03	0.03	0.03
Butisol [®]	0.01	0.02	0.02	0.01	0.02	0.03	0.01	0.02
Chloral Hydrate	0.02	0.03	0.03	0.03	0.01	0.03	0.03	0.04
Dalmane [®]	0.03	0.04	0.03	0.03	0.02	0.03	0.04	0.05
Halcion [®]	0.03	0.03	0.04	0.02	0.03	0.04	0.04	0.04
Phenobarbital	0.05	0.04	0.03	0.03	0.04	0.04	0.07	0.06
Placidyl [®]	0.04	0.04	0.01	0.01	0.01	0.03	0.06	0.05
Tuinal [®]	0.05	0.05	0.01	0.01	0.01	0.02	0.07	0.07
SELECTED GROUPS OF DRUGS								
Temazepam, Flurazepam, or Triazolam ^{2,3}	0.05	0.06	0.06	0.05	0.06	0.07	0.06	0.08
Any Barbiturates ^{2,4}	0.09	0.08	0.05	0.04	0.06	0.07	0.11	0.11

*Low precision; no estimate reported.

¹ Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal[®], pentobarbital, Seconal[®], secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

² Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

³ Includes Restoril[®], temazepam, Dalmane[®], or Halcion[®].

⁴ Includes barbiturates, Amytal[®], Butisol[®], phenobarbital, and Tuinal[®].

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.132P Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2005 Versus 2004

Sedative	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Methaqualone, Sopor [®] , or Quaalude [®]	0.0072		0.8092		0.5423		0.0074	
Barbiturates ¹	0.6719		0.4539		0.9523		0.6897	
Restoril [®] or Temazepam	0.1008		0.7347		0.1556		0.1635	
Amytal [®]	0.5957		0.0580		0.5987		0.6595	
Butisol [®]	0.1642		0.1891		0.8822		0.0992	
Chloral Hydrate	0.5265		0.7583		0.3036		0.5906	
Dalmane [®]	0.7852		0.2286		0.3317		0.7990	
Halcion [®]	0.1893		0.0785		0.7796		0.2280	
Phenobarbital	0.3180		0.8678		0.5444		0.2757	
Placidyl [®]	0.1676		0.6251		0.6005		0.1576	
Tuinal [®]	0.6068		0.7189		0.7094		0.5972	
SELECTED GROUPS OF DRUGS								
Temazepam, Flurazepam, or Triazolam ^{2,3}	0.7479		0.4354		0.1709		0.8530	
Any Barbiturates ^{2,4}	0.5697		0.1943		0.5655		0.5684	

*Low precision; no estimate reported.

¹ Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal[®], pentobarbital, Seconal[®], secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

² Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

³ Includes Restoril[®], temazepam, Dalmane[®], or Halcion[®].

⁴ Includes barbiturates, Amytal[®], Butisol[®], phenobarbital, and Tuinal[®].

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.