

Table 1.19C Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	969	940	535	550	415	432
12	29	32	22	25	14	13
13	41	38	30	32	19	19
14	50	44	42	35	30	21
15	54	51	46	43	34	30
16	59	58	48	51	35	37
17	67	63	56	51	37	35
18	85	83	69	67	50	53
19	82	90	69	71	51	54
20	84	88	69	70	49	54
21	92	91	70	71	55	47
22	85	85	64	60	51	45
23	76	84	51	59	38	44
24	75	83	50	52	40	41
25	79	83	54	54	38	44
26-29	327	319	188	188	137	145
30-34	314	326	159	200	132	148
35-39	392	360	172	160	113	122
40-44	395	410	198	172	140	127
45-49	371	402	167	158	134	121
50-54	513	473	210	180	142	151
55-59	374	386	119	141	79	115
60-64	269	280	73	105	54	79
65 or Older	272	286	83	147	46	86

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.19D Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.40	0.39	0.22	0.23	0.17	0.18
12	0.68	0.72	0.53	0.58	0.36	0.32
13	0.86	0.81	0.65	0.70	0.44	0.44
14	0.92	0.92	0.81	0.77	0.62	0.47
15	1.01	0.96	0.91	0.87	0.74	0.66
16	1.04	1.05	0.95	0.96	0.77	0.78
17	1.13	1.10	1.08	1.04	0.83	0.79
18	1.17	1.25	1.11	1.13	0.92	0.99
19	1.30	1.25	1.31	1.21	1.07	1.06
20	1.27	1.26	1.27	1.28	1.05	1.13
21	1.21	1.23	1.35	1.24	1.15	0.98
22	1.18	1.25	1.24	1.19	1.10	0.98
23	1.23	1.26	1.08	1.20	0.88	1.00
24	1.29	1.27	1.03	1.13	0.90	0.95
25	1.34	1.31	1.18	1.16	0.93	0.99
26-29	1.18	1.23	1.05	1.07	0.81	0.87
30-34	1.06	1.19	0.73	0.91	0.63	0.72
35-39	1.13	1.14	0.78	0.75	0.52	0.59
40-44	1.04	1.07	0.83	0.71	0.60	0.54
45-49	1.06	1.07	0.73	0.68	0.60	0.52
50-54	1.76	1.71	1.02	0.87	0.71	0.73
55-59	1.70	1.70	0.69	0.80	0.47	0.66
60-64	1.82	1.82	0.58	0.78	0.43	0.59
65 or Older	0.79	0.81	0.24	0.42	0.13	0.24

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.19P Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.5606		0.8381		0.4711	
12	0.5564		0.4246		0.5772	
13	0.2015		0.8646		0.6321	
14	0.0433		0.0352		0.0023	
15	0.0019		0.0361		0.0893	
16	0.0030		0.2207		0.7892	
17	0.1084		0.3437		0.4014	
18	0.3808		0.1512		0.5839	
19	0.8507		0.9198		0.8973	
20	0.6676		0.7994		0.0708	
21	0.1414		0.6935		0.5117	
22	0.4448		0.5094		0.7311	
23	0.0219		0.0012		0.0024	
24	0.5719		0.6230		0.7584	
25	0.9623		0.9875		0.3569	
26-29	0.9656		0.6494		0.7756	
30-34	0.8116		0.1069		0.8221	
35-39	0.4691		0.6171		0.5539	
40-44	0.0435		0.1932		0.7356	
45-49	0.2749		0.9803		0.8390	
50-54	0.7683		0.5948		0.7035	
55-59	0.0141		0.6167		0.2988	
60-64	0.1202		0.2244		0.3404	
65 or Older	0.0239		0.1022		0.1044	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.20C Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	912	925	451	473	360	360
12	12	8	9	7	5	4
13	20	21	18	18	12	11
14	34	29	31	25	21	17
15	42	41	37	36	30	26
16	53	50	44	45	31	32
17	57	55	49	46	34	33
18	77	76	62	61	44	48
19	77	84	64	65	46	49
20	77	81	61	63	46	50
21	87	85	62	63	51	42
22	82	79	59	55	46	42
23	72	78	46	52	34	40
24	70	79	45	48	36	36
25	76	77	49	47	35	38
26-29	300	296	163	166	123	128
30-34	299	306	127	166	111	131
35-39	375	338	143	128	96	98
40-44	379	390	158	145	119	99
45-49	359	384	144	126	116	102
50-54	489	459	186	146	135	125
55-59	352	361	88	119	63	97
60-64	238	253	31	86	16	48
65 or Older	240	243	42	115	30	65

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.20D Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.38	0.38	0.19	0.19	0.15	0.15
12	0.29	0.21	0.23	0.19	0.13	0.10
13	0.46	0.48	0.41	0.41	0.28	0.26
14	0.70	0.67	0.65	0.57	0.44	0.40
15	0.86	0.86	0.76	0.77	0.64	0.57
16	1.00	0.97	0.90	0.90	0.70	0.69
17	1.11	1.08	1.00	0.98	0.77	0.75
18	1.16	1.22	1.06	1.07	0.84	0.94
19	1.34	1.25	1.29	1.19	1.02	1.00
20	1.31	1.31	1.20	1.23	1.00	1.08
21	1.26	1.32	1.23	1.20	1.08	0.91
22	1.28	1.28	1.19	1.11	1.01	0.91
23	1.26	1.25	1.01	1.11	0.79	0.93
24	1.24	1.30	0.98	1.06	0.84	0.86
25	1.34	1.32	1.12	1.05	0.88	0.88
26-29	1.24	1.30	0.93	0.96	0.74	0.77
30-34	1.08	1.20	0.62	0.80	0.55	0.65
35-39	1.12	1.14	0.67	0.62	0.45	0.48
40-44	1.06	1.10	0.67	0.61	0.51	0.42
45-49	1.06	1.07	0.64	0.55	0.53	0.45
50-54	1.70	1.72	0.92	0.72	0.68	0.62
55-59	1.68	1.64	0.53	0.68	0.38	0.56
60-64	1.67	1.70	0.24	0.65	0.12	0.36
65 or Older	0.69	0.69	0.12	0.33	0.09	0.18

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.20P Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.8089		0.5807		0.8247	
12	0.4100		0.8329		0.4235	
13	0.8368		0.6691		0.4528	
14	0.0082		0.0074		0.0512	
15	0.0810		0.2861		0.0516	
16	0.0054		0.0774		0.4711	
17	0.1995		0.1878		0.3462	
18	0.8718		0.8281		0.8458	
19	0.7550		0.5971		0.8666	
20	0.5124		0.7122		0.1449	
21	0.0715		0.7492		0.2678	
22	0.3805		0.7040		0.9000	
23	0.0091		0.0036		0.0280	
24	0.5771		0.9426		0.7843	
25	0.6628		0.9474		0.6755	
26-29	0.9201		0.7236		0.4301	
30-34	0.3838		0.0605		0.1858	
35-39	0.2775		0.1339		0.6543	
40-44	0.0229		0.3183		0.1292	
45-49	0.2616		0.8124		0.6324	
50-54	0.4349		0.3554		0.9378	
55-59	0.0255		0.4374		0.2292	
60-64	0.1799		0.0761		0.0923	
65 or Older	0.1277		0.1875		0.3412	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.21C Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	636	595	205	205	119	142
12	2	3	2	3	*	0
13	5	5	5	5	3	1
14	11	8	6	7	3	6
15	13	11	11	10	7	6
16	18	19	15	17	6	9
17	23	23	18	19	11	11
18	31	34	24	24	14	14
19	37	38	28	25	14	15
20	33	42	25	31	13	18
21	45	41	33	28	18	16
22	45	42	28	30	14	19
23	39	42	24	27	14	15
24	40	42	23	24	14	14
25	40	43	20	26	11	17
26-29	147	143	86	68	43	44
30-34	166	176	76	67	48	35
35-39	246	206	66	59	45	44
40-44	269	281	88	70	47	55
45-49	250	264	55	74	38	51
50-54	280	257	54	67	28	57
55-59	157	191	25	44	23	21
60-64	80	115	23	45	22	*
65 or Older	94	79	16	15	12	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.21D Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.26	0.24	0.09	0.08	0.05	0.06
12	0.06	0.07	0.05	0.07	*	0.01
13	0.12	0.11	0.11	0.11	0.08	0.04
14	0.24	0.18	0.14	0.16	0.07	0.13
15	0.30	0.25	0.24	0.23	0.17	0.14
16	0.42	0.44	0.36	0.38	0.13	0.21
17	0.53	0.56	0.44	0.44	0.27	0.26
18	0.65	0.72	0.52	0.51	0.31	0.31
19	0.90	0.82	0.69	0.55	0.35	0.35
20	0.77	0.94	0.61	0.70	0.34	0.43
21	0.95	0.90	0.76	0.63	0.44	0.39
22	0.97	0.91	0.66	0.69	0.34	0.45
23	0.92	0.97	0.58	0.66	0.36	0.38
24	0.89	0.96	0.54	0.61	0.34	0.38
25	1.00	0.97	0.52	0.63	0.30	0.43
26-29	0.90	0.85	0.54	0.43	0.28	0.28
30-34	0.78	0.83	0.38	0.34	0.24	0.18
35-39	0.99	0.92	0.31	0.29	0.22	0.22
40-44	1.00	1.00	0.38	0.30	0.21	0.23
45-49	1.03	0.98	0.26	0.33	0.18	0.23
50-54	1.26	1.20	0.28	0.34	0.15	0.29
55-59	0.91	1.05	0.15	0.26	0.14	0.12
60-64	0.62	0.86	0.18	0.34	0.18	*
65 or Older	0.27	0.22	0.05	0.04	0.03	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.21P Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.3244		0.4945		0.0585	
12	0.9230		0.6065		*	
13	0.4311		0.6410		0.5045	
14	0.2786		0.5994		0.1435	
15	0.0875		0.5404		0.6629	
16	0.7517		0.7938		0.2731	
17	0.5937		0.5499		0.9668	
18	0.9158		0.9331		0.5814	
19	0.9517		0.9540		0.8909	
20	0.8032		0.5379		0.5447	
21	0.4420		0.0996		0.6054	
22	0.1432		0.6898		0.1367	
23	0.4220		0.0842		0.3458	
24	0.7053		0.2683		0.2424	
25	0.6694		0.0108		0.0306	
26-29	0.9237		0.0829		0.3660	
30-34	0.5442		0.2179		0.1204	
35-39	0.0115		0.5777		0.8081	
40-44	0.1197		0.1241		0.5238	
45-49	0.0484		0.1552		0.1333	
50-54	0.1645		0.4931		0.1898	
55-59	0.0503		0.2673		0.9816	
60-64	0.2524		0.7379		*	
65 or Older	0.4692		0.7317		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.22C Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	280	292	117	114	66	87
12	1	*	1	*	*	*
13	3	2	3	1	3	1
14	5	4	4	4	3	3
15	6	3	3	3	1	1
16	9	7	7	6	1	4
17	9	10	5	6	3	5
18	14	17	11	10	3	7
19	18	18	9	14	6	8
20	14	20	5	9	4	6
21	20	16	12	6	7	3
22	23	15	7	10	3	7
23	18	23	8	14	5	4
24	18	20	7	7	4	4
25	19	19	4	8	3	3
26-29	77	74	41	31	18	23
30-34	87	95	38	28	26	19
35-39	124	108	42	42	24	30
40-44	128	139	43	47	26	38
45-49	108	125	28	43	22	30
50-54	107	115	50	28	24	21
55-59	63	74	14	20	14	15
60-64	22	46	7	*	*	*
65 or Older	54	15	16	15	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.22D Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.12	0.12	0.05	0.05	0.03	0.04
12	0.03	*	0.03	*	*	*
13	0.07	0.05	0.07	0.04	0.07	0.03
14	0.11	0.10	0.08	0.08	0.06	0.08
15	0.14	0.08	0.08	0.06	0.03	0.02
16	0.22	0.16	0.17	0.13	0.03	0.10
17	0.21	0.25	0.12	0.15	0.07	0.11
18	0.31	0.38	0.23	0.22	0.06	0.16
19	0.44	0.42	0.23	0.32	0.14	0.18
20	0.34	0.47	0.13	0.23	0.11	0.15
21	0.47	0.39	0.29	0.15	0.18	0.07
22	0.55	0.38	0.18	0.25	0.06	0.18
23	0.45	0.59	0.21	0.37	0.13	0.11
24	0.43	0.52	0.18	0.19	0.09	0.10
25	0.54	0.48	0.12	0.22	0.09	0.07
26-29	0.49	0.47	0.26	0.20	0.11	0.15
30-34	0.44	0.47	0.19	0.14	0.13	0.10
35-39	0.58	0.53	0.20	0.21	0.11	0.15
40-44	0.56	0.57	0.19	0.20	0.11	0.16
45-49	0.49	0.55	0.13	0.19	0.10	0.14
50-54	0.54	0.57	0.26	0.14	0.13	0.11
55-59	0.38	0.43	0.09	0.12	0.09	0.09
60-64	0.17	0.35	0.05	*	*	*
65 or Older	0.16	0.04	0.05	0.04	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.22P Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	1.0000		0.7040		0.0560	
12	*		*		*	
13	0.9570		0.7573		0.5812	
14	0.3522		0.6202		0.9752	
15	0.0123		0.1512		0.6930	
16	0.7418		0.6405		0.1590	
17	0.5539		0.5650		0.2909	
18	0.3431		0.9152		0.0943	
19	0.8827		0.1868		0.5561	
20	0.3683		0.2494		0.8374	
21	0.0881		0.0222		0.0465	
22	0.0340		0.1930		0.0209	
23	0.1326		0.2582		0.8877	
24	0.4285		0.7243		0.6541	
25	1.0000		0.0439		0.7685	
26-29	0.9386		0.6652		0.2653	
30-34	0.8656		0.7241		0.6194	
35-39	0.2369		0.8751		0.3202	
40-44	0.7278		0.9331		0.3797	
45-49	0.3640		0.2321		0.3495	
50-54	0.4500		0.3094		0.9276	
55-59	0.6987		0.4035		0.7245	
60-64	0.7609		*		*	
65 or Older	0.0985		0.7317		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.23C Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	618	608	153	146	58	83
12	6	5	4	3	3	2
13	12	11	10	8	5	6
14	16	11	14	10	7	4
15	19	18	16	15	5	7
16	23	23	20	19	11	10
17	28	28	21	23	12	12
18	38	35	27	27	13	13
19	39	46	27	31	12	15
20	42	51	25	30	12	17
21	54	50	31	27	14	17
22	55	47	24	26	10	10
23	47	50	22	22	11	11
24	47	51	19	19	9	6
25	51	52	18	18	11	8
26-29	191	176	70	63	30	36
30-34	176	195	54	47	20	13
35-39	214	196	55	32	20	22
40-44	228	216	39	32	14	9
45-49	220	216	25	14	6	9
50-54	299	262	5	52	*	45
55-59	154	195	18	*	*	*
60-64	72	116	*	37	*	35
65 or Older	85	54	24	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.23D Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.26	0.25	0.06	0.06	0.02	0.03
12	0.15	0.12	0.11	0.07	0.08	0.04
13	0.26	0.25	0.23	0.20	0.12	0.13
14	0.36	0.25	0.31	0.22	0.16	0.09
15	0.42	0.41	0.36	0.33	0.13	0.16
16	0.53	0.52	0.47	0.44	0.26	0.24
17	0.65	0.65	0.50	0.54	0.30	0.28
18	0.79	0.71	0.57	0.56	0.28	0.28
19	0.90	0.96	0.63	0.67	0.31	0.35
20	0.94	1.11	0.60	0.70	0.31	0.41
21	1.11	1.02	0.70	0.63	0.33	0.41
22	1.15	1.03	0.57	0.63	0.24	0.24
23	1.06	1.08	0.55	0.55	0.27	0.27
24	1.01	1.10	0.46	0.49	0.23	0.17
25	1.18	1.13	0.49	0.47	0.29	0.19
26-29	1.09	0.95	0.44	0.40	0.20	0.23
30-34	0.81	0.91	0.27	0.24	0.10	0.07
35-39	0.91	0.88	0.27	0.16	0.10	0.11
40-44	0.89	0.83	0.17	0.14	0.06	0.04
45-49	0.91	0.88	0.11	0.06	0.03	0.04
50-54	1.36	1.22	0.03	0.26	*	0.23
55-59	0.90	1.07	0.11	*	*	*
60-64	0.56	0.87	*	0.28	*	0.26
65 or Older	0.25	0.15	0.07	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.23P Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.2560		0.5978		0.1483	
12	0.7935		0.2968		0.1040	
13	0.5982		0.2158		0.8126	
14	0.0135		0.1162		0.3333	
15	0.1467		0.1077		0.7429	
16	0.4549		0.4944		0.9396	
17	0.1023		0.9916		0.9327	
18	0.1088		0.7893		0.9246	
19	0.2006		0.3505		0.9867	
20	0.4358		0.7346		0.2279	
21	0.4405		0.7306		0.3840	
22	0.0094		0.9461		0.3650	
23	0.3001		0.2277		1.0000	
24	0.4989		0.5085		0.1363	
25	0.9486		0.8106		0.8280	
26-29	0.6800		0.4403		0.6236	
30-34	0.8457		0.5860		0.7340	
35-39	0.6801		0.3383		0.5765	
40-44	0.0203		0.6779		0.4953	
45-49	0.8164		0.8056		0.3965	
50-54	0.1051		0.0416		*	
55-59	0.0065		*		*	
60-64	0.0210		*		*	
65 or Older	0.3381		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.24C Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	480	460	90	95	64	57
12	24	26	15	19	9	9
13	30	26	19	18	10	11
14	35	33	24	21	10	8
15	33	32	22	20	13	9
16	31	28	16	19	7	12
17	33	35	23	16	10	7
18	34	35	19	21	9	11
19	34	38	16	19	5	10
20	35	37	16	15	5	8
21	42	37	17	16	4	6
22	43	39	14	16	10	8
23	37	39	10	10	7	4
24	41	37	11	9	4	4
25	36	38	8	7	5	5
26-29	130	151	19	26	10	6
30-34	144	173	29	26	24	21
35-39	171	162	25	27	18	26
40-44	192	178	25	27	13	11
45-49	148	164	16	18	14	1
50-54	205	176	21	*	*	*
55-59	120	132	*	*	*	*
60-64	70	82	40	*	40	*
65 or Older	83	71	*	33	*	33

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.24D Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.20	0.19	0.04	0.04	0.03	0.02
12	0.57	0.59	0.38	0.45	0.23	0.22
13	0.64	0.58	0.42	0.42	0.22	0.25
14	0.72	0.71	0.52	0.47	0.23	0.18
15	0.71	0.69	0.49	0.46	0.31	0.21
16	0.68	0.61	0.38	0.43	0.16	0.28
17	0.73	0.77	0.53	0.39	0.23	0.18
18	0.71	0.72	0.42	0.46	0.20	0.25
19	0.80	0.83	0.40	0.44	0.13	0.23
20	0.82	0.83	0.39	0.37	0.13	0.20
21	0.92	0.83	0.41	0.38	0.11	0.15
22	0.98	0.86	0.35	0.38	0.24	0.19
23	0.88	0.89	0.25	0.27	0.17	0.09
24	0.90	0.90	0.28	0.23	0.09	0.10
25	0.91	0.88	0.21	0.19	0.14	0.13
26-29	0.81	0.86	0.12	0.17	0.07	0.04
30-34	0.68	0.83	0.15	0.13	0.12	0.11
35-39	0.77	0.76	0.12	0.13	0.08	0.13
40-44	0.79	0.69	0.11	0.12	0.06	0.05
45-49	0.67	0.69	0.08	0.08	0.06	0.01
50-54	1.00	0.86	0.11	*	*	*
55-59	0.70	0.75	*	*	*	*
60-64	0.55	0.61	0.32	*	0.32	*
65 or Older	0.24	0.20	*	0.09	*	0.09

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.24P Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.6435		0.4770		0.6944	
12	0.7106		0.4413		0.7607	
13	0.2826		0.4697		0.0636	
14	0.6632		0.5249		0.3111	
15	0.1226		0.4125		0.2857	
16	0.3886		0.8849		0.4170	
17	0.5533		0.3247		0.7512	
18	0.2439		0.7544		0.1828	
19	0.6819		0.5935		0.2690	
20	0.7763		0.9364		0.5420	
21	0.0702		0.5246		0.4398	
22	0.3553		0.5407		0.9397	
23	0.6261		0.5943		0.1244	
24	0.7176		0.4926		0.8521	
25	0.4754		0.9107		0.4482	
26-29	0.3934		0.5199		0.7077	
30-34	0.3400		0.3565		0.7115	
35-39	0.4840		0.7730		0.9344	
40-44	0.4539		0.8163		0.7557	
45-49	0.4793		0.9526		0.3784	
50-54	0.5250		*		*	
55-59	0.7738		*		*	
60-64	0.2389		*		*	
65 or Older	0.8298		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.25C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	729	712	353	342	210	229
12	17	20	14	16	10	11
13	27	25	22	20	14	13
14	32	25	26	22	19	14
15	35	32	27	26	18	16
16	38	37	31	31	19	19
17	42	41	34	34	20	21
18	59	53	43	42	27	26
19	55	57	42	45	28	32
20	61	64	45	48	23	28
21	62	62	45	43	29	27
22	60	57	42	39	26	24
23	54	60	36	36	22	25
24	50	57	32	34	21	21
25	48	57	30	34	17	22
26-29	201	195	119	117	77	71
30-34	191	210	114	123	77	79
35-39	237	222	116	114	66	74
40-44	251	249	120	121	71	78
45-49	245	271	103	103	62	73
50-54	347	312	102	117	52	91
55-59	212	249	84	96	52	58
60-64	145	183	53	58	25	40
65 or Older	162	173	68	86	33	46

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.25D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.30	0.29	0.15	0.14	0.09	0.09
12	0.44	0.48	0.36	0.40	0.24	0.26
13	0.59	0.58	0.49	0.46	0.32	0.30
14	0.65	0.54	0.55	0.50	0.42	0.32
15	0.76	0.67	0.59	0.57	0.40	0.35
16	0.81	0.77	0.67	0.68	0.44	0.44
17	0.92	0.91	0.79	0.76	0.48	0.49
18	1.05	1.00	0.85	0.83	0.57	0.53
19	1.13	1.10	0.96	0.92	0.67	0.71
20	1.21	1.22	1.00	1.04	0.55	0.64
21	1.23	1.14	1.00	0.94	0.69	0.64
22	1.17	1.15	0.92	0.90	0.63	0.58
23	1.13	1.22	0.84	0.85	0.54	0.61
24	1.02	1.20	0.73	0.82	0.50	0.56
25	1.10	1.18	0.78	0.80	0.47	0.54
26-29	1.08	1.03	0.73	0.72	0.49	0.45
30-34	0.90	0.93	0.56	0.60	0.38	0.40
35-39	0.95	0.97	0.54	0.54	0.31	0.36
40-44	0.99	0.93	0.53	0.51	0.31	0.33
45-49	0.98	1.05	0.47	0.46	0.29	0.32
50-54	1.51	1.44	0.52	0.58	0.27	0.45
55-59	1.15	1.31	0.50	0.55	0.31	0.34
60-64	1.09	1.32	0.42	0.44	0.20	0.30
65 or Older	0.47	0.49	0.20	0.24	0.10	0.13

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.25P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.8785		0.4599		0.2904	
12	0.5008		0.8531		0.5051	
13	0.0835		0.0942		0.5644	
14	0.0028		0.0688		0.0025	
15	0.0016		0.0890		0.0776	
16	0.0317		0.5776		0.4580	
17	0.8700		0.3819		0.7838	
18	0.0342		0.0456		0.1009	
19	0.7199		0.4063		0.8021	
20	0.0155		0.2144		0.5983	
21	0.9538		0.4055		0.3624	
22	0.2313		0.8793		0.9322	
23	0.0098		0.3316		0.0615	
24	0.0556		0.5727		0.3703	
25	0.0383		0.3244		0.2855	
26-29	0.5093		0.5209		0.9416	
30-34	0.5230		0.6489		0.7516	
35-39	0.5194		0.2912		0.4925	
40-44	0.0327		0.6644		0.4765	
45-49	0.8165		0.4152		0.8727	
50-54	0.1634		0.2090		0.1029	
55-59	0.0393		0.7252		0.9351	
60-64	0.0821		0.4765		0.4352	
65 or Older	0.1677		0.3944		0.2978	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.26C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	564	562	300	304	170	194
12	16	18	13	16	9	11
13	24	23	21	18	13	11
14	30	23	25	21	18	13
15	31	28	24	24	16	14
16	36	34	29	28	16	17
17	39	37	30	30	17	18
18	55	49	38	38	25	24
19	49	54	37	41	24	26
20	54	58	37	42	20	23
21	59	57	41	39	26	23
22	54	52	37	36	24	19
23	49	52	30	31	20	20
24	46	51	30	28	18	19
25	40	50	24	30	15	20
26-29	174	172	100	101	55	61
30-34	160	184	97	110	65	70
35-39	186	179	94	97	50	56
40-44	192	184	105	105	63	59
45-49	190	191	91	78	55	51
50-54	253	239	74	109	45	88
55-59	151	181	79	89	28	42
60-64	92	110	22	51	18	27
65 or Older	123	102	42	72	33	34

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.26D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.23	0.23	0.12	0.12	0.07	0.08
12	0.41	0.44	0.33	0.39	0.23	0.26
13	0.54	0.53	0.46	0.42	0.31	0.27
14	0.61	0.50	0.54	0.46	0.40	0.29
15	0.68	0.60	0.53	0.53	0.36	0.33
16	0.76	0.73	0.64	0.61	0.38	0.38
17	0.87	0.83	0.70	0.68	0.42	0.43
18	1.01	0.94	0.77	0.77	0.52	0.50
19	1.04	1.06	0.86	0.85	0.59	0.58
20	1.12	1.14	0.85	0.94	0.49	0.55
21	1.20	1.08	0.93	0.87	0.61	0.54
22	1.11	1.08	0.83	0.82	0.58	0.46
23	1.06	1.10	0.72	0.75	0.49	0.51
24	0.99	1.13	0.68	0.73	0.43	0.50
25	0.97	1.07	0.62	0.72	0.41	0.50
26-29	0.98	0.97	0.62	0.62	0.35	0.39
30-34	0.77	0.84	0.48	0.54	0.32	0.35
35-39	0.79	0.81	0.45	0.47	0.24	0.28
40-44	0.79	0.74	0.46	0.44	0.27	0.25
45-49	0.81	0.79	0.42	0.35	0.25	0.23
50-54	1.21	1.14	0.38	0.54	0.23	0.44
55-59	0.85	1.01	0.47	0.51	0.17	0.25
60-64	0.72	0.81	0.17	0.39	0.15	0.21
65 or Older	0.35	0.29	0.12	0.20	0.10	0.10

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.26P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.4798		0.3133		0.4302	
12	0.5055		0.6890		0.2995	
13	0.2755		0.1690		0.3742	
14	0.0051		0.0558		0.0008	
15	0.0020		0.1654		0.1439	
16	0.0111		0.4683		0.4874	
17	0.7707		0.7551		0.5293	
18	0.0180		0.1379		0.4226	
19	0.3050		0.3343		0.8061	
20	0.0490		0.2909		0.8556	
21	0.8322		0.5779		0.3202	
22	0.5789		0.4613		0.5840	
23	0.0345		0.1613		0.2101	
24	0.0381		0.9721		0.5740	
25	0.0035		0.0851		0.1126	
26-29	0.3427		0.7185		0.4172	
30-34	0.3626		0.6847		0.8468	
35-39	0.3889		0.1557		0.4424	
40-44	0.1600		0.9749		0.9337	
45-49	0.9847		0.1423		0.4439	
50-54	0.9217		0.0767		0.1147	
55-59	0.1922		0.7627		0.6974	
60-64	0.1544		0.0860		0.4859	
65 or Older	0.5029		0.2760		0.8925	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.27C Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	827	835	410	398	255	279
12	27	31	20	24	13	13
13	38	33	28	26	17	17
14	45	39	35	29	23	16
15	43	42	33	33	22	19
16	46	44	34	36	22	24
17	53	52	43	39	25	23
18	67	63	49	47	31	31
19	64	67	49	50	31	36
20	70	72	52	55	28	37
21	73	72	56	53	33	33
22	70	66	47	46	32	30
23	61	68	40	44	25	29
24	62	67	37	40	27	25
25	65	68	36	40	21	29
26-29	247	248	142	132	90	89
30-34	242	271	136	142	93	86
35-39	310	284	136	127	80	93
40-44	322	326	147	138	89	93
45-49	298	318	114	121	71	85
50-54	403	366	115	128	58	105
55-59	256	295	85	105	57	62
60-64	160	205	71	79	52	66
65 or Older	187	200	73	93	35	56

*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.27D Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.34	0.34	0.17	0.16	0.11	0.11
12	0.65	0.71	0.50	0.57	0.33	0.31
13	0.80	0.71	0.60	0.59	0.39	0.38
14	0.88	0.82	0.73	0.64	0.50	0.37
15	0.88	0.85	0.72	0.70	0.49	0.42
16	0.92	0.86	0.74	0.76	0.51	0.54
17	1.03	1.03	0.91	0.87	0.60	0.55
18	1.12	1.12	0.92	0.90	0.65	0.63
19	1.26	1.20	1.07	1.00	0.74	0.78
20	1.30	1.28	1.11	1.14	0.66	0.82
21	1.32	1.21	1.19	1.07	0.76	0.76
22	1.22	1.20	1.01	1.00	0.75	0.72
23	1.19	1.25	0.92	0.99	0.62	0.71
24	1.18	1.25	0.80	0.94	0.63	0.63
25	1.28	1.31	0.90	0.93	0.57	0.70
26-29	1.22	1.13	0.85	0.80	0.56	0.55
30-34	1.00	1.11	0.65	0.68	0.46	0.43
35-39	1.09	1.11	0.64	0.60	0.37	0.45
40-44	1.11	1.05	0.63	0.58	0.38	0.40
45-49	1.07	1.07	0.53	0.53	0.32	0.38
50-54	1.63	1.59	0.59	0.64	0.30	0.52
55-59	1.36	1.49	0.51	0.60	0.34	0.36
60-64	1.19	1.45	0.55	0.59	0.41	0.50
65 or Older	0.54	0.57	0.21	0.26	0.10	0.16

*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 1.27P Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.7537		0.6892		0.1014	
12	0.4577		0.3342		0.7262	
13	0.0784		0.4435		0.4377	
14	0.0723		0.0826		0.0023	
15	0.0003		0.0284		0.0886	
16	0.0256		0.6850		0.2520	
17	0.9282		0.8116		0.8622	
18	0.0736		0.0593		0.2339	
19	0.3953		0.3794		0.9803	
20	0.0843		0.2726		0.1936	
21	0.3004		0.6591		0.8245	
22	0.0798		0.9798		0.7674	
23	0.0352		0.0547		0.0093	
24	0.5861		0.2614		0.7484	
25	0.6963		0.2229		0.0558	
26-29	0.5724		0.3450		0.7376	
30-34	0.7118		0.8609		0.2304	
35-39	0.3883		0.8289		0.3358	
40-44	0.0560		0.9442		0.5007	
45-49	0.1228		0.8823		0.4068	
50-54	0.2067		0.2505		0.0522	
55-59	0.0185		0.3738		0.9328	
60-64	0.0231		0.6181		0.7314	
65 or Older	0.1848		0.3216		0.1739	

*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.