

Table 2.19A Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	171,827	172,217	83,066	84,958	70,257	71,519
12	339	284	169	150	61	64
13	757	679	419	431	222	187
14	1,231 ^a	1,081	767 ^a	660	447 ^b	335
15	1,653	1,568	1,110	1,066	697	645
16	1,965	1,973	1,404	1,431	946	927
17	2,304	2,210	1,692	1,626	1,262	1,167
18	3,026	2,867	2,377	2,302	1,874	1,765
19	2,745	2,956	2,123	2,270	1,690	1,781
20	2,911	3,007	2,240	2,308	1,800	1,877
21	3,073	2,967	2,253	2,270	1,878	1,893
22	3,096	3,005	2,274	2,281	1,908	1,896
23	2,912	2,954	2,105	2,158	1,762	1,774
24	3,129 ^b	2,794	2,167	1,972	1,809	1,661
25	2,765	2,947	1,942	2,075	1,637	1,735
26-29	11,703	11,865	7,418	7,627	6,314	6,241
30-34	14,642	14,333	8,044	8,196	6,708	6,906
35-39	16,161	15,201	8,521	8,171	7,270	7,008
40-44	18,332	18,474	9,188	9,497	8,018	8,534
45-49	17,261	17,487	7,949	8,652	7,120	7,635
50-54	15,622	15,653	6,090	6,389	5,427	5,610
55-59	12,865	13,275	4,662	4,744	4,196	4,106
60-64	9,865	10,333	3,014	3,338	2,737	2,974
65 or Older	23,468	24,304	5,137	5,343	4,472	4,800

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.19B Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	71.4	70.8	34.5	34.9	29.2	29.4
12	8.5	7.1	4.2	3.7	1.5	1.6
13	17.2	16.1	9.5	10.2	5.1	4.4
14	28.1 ^a	24.9	17.5 ^a	15.2	10.2 ^b	7.7
15	38.5	36.0	25.9	24.5	16.2	14.8
16	47.7	45.7	34.1	33.2	23.0	21.5
17	56.7	53.7	41.6	39.6	31.0	28.4
18	65.8	62.8	51.7	50.4	40.8	38.6
19	69.7	68.9	53.9	52.9	42.9	41.5
20	73.0	73.2	56.1	56.2	45.1	45.7
21	75.3	73.9	55.2	56.5	46.0	47.1
22	77.0	74.1	56.6	56.3	47.5	46.8
23	74.2	76.0	53.6	55.5	44.9	45.7
24	77.3	75.2	53.5	53.1	44.7	44.7
25	76.9	76.7	54.0	54.0	45.5	45.1
26-29	76.3	76.4	48.3	49.1	41.1	40.2
30-34	74.6	74.2	41.0	42.4	34.2	35.7
35-39	77.2	76.2	40.7	41.0	34.7	35.1
40-44	80.8	79.0	40.5	40.6	35.4	36.5
45-49	80.3	79.7	37.0	39.4	33.1	34.8
50-54	80.6	79.4	31.4	32.4	28.0	28.5
55-59	78.4	77.4	28.4	27.7	25.6	23.9
60-64	78.7	78.1	24.0	25.2	21.8	22.5
65 or Older	67.7	69.1	14.8	15.2	12.9	13.6

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.20A Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	161,842	161,863	69,909	70,832	59,896	60,532
12	277	230	132	122	45	52
13	653	593	362	363	190	145
14	1,097 ^b	898	656 ^b	517	368 ^b	269
15	1,467	1,352	913	866	578	533
16	1,790	1,724	1,193	1,158	787	759
17	2,079	1,985	1,390	1,361	1,031	975
18	2,722	2,575	1,983	1,918	1,597	1,462
19	2,554	2,723	1,855	1,922	1,464	1,534
20	2,707	2,796	1,955	1,997	1,608	1,633
21	2,912	2,768	2,021	2,011	1,680	1,673
22	2,922	2,817	2,020	2,008	1,725	1,697
23	2,731	2,740	1,875	1,920	1,595	1,612
24	2,942 ^a	2,658	1,903	1,729	1,605	1,491
25	2,628	2,803	1,679	1,842	1,456	1,565
26-29	11,046	11,275	6,578	6,602	5,643	5,561
30-34	13,664	13,419	6,724	6,980	5,702	5,955
35-39	15,262	14,125	7,148	6,630	6,202	5,759
40-44	17,457	17,287	7,748	7,706	6,891	7,124
45-49	16,422	16,706	6,624	7,202	6,048	6,516
50-54	14,946	14,946	5,216	5,343	4,777	4,840
55-59	12,251	12,710	3,752	3,882	3,416	3,346
60-64	9,376	9,815	2,486	2,748	2,285	2,515
65 or Older	21,936	22,919	3,698	4,004	3,203	3,517

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.20B Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	67.3	66.6	29.1	29.1	24.9	24.9
12	7.0	5.8	3.3	3.0	1.1	1.3
13	14.9	14.0	8.2	8.6	4.3	3.4
14	25.1 ^b	20.7	15.0 ^b	11.9	8.4 ^b	6.2
15	34.2 ^a	31.0	21.3	19.9	13.5	12.2
16	43.5 ^a	40.0	29.0	26.8	19.1	17.6
17	51.1	48.3	34.2	33.1	25.4	23.7
18	59.2	56.4	43.1	42.0	34.8	32.0
19	64.9	63.4	47.1	44.8	37.2	35.7
20	67.8	68.1	49.0	48.6	40.3	39.8
21	71.4	68.9	49.5	50.1	41.2	41.6
22	72.7 ^a	69.5	50.2	49.5	42.9	41.8
23	69.6	70.5	47.7	49.4	40.6	41.5
24	72.7	71.5	47.0	46.5	39.7	40.1
25	73.0	72.9	46.7	47.9	40.5	40.7
26-29	72.0	72.6	42.9	42.5	36.8	35.8
30-34	69.6	69.4	34.3	36.1	29.0	30.8
35-39	72.9	70.8	34.1	33.3	29.6	28.9
40-44	77.0 ^a	73.9	34.2	33.0	30.4	30.5
45-49	76.4	76.2	30.8	32.8	28.1	29.7
50-54	77.1	75.8	26.9	27.1	24.6	24.5
55-59	74.7	74.1	22.9	22.6	20.8	19.5
60-64	74.8	74.2	19.8	20.8	18.2	19.0
65 or Older	63.3	65.2	10.7	11.4	9.2	10.0

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.21A Daily Cigarette Use in Lifetime and Past Month Smoking Status, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	Daily Cigarette Use in Lifetime		Daily Cigarette Use in Lifetime and Cigarette Use in Past Month		Daily Cigarette Use in Lifetime but No Cigarette Use in Past Month		Percentage Who Did Not Smoke Cigarettes in the Past Month among Lifetime Daily Smokers	
	2004	2005	2004	2005	2004	2005	2004	2005
TOTAL	93,409	93,470	50,278	50,927	43,130	42,543	46.2	45.5
12	13	7	10	6	3	2	*	*
13	71	60	48	41	22	19	*	*
14	159 ^a	114	134 ^b	83	25	30	15.6	*
15	325 ^a	249	271 ^a	196	54	52	16.8	21.1
16	464	404	401	343	63	61	13.6	15.1
17	676	637	589	559	87	78	12.9	12.3
18	1,140	998	1,014 ^a	880	126	118	11.0	11.9
19	1,134	1,215	993	1,040	141	174	12.4	14.4
20	1,316	1,356	1,169	1,180	148	176	11.2	13.0
21	1,506	1,350	1,269	1,166	237	184	15.7	13.6
22	1,484	1,525	1,262	1,292	222	234	14.9	15.3
23	1,534	1,485	1,260	1,256	274	229	17.9	15.4
24	1,643	1,522	1,282	1,239	361	284	22.0	18.6
25	1,456	1,618	1,178	1,285	278	333	19.1	20.6
26-29	6,042	5,965	4,629	4,664	1,414	1,302	23.4	21.8
30-34	6,850	7,416	4,745	5,082	2,105	2,334	30.7	31.5
35-39	7,935	7,218	5,307	4,990	2,628	2,228	33.1	30.9
40-44	10,037	9,483	6,338	6,303	3,699 ^a	3,180	36.9	33.5
45-49	10,137	10,413	5,613	5,920	4,524	4,493	44.6	43.1
50-54	9,475	9,385	4,377	4,562	5,097	4,823	53.8	51.4
55-59	8,220	8,412	3,294	3,179	4,925	5,233	59.9	62.2
60-64	6,518	6,713	2,199	2,374	4,319	4,339	66.3	64.6
65 or Older	15,274	15,927	2,897	3,288	12,378	12,639	81.0	79.4

*Low precision; no estimate reported.

NOTE: Daily Cigarette Use in Lifetime is defined as ever smoking every day for at least 30 days.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.21B Daily Cigarette Use in Lifetime and Past Month Smoking Status, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	Daily Cigarette Use in Lifetime		Daily Cigarette Use in Lifetime and Cigarette Use in Past Month		Daily Cigarette Use in Lifetime but No Cigarette Use in Past Month		Percentage Who Did Not Smoke Cigarettes in the Past Month among Lifetime Daily Smokers	
	2004	2005	2004	2005	2004	2005	2004	2005
TOTAL	38.8	38.4	20.9	20.9	17.9	17.5	46.2	45.5
12	0.3	0.2	0.3	0.1	0.1	0.0	*	*
13	1.6	1.4	1.1	1.0	0.5	0.4	*	*
14	3.6 ^a	2.6	3.1 ^b	1.9	0.6	0.7	15.6	*
15	7.6 ^a	5.7	6.3 ^b	4.5	1.3	1.2	16.8	21.1
16	11.3 ^a	9.4	9.7 ^a	7.9	1.5	1.4	13.6	15.1
17	16.6	15.5	14.5	13.6	2.2	1.9	12.9	12.3
18	24.8 ^a	21.9	22.1 ^a	19.3	2.7	2.6	11.0	11.9
19	28.8	28.3	25.2	24.2	3.6	4.1	12.4	14.4
20	33.0	33.0	29.3	28.7	3.7	4.3	11.2	13.0
21	36.9 ^a	33.6	31.1	29.0	5.8	4.6	15.7	13.6
22	36.9	37.6	31.4	31.9	5.5	5.8	14.9	15.3
23	39.1	38.2	32.1	32.3	7.0	5.9	17.9	15.4
24	40.6	41.0	31.7	33.3	8.9	7.6	22.0	18.6
25	40.5	42.1	32.7	33.4	7.7	8.7	19.1	20.6
26-29	39.4	38.4	30.2	30.0	9.2	8.4	23.4	21.8
30-34	34.9 ^a	38.4	24.2	26.3	10.7	12.1	30.7	31.5
35-39	37.9	36.2	25.3	25.0	12.5	11.2	33.1	30.9
40-44	44.3 ^a	40.6	27.9	27.0	16.3 ^b	13.6	36.9	33.5
45-49	47.1	47.5	26.1	27.0	21.0	20.5	44.6	43.1
50-54	48.9	47.6	22.6	23.1	26.3	24.5	53.8	51.4
55-59	50.1	49.0	20.1	18.5	30.0	30.5	59.9	62.2
60-64	52.0	50.7	17.5	17.9	34.4	32.8	66.3	64.6
65 or Older	44.0	45.3	8.4	9.3	35.7	35.9	81.0	79.4

*Low precision; no estimate reported.

NOTE: Daily Cigarette Use in Lifetime is defined as ever smoking every day for at least 30 days.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.22A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	44,625	44,793	9,756	10,442	7,154	7,682
12	76	56	39	24	13	7
13	139	130	78	73	36	33
14	223	224	131	140	61	59
15	364	386	220	249	108	96
16	431	464	281	292	164	167
17	558	542	335	328	188	161
18	817	716	445	456	264	251
19	685	743	354	409	194	230
20	777	864	333	360	190	203
21	934	849	314	362	208	223
22	928	888	318	376	188	222
23	884	920	253	279	172	182
24	978	888	293	263	196	176
25	874	900	252	261	177	177
26-29	4,159	3,967	955	883	686	679
30-34	4,978	5,205	1,010	1,072	797	874
35-39	5,442	5,410	939	1,005	759	798
40-44	4,940	5,525	779 ^a	1,064	654 ^a	907
45-49	4,382	4,058	589	569	447	510
50-54	3,108	2,978	418	491	342	398
55-59	2,511	2,617	377	376	308	358
60-64	1,717	2,070	267	427	228	353
65 or Older	4,719	4,394	777	685	774	618

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.22B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	18.6	18.4	4.1	4.3	3.0	3.2
12	1.9	1.4	1.0	0.6	0.3	0.2
13	3.2	3.1	1.8	1.7	0.8	0.8
14	5.1	5.2	3.0	3.2	1.4	1.4
15	8.5	8.8	5.1	5.7	2.5	2.2
16	10.5	10.7	6.8	6.8	4.0	3.9
17	13.7	13.2	8.2	8.0	4.6	3.9
18	17.8	15.7	9.7	10.0	5.7	5.5
19	17.4	17.3	9.0	9.5	4.9	5.4
20	19.5	21.0	8.4	8.8	4.8	4.9
21	22.9	21.1	7.7	9.0	5.1	5.6
22	23.1	21.9	7.9	9.3	4.7	5.5
23	22.5	23.7	6.4	7.2	4.4	4.7
24	24.2	23.9	7.2	7.1	4.9	4.7
25	24.3	23.4	7.0	6.8	4.9	4.6
26-29	27.1	25.5	6.2	5.7	4.5	4.4
30-34	25.4	26.9	5.1	5.5	4.1	4.5
35-39	26.0	27.1	4.5	5.0	3.6	4.0
40-44	21.8	23.6	3.4 ^a	4.5	2.9	3.9
45-49	20.4	18.5	2.7	2.6	2.1	2.3
50-54	16.0	15.1	2.2	2.5	1.8	2.0
55-59	15.3	15.3	2.3	2.2	1.9	2.1
60-64	13.7	15.6	2.1	3.2	1.8	2.7
65 or Older	13.6	12.5	2.2	1.9	2.2	1.8

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.23A Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	87,409	88,260	25,974	26,908	13,727	13,640
12	81	68	44	36	26	12
13	252	210	141	112	53	45
14	452	440	285	290	125	96
15	748	687	494	481	204	189
16	943	958	695	708	335	309
17	1,267	1,210	911	858	458	413
18	1,779	1,696	1,330	1,283	766	695
19	1,631	1,735	1,102	1,166	595	598
20	1,768	1,773	1,094	1,105	529	589
21	1,815	1,822	962	1,043	541	482
22	1,904	1,764	868	923	439	407
23	1,752	1,780	809	831	421	405
24	1,909	1,765	844	780	437	375
25	1,654	1,699	716	725	375	344
26-29	6,942	6,895	2,447	2,583	1,196	1,171
30-34	7,808	7,886	2,342	2,483	1,087	1,201
35-39	8,350	8,238	2,294	2,719	1,154	1,383
40-44	9,076	9,082	2,277	2,584	1,204	1,382
45-49	8,824	9,119	2,296	2,427	1,293	1,244
50-54	7,809	7,646	1,399	1,495	864	945
55-59	6,530	6,529	1,196	1,002	741	507
60-64	4,911	4,872	639	508	388	294
65 or Older	9,203 ^a	10,383	791	766	497	555

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.23B Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	36.3	36.3	10.8	11.1	5.7	5.6
12	2.0	1.7	1.1	0.9	0.7	0.3
13	5.7	5.0	3.2	2.6	1.2	1.1
14	10.3	10.1	6.5	6.7	2.9	2.2
15	17.4	15.8	11.5	11.0	4.8	4.3
16	22.9	22.2	16.9	16.4	8.1	7.2
17	31.2	29.4	22.4	20.9	11.3	10.0
18	38.7	37.1	28.9	28.1	16.7	15.2
19	41.4	40.4	28.0	27.2	15.1	13.9
20	44.3	43.2	27.4	26.9	13.3	14.3
21	44.5	45.4	23.6	26.0	13.3	12.0
22	47.4 ^a	43.5	21.6	22.8	10.9	10.0
23	44.6	45.8	20.6	21.4	10.7	10.4
24	47.2	47.5	20.9	21.0	10.8	10.1
25	46.0	44.2	19.9	18.9	10.4	8.9
26-29	45.2	44.4	15.9	16.6	7.8	7.5
30-34	39.8	40.8	11.9	12.8	5.5	6.2
35-39	39.9	41.3	11.0 ^a	13.6	5.5	6.9
40-44	40.0	38.8	10.0	11.1	5.3	5.9
45-49	41.0	41.6	10.7	11.1	6.0	5.7
50-54	40.3	38.8	7.2	7.6	4.5	4.8
55-59	39.8	38.1	7.3	5.8	4.5	3.0
60-64	39.2	36.8	5.1	3.8	3.1	2.2
65 or Older	26.5	29.5	2.3	2.2	1.4	1.6

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.24A Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	198,220 ^b	201,667	156,686 ^b	161,626	120,934 ^b	126,028
12	441	385	260	229	92	99
13	969	909	685	675	270	244
14	1,598 ^a	1,446	1,247	1,170	540	461
15	2,178	2,186	1,773	1,779	878	854
16	2,493	2,618	2,102	2,208	1,142	1,166
17	2,916	2,763	2,469	2,394	1,515	1,367
18	3,514	3,422	3,120	3,043	2,130	2,026
19	3,204	3,471	2,864	3,106	2,027	2,235
20	3,395	3,472	3,047	3,126	2,246	2,367
21	3,681	3,569	3,436	3,381	2,846	2,788
22	3,612	3,552	3,362	3,255	2,776	2,684
23	3,460	3,541	3,118	3,251	2,504	2,701
24	3,632 ^a	3,319	3,250 ^a	2,987	2,580	2,408
25	3,254	3,489	2,931	3,155	2,365	2,574
26-29	13,908	14,133	12,046	12,525	9,705	9,895
30-34	17,573	17,387	14,775	15,048	11,457	11,905
35-39	18,984	18,247	15,806	15,527	12,687	12,434
40-44	20,747	21,186	17,017	17,295	13,623	14,044
45-49	19,687	20,201	15,359	16,132	12,355	13,213
50-54	17,304	18,091	13,368	14,257	10,779	11,501
55-59	14,440	15,101	10,289	10,718	7,922	8,720
60-64	10,675	11,147	7,852	7,958	6,255	6,289
65 or Older	26,554 ^a	28,032	16,511 ^b	18,409	12,242 ^b	14,054

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.24B Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	82.4	82.9	65.1 ^a	66.5	50.3 ^b	51.8
12	11.1	9.6	6.6	5.7	2.3	2.5
13	22.1	21.5	15.6	16.0	6.1	5.8
14	36.5 ^a	33.3	28.5	27.0	12.3	10.6
15	50.7	50.2	41.3	40.8	20.4	19.6
16	60.6	60.7	51.1	51.2	27.7	27.0
17	71.7 ^b	67.2	60.7	58.2	37.3 ^b	33.2
18	76.5	74.9	67.9	66.6	46.3	44.4
19	81.4	80.9	72.7	72.4	51.5	52.1
20	85.1	84.5	76.4	76.1	56.3	57.6
21	90.2	88.9	84.2	84.2	69.8	69.4
22	89.9	87.6	83.6 ^a	80.3	69.1	66.2
23	88.1 ^b	91.1	79.4 ^b	83.6	63.8 ^b	69.5
24	89.7	89.3	80.3	80.4	63.7	64.8
25	90.4	90.8	81.4	82.1	65.7	67.0
26-29	90.6	91.0	78.5	80.7	63.2	63.7
30-34	89.5	89.9	75.3	77.9	58.4 ^a	61.6
35-39	90.6	91.5	75.5	77.9	60.6	62.4
40-44	91.5	90.6	75.0	74.0	60.1	60.1
45-49	91.5	92.1	71.4	73.6	57.5	60.2
50-54	89.3	91.8	69.0	72.3	55.6	58.3
55-59	88.0	88.0	62.7	62.5	48.3	50.8
60-64	85.1	84.3	62.6	60.2	49.9	47.5
65 or Older	76.6	79.7	47.6 ^a	52.3	35.3 ^a	40.0

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.25A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Numbers in Thousands, 2004 and 2005

Age Category	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2004	2005	2004	2005	2004	2005
TOTAL	120,934 ^b	126,028	54,725	55,090	16,689	16,035
12	92	99	42	53	4	5
13	270	244	129	109	16	15
14	540	461	244	224	43	41
15	878	854	544	470	93	103
16	1,142	1,166	754	726	187	179
17	1,515	1,367	1,080 ^a	931	328	268
18	2,130	2,026	1,522	1,408	507	486
19	2,027	2,235	1,472	1,636	548	601
20	2,246	2,367	1,610	1,640	649	597
21	2,846	2,788	1,965	2,006	781	794
22	2,776	2,684	1,882	1,891	715	777
23	2,504	2,701	1,694	1,853	562	643
24	2,580	2,408	1,632	1,549	552	532
25	2,365	2,574	1,494	1,614	533	536
26-29	9,705	9,895	5,643	5,483	1,648	1,665
30-34	11,457	11,905	5,623	5,991	1,627	1,692
35-39	12,687	12,434	6,236 ^a	5,526	1,523	1,395
40-44	13,623	14,044	6,099	6,197	1,894	1,676
45-49	12,355	13,213	4,986	5,120	1,483	1,392
50-54	10,779	11,501	3,663	3,739	1,210	1,080
55-59	7,922	8,720	2,309	2,433	701	562
60-64	6,255	6,289	1,705	1,570	467	404
65 or Older	12,242 ^b	14,054	2,398	2,921	616	591

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.25B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Percentages, 2004 and 2005

Age Category	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2004	2005	2004	2005	2004	2005
TOTAL	50.3 ^b	51.8	22.8	22.7	6.9	6.6
12	2.3	2.5	1.1	1.3	0.1	0.1
13	6.1	5.8	2.9	2.6	0.4	0.4
14	12.3	10.6	5.6	5.2	1.0	1.0
15	20.4	19.6	12.7 ^a	10.8	2.2	2.4
16	27.7	27.0	18.3	16.8	4.5	4.2
17	37.3 ^b	33.2	26.6 ^b	22.7	8.1 ^a	6.5
18	46.3	44.4	33.1	30.8	11.0	10.6
19	51.5	52.1	37.4	38.1	13.9	14.0
20	56.3	57.6	40.3	39.9	16.3	14.5
21	69.8	69.4	48.2	49.9	19.2	19.8
22	69.1	66.2	46.8	46.6	17.8	19.2
23	63.8 ^b	69.5	43.1 ^b	47.7	14.3	16.5
24	63.7	64.8	40.3	41.7	13.7	14.3
25	65.7	67.0	41.5	42.0	14.8	14.0
26-29	63.2	63.7	36.8	35.3	10.7	10.7
30-34	58.4 ^a	61.6	28.6	31.0	8.3	8.8
35-39	60.6	62.4	29.8	27.7	7.3	7.0
40-44	60.1	60.1	26.9	26.5	8.4	7.2
45-49	57.5	60.2	23.2	23.3	6.9	6.3
50-54	55.6	58.3	18.9	19.0	6.2	5.5
55-59	48.3	50.8	14.1	14.2	4.3	3.3
60-64	49.9	47.5	13.6	11.9	3.7	3.1
65 or Older	35.3 ^a	40.0	6.9	8.3	1.8	1.7

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.