

Table 2.1A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	171,827	172,217	83,066	84,958	70,257	71,519
Cigarettes	161,842	161,863	69,909	70,832	59,896	60,532
Smokeless Tobacco	44,625	44,793	9,756	10,442	7,154	7,682
Cigars	87,409	88,260	25,974	26,908	13,727	13,640
Pipe Tobacco ²	39,499	38,462	--	--	1,835	2,190
ALCOHOL	198,220 ^b	201,667	156,686 ^b	161,626	120,934 ^b	126,028
Binge Alcohol Use ³	--	--	--	--	54,725	55,090
Heavy Alcohol Use ³	--	--	--	--	16,689	16,035

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.1B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	71.4	70.8	34.5	34.9	29.2	29.4
Cigarettes	67.3	66.6	29.1	29.1	24.9	24.9
Smokeless Tobacco	18.6	18.4	4.1	4.3	3.0	3.2
Cigars	36.3	36.3	10.8	11.1	5.7	5.6
Pipe Tobacco ²	16.4	15.8	--	--	0.8	0.9
ALCOHOL	82.4	82.9	65.1 ^a	66.5	50.3 ^b	51.8
Binge Alcohol Use ³	--	--	--	--	22.8	22.7
Heavy Alcohol Use ³	--	--	--	--	6.9	6.6

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.2A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	8,249 ^b	7,795	5,562	5,365	3,635 ^b	3,324
Cigarettes	7,363 ^b	6,781	4,645 ^a	4,387	2,998 ^a	2,733
Smokeless Tobacco	1,791	1,801	1,085	1,105	571	523
Cigars	3,743	3,573	2,568	2,485	1,201 ^a	1,064
Pipe Tobacco ²	700	634	--	--	178	158
ALCOHOL	10,596	10,306	8,537	8,454	4,436 ^a	4,191
Binge Alcohol Use ³	--	--	--	--	2,794 ^b	2,513
Heavy Alcohol Use ³	--	--	--	--	671	611

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.2B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	32.7 ^b	30.7	22.1	21.2	14.4 ^b	13.1
Cigarettes	29.2 ^b	26.7	18.4 ^a	17.3	11.9 ^b	10.8
Smokeless Tobacco	7.1	7.1	4.3	4.4	2.3	2.1
Cigars	14.8	14.1	10.2	9.8	4.8 ^a	4.2
Pipe Tobacco ²	2.8	2.5	--	--	0.7	0.6
ALCOHOL	42.0 ^a	40.6	33.9	33.3	17.6 ^a	16.5
Binge Alcohol Use ³	--	--	--	--	11.1 ^b	9.9
Heavy Alcohol Use ³	--	--	--	--	2.7	2.4

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.3A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	23,658	23,496	17,481	17,637	14,359	14,382
Cigarettes	22,118	21,879	15,290	15,348	12,731	12,667
Smokeless Tobacco	6,878	6,768	2,563	2,766	1,589	1,664
Cigars	14,213	14,036	7,724	7,856	4,103	3,895
Pipe Tobacco ²	2,661	2,693	--	--	396	471
ALCOHOL	27,752	27,836	25,127	25,303	19,474	19,783
Binge Alcohol Use ³	--	--	--	--	13,271	13,597
Heavy Alcohol Use ³	--	--	--	--	4,848	4,967

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.3B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	73.5	72.3	54.3	54.3	44.6	44.3
Cigarettes	68.7 ^a	67.3	47.5	47.2	39.5	39.0
Smokeless Tobacco	21.4	20.8	8.0	8.5	4.9	5.1
Cigars	44.1	43.2	24.0	24.2	12.7	12.0
Pipe Tobacco ²	8.3	8.3	--	--	1.2	1.5
ALCOHOL	86.2	85.7	78.0	77.9	60.5	60.9
Binge Alcohol Use ³	--	--	--	--	41.2	41.9
Heavy Alcohol Use ³	--	--	--	--	15.1	15.3

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.4A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	139,921	140,926	60,023	61,957	52,263	53,813
Cigarettes	132,360	133,203	49,974	51,098	44,166	45,133
Smokeless Tobacco	35,956	36,224	6,109	6,572	4,994	5,495
Cigars	69,453	70,652	15,681	16,567	8,423	8,681
Pipe Tobacco ²	36,138	35,135	--	--	1,261	1,561
ALCOHOL	159,872 ^b	163,525	123,022 ^b	127,869	97,023 ^b	102,054
Binge Alcohol Use ³	--	--	--	--	38,661	38,980
Heavy Alcohol Use ³	--	--	--	--	11,170	10,457

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.4B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	76.4	76.0	32.8	33.4	28.5	29.0
Cigarettes	72.3	71.9	27.3	27.6	24.1	24.3
Smokeless Tobacco	19.6	19.5	3.3	3.5	2.7	3.0
Cigars	37.9	38.1	8.6	8.9	4.6	4.7
Pipe Tobacco ²	19.7	19.0	--	--	0.7	0.8
ALCOHOL	87.3	88.2	67.2 ^b	69.0	53.0 ^b	55.1
Binge Alcohol Use ³	--	--	--	--	21.1	21.0
Heavy Alcohol Use ³	--	--	--	--	6.1	5.6

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.5A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	1,095 ^a	963	588	581	284	251
Cigarettes	930	823	493	485	235	197
Smokeless Tobacco	216	186	117	97	49	41
Cigars	333	278	184	147	79	57
Pipe Tobacco ²	71	61	--	--	20	20
ALCOHOL	1,410	1,294	946	904	362	343
Binge Alcohol Use ³	--	--	--	--	171	162
Heavy Alcohol Use ³	--	--	--	--	21	20

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.5B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	13.1 ^a	11.7	7.0	7.1	3.4	3.0
Cigarettes	11.1	10.0	5.9	5.9	2.8	2.4
Smokeless Tobacco	2.6	2.3	1.4	1.2	0.6	0.5
Cigars	4.0	3.4	2.2	1.8	0.9	0.7
Pipe Tobacco ²	0.8	0.7	--	--	0.2	0.2
ALCOHOL	16.9	15.7	11.3	11.0	4.3	4.2
Binge Alcohol Use ³	--	--	--	--	2.0	2.0
Heavy Alcohol Use ³	--	--	--	--	0.2	0.2

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.6A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	2,884 ^a	2,649	1,877	1,726	1,144 ^b	980
Cigarettes	2,564 ^b	2,250	1,568 ^a	1,383	945 ^b	802
Smokeless Tobacco	586	609	352	389	170	155
Cigars	1,200	1,127	779	771	329	285
Pipe Tobacco ²	243	209	--	--	66	56
ALCOHOL	3,776	3,632	3,020	2,948	1,417	1,315
Binge Alcohol Use ³	--	--	--	--	788	694
Heavy Alcohol Use ³	--	--	--	--	136	144

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.6B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	33.3 ^b	30.5	21.7 ^a	19.8	13.2 ^b	11.3
Cigarettes	29.6 ^b	25.9	18.1 ^b	15.9	10.9 ^b	9.2
Smokeless Tobacco	6.8	7.0	4.1	4.5	2.0	1.8
Cigars	13.8	13.0	9.0	8.9	3.8	3.3
Pipe Tobacco ²	2.8	2.4	--	--	0.8	0.6
ALCOHOL	43.6	41.8	34.8	33.9	16.4	15.1
Binge Alcohol Use ³	--	--	--	--	9.1 ^a	8.0
Heavy Alcohol Use ³	--	--	--	--	1.6	1.7

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.7A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	4,269	4,183	3,097	3,058	2,208	2,093
Cigarettes	3,869	3,708	2,583	2,519	1,818	1,734
Smokeless Tobacco	988	1,005	616	619	352	328
Cigars	2,210	2,168	1,605	1,566	793	722
Pipe Tobacco ²	386	364	--	--	92	83
ALCOHOL	5,410	5,381	4,571	4,602	2,657	2,533
Binge Alcohol Use ³	--	--	--	--	1,834 ^a	1,657
Heavy Alcohol Use ³	--	--	--	--	515	447

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.7B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	52.2 ^a	49.6	37.8	36.3	27.0 ^a	24.8
Cigarettes	47.3 ^b	44.0	31.6	29.9	22.2	20.6
Smokeless Tobacco	12.1	11.9	7.5	7.4	4.3	3.9
Cigars	27.0	25.7	19.6	18.6	9.7	8.6
Pipe Tobacco ²	4.7	4.3	--	--	1.1	1.0
ALCOHOL	66.1 ^a	63.9	55.9	54.6	32.5 ^a	30.1
Binge Alcohol Use ³	--	--	--	--	22.4 ^b	19.7
Heavy Alcohol Use ³	--	--	--	--	6.3 ^a	5.3

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.8A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	8,682	8,830	6,740	6,881	5,364	5,423
Cigarettes	7,983	8,093	5,793	5,837	4,669	4,629
Smokeless Tobacco	2,280	2,324	1,133	1,225	647	684
Cigars	5,179	5,205	3,525	3,554	1,890	1,881
Pipe Tobacco ²	920	954	--	--	200	242
ALCOHOL	10,113	10,365	9,031	9,275	6,402	6,628
Binge Alcohol Use ³	--	--	--	--	4,604	4,684
Heavy Alcohol Use ³	--	--	--	--	1,704	1,685

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.8B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	69.3	68.1	53.8	53.1	42.8	41.8
Cigarettes	63.7	62.4	46.3	45.0	37.3	35.7
Smokeless Tobacco	18.2	17.9	9.0	9.4	5.2	5.3
Cigars	41.4	40.1	28.1	27.4	15.1	14.5
Pipe Tobacco ²	7.4	7.4	--	--	1.6	1.9
ALCOHOL	80.8	79.9	72.1	71.5	51.1	51.1
Binge Alcohol Use ³	--	--	--	--	36.8	36.1
Heavy Alcohol Use ³	--	--	--	--	13.6	13.0

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.9A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	14,976	14,666	10,741	10,756	8,995	8,959
Cigarettes	14,136	13,786	9,497	9,511	8,062	8,037
Smokeless Tobacco	4,598	4,444	1,430	1,541	941	980
Cigars	9,034	8,831	4,199	4,302	2,213	2,013
Pipe Tobacco ²	1,740	1,739	--	--	196	229
ALCOHOL	17,639	17,471	16,096	16,028	13,072	13,155
Binge Alcohol Use ³	--	--	--	--	8,667	8,913
Heavy Alcohol Use ³	--	--	--	--	3,144	3,282

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.9B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	76.1	75.1	54.6	55.1	45.7	45.9
Cigarettes	71.9	70.6	48.3	48.7	41.0	41.2
Smokeless Tobacco	23.4	22.8	7.3	7.9	4.8	5.0
Cigars	45.9	45.2	21.3	22.0	11.3	10.3
Pipe Tobacco ²	8.8	8.9	--	--	1.0	1.2
ALCOHOL	89.7	89.5	81.8	82.1	66.5	67.4
Binge Alcohol Use ³	--	--	--	--	44.1	45.7
Heavy Alcohol Use ³	--	--	--	--	16.0	16.8

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.10A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	26,346	26,198	15,462	15,823	13,022	13,147
Cigarettes	24,709	24,694	13,302	13,582	11,344	11,516
Smokeless Tobacco	9,137	9,172	1,964	1,955	1,483	1,553
Cigars	14,750	14,781	4,789	5,066	2,283	2,372
Pipe Tobacco ²	3,270	3,279	--	--	248	291
ALCOHOL	31,481	31,520	26,821 ^a	27,573	21,161	21,800
Binge Alcohol Use ³	--	--	--	--	11,265	11,474
Heavy Alcohol Use ³	--	--	--	--	3,275	3,357

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.10B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	75.3	75.2	44.2	45.4	37.2	37.7
Cigarettes	70.6	70.8	38.0	39.0	32.4	33.0
Smokeless Tobacco	26.1	26.3	5.6	5.6	4.2	4.5
Cigars	42.2	42.4	13.7	14.5	6.5	6.8
Pipe Tobacco ²	9.3	9.4	--	--	0.7	0.8
ALCOHOL	90.0	90.4	76.7 ^a	79.1	60.5	62.5
Binge Alcohol Use ³	--	--	--	--	32.2	32.9
Heavy Alcohol Use ³	--	--	--	--	9.4	9.6

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.11A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	113,575	114,727	44,562	46,134	39,241	40,666
Cigarettes	107,651	108,509	36,673	37,516	32,822	33,616
Smokeless Tobacco	26,819	27,052	4,145	4,617	3,512	3,942
Cigars	54,703	55,871	10,892	11,501	6,140	6,309
Pipe Tobacco ²	32,869	31,856	--	--	1,013	1,270
ALCOHOL	128,391 ^b	132,005	96,201 ^b	100,296	75,862 ^b	80,255
Binge Alcohol Use ³	--	--	--	--	27,396	27,505
Heavy Alcohol Use ³	--	--	--	--	7,895	7,100

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.11B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	76.7	76.2	30.1	30.6	26.5	27.0
Cigarettes	72.7	72.1	24.8	24.9	22.2	22.3
Smokeless Tobacco	18.1	18.0	2.8	3.1	2.4	2.6
Cigars	36.9	37.1	7.4	7.6	4.1	4.2
Pipe Tobacco ²	22.2	21.2	--	--	0.7	0.8
ALCOHOL	86.7	87.7	64.9 ^a	66.6	51.2 ^a	53.3
Binge Alcohol Use ³	--	--	--	--	18.5	18.3
Heavy Alcohol Use ³	--	--	--	--	5.3 ^a	4.7

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.12A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	163,579	164,422	77,504	79,593	66,622	68,195
Cigarettes	154,479	155,082	65,264	66,446	56,897	57,799
Smokeless Tobacco	42,835	42,992	8,672	9,338	6,583	7,158
Cigars	83,666	84,687	23,405	24,423	12,526	12,576
Pipe Tobacco ²	38,799	37,829	--	--	1,657	2,032
ALCOHOL	187,624 ^b	191,361	148,149 ^b	153,172	116,498 ^b	121,837
Binge Alcohol Use ³	--	--	--	--	51,931	52,577
Heavy Alcohol Use ³	--	--	--	--	16,018	15,424

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.12B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	76.0	75.5	36.0	36.5	30.9	31.3
Cigarettes	71.8	71.2	30.3	30.5	26.4	26.5
Smokeless Tobacco	19.9	19.7	4.0	4.3	3.1	3.3
Cigars	38.9	38.9	10.9	11.2	5.8	5.8
Pipe Tobacco ²	18.0	17.4	--	--	0.8	0.9
ALCOHOL	87.1	87.8	68.8 ^a	70.3	54.1 ^b	55.9
Binge Alcohol Use ³	--	--	--	--	24.1	24.1
Heavy Alcohol Use ³	--	--	--	--	7.4	7.1

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.13A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	92,403	92,316	49,326	49,876	41,569	42,175
Cigarettes	84,385	84,094	37,803	37,660	32,278	32,312
Smokeless Tobacco	37,569	38,121	8,987	9,497	6,730	7,174
Cigars	65,580	65,743	20,686	21,321	11,375	11,355
Pipe Tobacco ²	35,110	34,099	--	--	1,579	1,877
ALCOHOL	100,016 ^b	101,742	81,408	82,926	66,317 ^a	68,497
Binge Alcohol Use ³	--	--	--	--	36,195	36,025
Heavy Alcohol Use ³	--	--	--	--	12,388	12,172

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.13B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	79.3	78.3	42.3	42.3	35.7	35.8
Cigarettes	72.4	71.3	32.5	31.9	27.7	27.4
Smokeless Tobacco	32.3	32.3	7.7	8.1	5.8	6.1
Cigars	56.3	55.8	17.8	18.1	9.8	9.6
Pipe Tobacco ²	30.1	28.9	--	--	1.4	1.6
ALCOHOL	85.9	86.3	69.9	70.3	56.9	58.1
Binge Alcohol Use ³	--	--	--	--	31.1	30.5
Heavy Alcohol Use ³	--	--	--	--	10.6	10.3

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.14A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	79,425	79,901	33,740	35,082	28,688	29,344
Cigarettes	77,457	77,769	32,105	33,173	27,618	28,220
Smokeless Tobacco	7,057	6,672	769	946	424	508
Cigars	21,829	22,517	5,288	5,586	2,352	2,285
Pipe Tobacco ²	4,389	4,363	--	--	256	313
ALCOHOL	98,203 ^a	99,925	75,278 ^b	78,700	54,616 ^b	57,531
Binge Alcohol Use ³	--	--	--	--	18,530	19,065
Heavy Alcohol Use ³	--	--	--	--	4,301	3,863

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.14B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	64.0	63.8	27.2	28.0	23.1	23.4
Cigarettes	62.4	62.1	25.9	26.5	22.3	22.5
Smokeless Tobacco	5.7	5.3	0.6	0.8	0.3	0.4
Cigars	17.6	18.0	4.3	4.5	1.9	1.8
Pipe Tobacco ²	3.5	3.5	--	--	0.2	0.3
ALCOHOL	79.2	79.8	60.7 ^b	62.8	44.0 ^a	45.9
Binge Alcohol Use ³	--	--	--	--	14.9	15.2
Heavy Alcohol Use ³	--	--	--	--	3.5 ^a	3.1

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.15A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 to 17: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	4,345	4,136	2,921	2,917	1,972	1,843
Cigarettes	3,684 ^b	3,407	2,227	2,185	1,453	1,387
Smokeless Tobacco	1,411	1,413	909	926	521	476
Cigars	2,368	2,290	1,629	1,627	852 ^a	748
Pipe Tobacco ²	478	402	--	--	110	109
ALCOHOL	5,350	5,225	4,206	4,144	2,216	2,057
Binge Alcohol Use ³	--	--	--	--	1,496 ^a	1,341
Heavy Alcohol Use ³	--	--	--	--	407	383

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.15B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 to 17: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	33.7 ^a	31.9	22.7	22.5	15.3	14.2
Cigarettes	28.6 ^b	26.3	17.3	16.9	11.3	10.7
Smokeless Tobacco	11.0	10.9	7.1	7.2	4.0	3.7
Cigars	18.4	17.7	12.7	12.6	6.6 ^a	5.8
Pipe Tobacco ²	3.7	3.1	--	--	0.9	0.8
ALCOHOL	41.5	40.3	32.7	32.0	17.2 ^a	15.9
Binge Alcohol Use ³	--	--	--	--	11.6 ^a	10.4
Heavy Alcohol Use ³	--	--	--	--	3.2	3.0

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.16A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 to 17: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	3,904 ^a	3,659	2,640 ^a	2,447	1,663 ^a	1,481
Cigarettes	3,679 ^b	3,374	2,417 ^a	2,202	1,545 ^b	1,345
Smokeless Tobacco	380	388	176	179	50	47
Cigars	1,375	1,283	939	858	349	316
Pipe Tobacco ²	222	232	--	--	68	49
ALCOHOL	5,246	5,081	4,331	4,310	2,220	2,134
Binge Alcohol Use ³	--	--	--	--	1,298	1,172
Heavy Alcohol Use ³	--	--	--	--	264	229

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.16B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 to 17: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	31.6 ^a	29.5	21.4 ^a	19.7	13.5 ^b	11.9
Cigarettes	29.8 ^b	27.2	19.6 ^b	17.8	12.5 ^b	10.8
Smokeless Tobacco	3.1	3.1	1.4	1.4	0.4	0.4
Cigars	11.1	10.3	7.6	6.9	2.8	2.5
Pipe Tobacco ²	1.8	1.9	--	--	0.5	0.4
ALCOHOL	42.5	41.0	35.1	34.7	18.0	17.2
Binge Alcohol Use ³	--	--	--	--	10.5 ^a	9.4
Heavy Alcohol Use ³	--	--	--	--	2.1	1.8

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.17A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 18 to 25: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	12,764	12,618	10,063	10,128	8,368	8,425
Cigarettes	11,654	11,456	8,348	8,350	7,041	7,007
Smokeless Tobacco	5,605	5,479	2,366	2,518	1,532	1,588
Cigars	9,140	9,090	5,651	5,742	3,179	2,997
Pipe Tobacco ²	2,157	2,136	--	--	335	372
ALCOHOL	14,101	14,112	12,900	13,069	10,501 ^a	10,830
Binge Alcohol Use ³	--	--	--	--	8,101 ^a	8,442
Heavy Alcohol Use ³	--	--	--	--	3,434	3,545

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.17B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 18 to 25: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	78.9 ^a	77.2	62.2	62.0	51.7	51.6
Cigarettes	72.1 ^a	70.1	51.6	51.1	43.5	42.9
Smokeless Tobacco	34.7	33.5	14.6	15.4	9.5	9.7
Cigars	56.5	55.7	34.9	35.1	19.7	18.3
Pipe Tobacco ²	13.3	13.1	--	--	2.1	2.3
ALCOHOL	87.2	86.4	79.8	80.0	64.9	66.3
Binge Alcohol Use ³	--	--	--	--	50.1	51.7
Heavy Alcohol Use ³	--	--	--	--	21.2	21.7

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.18A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 18 to 25: Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	10,893	10,878	7,418	7,509	5,990	5,957
Cigarettes	10,464	10,423	6,942	6,998	5,690	5,660
Smokeless Tobacco	1,273	1,289	197	248	56	75
Cigars	5,073	4,945	2,073	2,115	924	898
Pipe Tobacco ²	504	557	--	--	61	100
ALCOHOL	13,651	13,724	12,228	12,234	8,974	8,953
Binge Alcohol Use ³	--	--	--	--	5,170	5,155
Heavy Alcohol Use ³	--	--	--	--	1,414	1,422

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.18B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 18 to 25: Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	68.0	67.4	46.3	46.5	37.4	36.9
Cigarettes	65.3	64.5	43.3	43.3	35.5	35.0
Smokeless Tobacco	7.9	8.0	1.2	1.5	0.4	0.5
Cigars	31.7	30.6	12.9	13.1	5.8	5.6
Pipe Tobacco ²	3.1	3.5	--	--	0.4	0.6
ALCOHOL	85.2	85.0	76.3	75.7	56.0	55.4
Binge Alcohol Use ³	--	--	--	--	32.3	31.9
Heavy Alcohol Use ³	--	--	--	--	8.8	8.8

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.