

**Table 2.72A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	171,827	172,217	83,066	84,958	70,257	71,519
<b>GEOGRAPHIC DIVISION</b>						
Northeast	31,951	32,504	15,180	15,530	12,642	12,941
New England	8,826	8,820	3,793	3,969	3,206	3,295
Middle Atlantic	23,125	23,685	11,387	11,560	9,437	9,646
Midwest	40,761	40,586	20,423	21,077	17,549	18,012
East North Central	28,399	27,974	14,271	14,557	12,273	12,518
West North Central	12,362	12,612	6,152	6,519	5,275	5,494
South	61,896	61,944	31,501	31,471	27,359	27,032
South Atlantic	32,981	32,377	16,439	15,469	14,219	13,354
East South Central	10,384 <sup>b</sup>	10,988	5,471	6,048	4,873	5,265
West South Central	18,531	18,579	9,591	9,954	8,267	8,413
West	37,218	37,184	15,962	16,880	12,707	13,534
Mountain	11,547	11,797	5,308	5,440	4,366	4,481
Pacific	25,672	25,387	10,654	11,441	8,341	9,053
<b>COUNTY TYPE</b>						
Large Metro	88,589	90,327	41,730	43,108	34,379	35,562
Small Metro	52,580	51,563	25,815	25,562	22,211	21,883
250K – 1 Mil. Pop.	33,853	34,347	16,377	16,769	13,895	14,262
< 250K Pop.	18,726	17,216	9,438	8,793	8,316	7,620
Nonmetro	30,658	30,328	15,521	16,288	13,666	14,074
Urbanized	12,518	11,354	6,544	6,180	5,652	5,358
Less Urbanized	15,418	15,444	7,812	8,442	6,980	7,352
Completely Rural	2,723	3,530	1,165 <sup>a</sup>	1,665	1,033	1,364

\*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 2.72B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	71.4	70.8	34.5	34.9	29.2	29.4
<b>GEOGRAPHIC DIVISION</b>						
Northeast	70.2	71.2	33.4	34.0	27.8	28.4
New England	73.9	73.7	31.7	33.2	26.8	27.5
Middle Atlantic	68.9	70.4	33.9	34.3	28.1	28.7
Midwest	75.2	74.4	37.7	38.7	32.4	33.0
East North Central	74.9	73.4	37.6	38.2	32.4	32.8
West North Central	76.0	76.8	37.8	39.7	32.4	33.5
South	71.9	70.7	36.6	35.9	31.8	30.9
South Atlantic	73.0 <sup>a</sup>	70.2	36.4 <sup>a</sup>	33.6	31.5	29.0
East South Central	72.1 <sup>a</sup>	75.6	38.0	41.6	33.9	36.2
West South Central	69.8	68.9	36.1	36.9	31.1	31.2
West	68.1	67.0	29.2	30.4	23.2	24.4
Mountain	72.2	71.8	33.2	33.1	27.3	27.3
Pacific	66.4	65.1	27.5	29.3	21.6	23.2
<b>COUNTY TYPE</b>						
Large Metro	69.0	68.9	32.5	32.9	26.8	27.1
Small Metro	73.4	72.0	36.0	35.7	31.0	30.6
250K – 1 Mil. Pop.	71.8	72.0	34.7	35.1	29.5	29.9
< 250K Pop.	76.5 <sup>b</sup>	72.0	38.6	36.8	34.0	31.9
Nonmetro	75.6	74.8	38.3	40.2	33.7	34.7
Urbanized	75.0	73.8	39.2	40.2	33.9	34.8
Less Urbanized	76.0	75.7	38.5	41.4	34.4	36.0
Completely Rural	76.3	74.1	32.7	35.0	29.0	28.6

\*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 2.73A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	8,249 <sup>b</sup>	7,795	5,562	5,365	3,635 <sup>b</sup>	3,324
<b>GEOGRAPHIC DIVISION</b>						
Northeast	1,419	1,369	1,052	977	660	623
New England	380	351	294	258	183	164
Middle Atlantic	1,039	1,018	758	719	477	459
Midwest	1,959 <sup>a</sup>	1,814	1,311	1,255	882	821
East North Central	1,345 <sup>a</sup>	1,246	901	845	592	542
West North Central	614	568	410	410	290	280
South	3,046	2,972	2,009	1,974	1,314	1,225
South Atlantic	1,513	1,460	974	972	645	620
East South Central	557	550	399	380	284	238
West South Central	976	962	636	622	385	366
West	1,825 <sup>a</sup>	1,640	1,190	1,158	779 <sup>a</sup>	656
Mountain	567	597	372	436	247	245
Pacific	1,258 <sup>b</sup>	1,043	818	722	532 <sup>a</sup>	411
<b>COUNTY TYPE</b>						
Large Metro	3,869	3,752	2,560	2,595	1,600	1,584
Small Metro	2,668 <sup>b</sup>	2,360	1,849 <sup>b</sup>	1,602	1,220 <sup>b</sup>	988
250K – 1 Mil. Pop.	1,735 <sup>a</sup>	1,552	1,214 <sup>a</sup>	1,064	783 <sup>a</sup>	660
< 250K Pop.	932	807	635	539	437 <sup>b</sup>	328
Nonmetro	1,712	1,683	1,153	1,167	815	752
Urbanized	628	577	429	393	305	258
Less Urbanized	938	883	637	618	450	395
Completely Rural	145 <sup>a</sup>	223	86 <sup>a</sup>	156	60	99

\*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 2.73B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	32.7 <sup>b</sup>	30.7	22.1	21.2	14.4 <sup>b</sup>	13.1
<b>GEOGRAPHIC DIVISION</b>						
Northeast	31.3	30.1	23.2	21.5	14.6	13.7
New England	32.4	29.9	25.1	22.0	15.6	13.9
Middle Atlantic	30.9	30.2	22.5	21.3	14.2	13.6
Midwest	34.5 <sup>a</sup>	32.0	23.1	22.2	15.5	14.5
East North Central	33.6 <sup>a</sup>	31.0	22.5	21.1	14.8	13.5
West North Central	36.7	34.4	24.5	24.8	17.3	16.9
South	33.8	32.8	22.3	21.8	14.6	13.5
South Atlantic	33.0	31.5	21.3	21.0	14.1	13.4
East South Central	38.5	37.9	27.5	26.2	19.6	16.4
West South Central	32.8	32.3	21.4	20.9	12.9	12.3
West	30.4 <sup>a</sup>	27.0	19.8	19.1	13.0 <sup>a</sup>	10.8
Mountain	32.5	33.9	21.3	24.8	14.2	13.9
Pacific	29.6 <sup>b</sup>	24.2	19.2	16.7	12.5 <sup>a</sup>	9.5
<b>COUNTY TYPE</b>						
Large Metro	29.0	27.5	19.2	19.0	12.0	11.6
Small Metro	34.9 <sup>b</sup>	31.6	24.2 <sup>b</sup>	21.5	16.0 <sup>b</sup>	13.2
250K – 1 Mil. Pop.	34.0 <sup>b</sup>	30.4	23.8 <sup>b</sup>	20.9	15.3 <sup>b</sup>	12.9
< 250K Pop.	36.7	34.2	25.0	22.8	17.2 <sup>a</sup>	13.9
Nonmetro	40.5	39.8	27.3	27.6	19.3	17.8
Urbanized	36.9	36.5	25.3	24.9	17.9	16.3
Less Urbanized	43.7	41.2	29.7	28.9	21.0	18.4
Completely Rural	38.5	44.3	22.9	31.0	16.0	19.7

\*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 2.74A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	23,658	23,496	17,481	17,637	14,359	14,382
<b>GEOGRAPHIC DIVISION</b>						
Northeast	4,206	4,087	3,208	3,079	2,645	2,545
New England	1,126	1,092	871	842	729	687
Middle Atlantic	3,080	2,995	2,337	2,237	1,916	1,859
Midwest	5,910 <sup>a</sup>	5,722	4,506	4,400	3,779 <sup>a</sup>	3,597
East North Central	4,007	3,897	3,068	2,995	2,582	2,450
West North Central	1,903	1,825	1,438	1,405	1,197	1,147
South	8,337	8,413	6,137	6,348	5,132	5,315
South Atlantic	4,164	4,132	3,065	3,102	2,550	2,561
East South Central	1,439	1,465	1,109	1,115	946	969
West South Central	2,734	2,815	1,963 <sup>a</sup>	2,130	1,635	1,785
West	5,204	5,275	3,631	3,810	2,803	2,924
Mountain	1,664	1,696	1,187	1,204	963	966
Pacific	3,540	3,579	2,444	2,607	1,840	1,958
<b>COUNTY TYPE</b>						
Large Metro	11,826	12,003	8,565	8,954	6,906	7,183
Small Metro	7,931	7,616	5,907	5,707	4,856	4,693
250K – 1 Mil. Pop.	5,024	4,892	3,722	3,649	3,015	2,988
< 250K Pop.	2,907	2,725	2,186	2,058	1,841	1,705
Nonmetro	3,901	3,877	3,009	2,975	2,597	2,506
Urbanized	1,798	1,636	1,415	1,224	1,214 <sup>a</sup>	1,019
Less Urbanized	1,827	1,846	1,376	1,440	1,193	1,236
Completely Rural	276 <sup>a</sup>	395	217	311	190	251

\*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 2.74B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	73.5	72.3	54.3	54.3	44.6	44.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	74.0	71.6	56.4	53.9	46.5	44.6
New England	75.7	73.1	58.5	56.3	49.0	46.0
Middle Atlantic	73.4	71.1	55.7	53.1	45.7	44.1
Midwest	79.4 <sup>b</sup>	76.8	60.6	59.1	50.8 <sup>a</sup>	48.3
East North Central	78.2	76.1	59.8	58.5	50.4	47.8
West North Central	82.2 <sup>a</sup>	78.4	62.1	60.4	51.7	49.3
South	72.3	71.9	53.2	54.2	44.5	45.4
South Atlantic	73.0	70.9	53.7	53.2	44.7	43.9
East South Central	73.8	75.2	56.8	57.2	48.5	49.7
West South Central	70.4	71.7	50.6	54.3	42.1	45.5
West	69.1	69.2	48.2	50.0	37.2	38.4
Mountain	72.7	72.3	51.9	51.3	42.1	41.2
Pacific	67.5	67.8	46.6	49.4	35.1	37.1
<b>COUNTY TYPE</b>						
Large Metro	71.0	69.7	51.4	52.0	41.5	41.7
Small Metro	74.8	73.8	55.7	55.3	45.8	45.5
250K – 1 Mil. Pop.	72.9	72.2	54.0	53.8	43.8	44.1
< 250K Pop.	78.2	76.9	58.8	58.1	49.5	48.1
Nonmetro	79.0	78.3	60.9	60.1	52.6	50.6
Urbanized	78.1	77.7	61.5	58.1	52.7	48.4
Less Urbanized	79.0	78.4	59.5	61.1	51.6	52.5
Completely Rural	85.0	80.5	67.0	63.3	58.7	51.0

\*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 2.75A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	139,921	140,926	60,023	61,957	52,263	53,813
<b>GEOGRAPHIC DIVISION</b>						
Northeast	26,326	27,048	10,921	11,474	9,337	9,773
New England	7,319	7,377	2,627	2,869	2,293	2,445
Middle Atlantic	19,006	19,671	8,293	8,605	7,044	7,329
Midwest	32,892	33,050	14,606	15,422	12,888	13,594
East North Central	23,047	22,831	10,302	10,717	9,099	9,526
West North Central	9,845	10,219	4,304	4,704	3,789	4,068
South	50,514	50,559	23,355	23,149	20,913	20,492
South Atlantic	27,305	26,784	12,400	11,395	11,023	10,172
East South Central	8,387 <sup>b</sup>	8,973	3,963 <sup>a</sup>	4,553	3,643	4,057
West South Central	14,821	14,802	6,992	7,201	6,247	6,262
West	30,189	30,269	11,142	11,912	9,124	9,954
Mountain	9,316	9,505	3,749	3,800	3,155	3,269
Pacific	20,874	20,764	7,392	8,112	5,969	6,684
<b>COUNTY TYPE</b>						
Large Metro	72,894	74,571	30,606	31,559	25,874	26,795
Small Metro	41,981	41,587	18,058	18,252	16,135	16,202
250K – 1 Mil. Pop.	27,094	27,903	11,441	12,056	10,097	10,614
< 250K Pop.	14,887	13,683	6,617	6,196	6,038	5,588
Nonmetro	25,046	24,768	11,359	12,146	10,254	10,816
Urbanized	10,092	9,141	4,700	4,563	4,134	4,081
Less Urbanized	12,653	12,715	5,799	6,384	5,338	5,721
Completely Rural	2,302	2,912	861	1,198	783	1,014

\*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 2.75B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	76.4	76.0	32.8	33.4	28.5	29.0
<b>GEOGRAPHIC DIVISION</b>						
Northeast	74.6	76.5	31.0	32.4	26.5	27.6
New England	78.8	79.4	28.3	30.9	24.7	26.3
Middle Atlantic	73.1	75.4	31.9	33.0	27.1	28.1
Midwest	80.1	79.8	35.5	37.2	31.4	32.8
East North Central	80.0	78.8	35.8	37.0	31.6	32.9
West North Central	80.1	82.2	35.0	37.8	30.8	32.7
South	77.0	75.7	35.6	34.6	31.9	30.7
South Atlantic	78.2 <sup>a</sup>	75.2	35.5 <sup>a</sup>	32.0	31.6	28.5
East South Central	76.3 <sup>a</sup>	80.6	36.1	40.9	33.1	36.4
West South Central	75.2	73.8	35.5	35.9	31.7	31.2
West	73.4	72.5	27.1	28.5	22.2	23.8
Mountain	77.9	77.1	31.4	30.8	26.4	26.5
Pacific	71.5	70.6	25.3	27.6	20.5	22.7
<b>COUNTY TYPE</b>						
Large Metro	74.1	74.4	31.1	31.5	26.3	26.7
Small Metro	78.6	77.3	33.8	33.9	30.2	30.1
250K – 1 Mil. Pop.	77.0	77.9	32.5	33.6	28.7	29.6
< 250K Pop.	81.7 <sup>b</sup>	76.0	36.3	34.4	33.2	31.0
Nonmetro	79.8	79.0	36.2	38.7	32.7	34.5
Urbanized	79.5	78.2	37.0	39.0	32.6	34.9
Less Urbanized	79.9	79.9	36.6	40.1	33.7	36.0
Completely Rural	80.3	77.2	30.0	31.8	27.3	26.9

\*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.



**Table 2.76A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	163,579	164,422	77,504	79,593	66,622	68,195
<b>GEOGRAPHIC DIVISION</b>						
Northeast	30,532	31,135	14,128	14,553	11,982	12,318
New England	8,446	8,469	3,498	3,711	3,022	3,131
Middle Atlantic	22,087	22,666	10,630	10,842	8,960	9,187
Midwest	38,803	38,772	19,112	19,822	16,667	17,191
East North Central	27,054	26,728	13,370	13,712	11,681	11,976
West North Central	11,748	12,044	5,742	6,109	4,986	5,215
South	58,850	58,972	29,492	29,497	26,045	25,807
South Atlantic	31,469	30,917	15,465	14,497	13,574	12,733
East South Central	9,827 <sup>b</sup>	10,438	5,072 <sup>a</sup>	5,668	4,589	5,027
West South Central	17,555	17,617	8,954	9,332	7,882	8,047
West	35,394	35,544	14,773	15,722	11,928	12,878
Mountain	10,980	11,200	4,936	5,004	4,118	4,236
Pacific	24,414	24,344	9,837	10,718	7,809	8,642
<b>COUNTY TYPE</b>						
Large Metro	84,720	86,574	39,170	40,513	32,780	33,978
Small Metro	49,912	49,203	23,966	23,960	20,991	20,895
250K – 1 Mil. Pop.	32,118	32,795	15,162	15,705	13,112	13,602
< 250K Pop.	17,794	16,408	8,803	8,255	7,879	7,293
Nonmetro	28,947	28,645	14,368	15,121	12,851	13,322
Urbanized	11,890	10,777	6,115	5,787	5,348	5,100
Less Urbanized	14,480	14,561	7,175	7,824	6,530	6,958
Completely Rural	2,577	3,307	1,078	1,509	973	1,265

\*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 2.76B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2004 and 2005**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	76.0	75.5	36.0	36.5	30.9	31.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	74.5	75.8	34.5	35.4	29.3	30.0
New England	78.4	78.5	32.5	34.4	28.1	29.0
Middle Atlantic	73.2	74.8	35.2	35.8	29.7	30.3
Midwest	80.0	79.4	39.4	40.6	34.3	35.2
East North Central	79.7	78.4	39.4	40.2	34.4	35.1
West North Central	80.5	81.6	39.3	41.4	34.1	35.3
South	76.3	75.1	38.2	37.6	33.8	32.9
South Atlantic	77.5 <sup>a</sup>	74.6	38.1 <sup>a</sup>	35.0	33.4	30.7
East South Central	75.9 <sup>a</sup>	79.8	39.2	43.3	35.5	38.4
West South Central	74.4	73.5	38.0	38.9	33.4	33.6
West	72.7	72.0	30.4	31.8	24.5	26.1
Mountain	77.1	76.3	34.7	34.1	28.9	28.9
Pacific	70.9	70.1	28.6	30.9	22.7	24.9
<b>COUNTY TYPE</b>						
Large Metro	73.7	73.7	34.1	34.5	28.5	28.9
Small Metro	78.0	76.7	37.5	37.4	32.8	32.6
250K – 1 Mil. Pop.	76.4	77.0	36.0	36.9	31.2	31.9
< 250K Pop.	81.1 <sup>b</sup>	76.2	40.1	38.3	35.9	33.9
Nonmetro	79.7	78.9	39.5	41.6	35.4	36.7
Urbanized	79.3	78.1	40.8	41.9	35.7	37.0
Less Urbanized	79.8	79.7	39.5	42.9	36.0	38.1
Completely Rural	80.8	77.6	33.8	35.4	30.5	29.7

\*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.