

Table 2.103C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	272	286	246	268	191	208
GENDER						
Male	177	184	162	171	130	137
Female	170	180	155	171	119	128
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	243	261	220	243	171	189
White	210	232	195	219	154	175
Black or African American	100	106	83	86	56	58
American Indian or Alaska Native	15	15	12	14	7	8
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	4
Asian	49	52	44	47	27	29
Two or More Races	29	23	26	22	17	15
Hispanic or Latino	112	123	96	109	76	79
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	136	148	126	140	104	115
Female, White, Not Hispanic	133	141	124	134	94	106
Male, Black, Not Hispanic	66	65	56	54	38	40
Female, Black, Not Hispanic	59	65	52	55	36	36
Male, Hispanic	77	87	69	76	54	59
Female, Hispanic	74	77	62	72	51	51

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.103D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Percentages, 2004 and 2005

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.41	0.41	0.41	0.43	0.39	0.41
GENDER						
Male	0.56	0.55	0.56	0.56	0.55	0.53
Female	0.55	0.56	0.55	0.58	0.51	0.54
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.43	0.44	0.44	0.45	0.43	0.44
White	0.49	0.50	0.52	0.53	0.50	0.52
Black or African American	1.06	1.02	0.98	0.96	0.81	0.83
American Indian or Alaska Native	4.44	3.72	3.81	3.53	2.92	2.85
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	3.54
Asian	2.15	2.40	2.01	2.29	1.57	1.59
Two or More Races	2.59	2.79	2.75	2.81	2.28	2.24
Hispanic or Latino	1.02	1.06	0.98	1.06	0.97	0.97
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.68	0.67	0.68	0.71	0.68	0.69
Female, White, Not Hispanic	0.67	0.64	0.70	0.67	0.65	0.68
Male, Black, Not Hispanic	1.50	1.43	1.38	1.34	1.10	1.21
Female, Black, Not Hispanic	1.33	1.38	1.28	1.35	1.10	1.03
Male, Hispanic	1.52	1.52	1.48	1.48	1.41	1.37
Female, Hispanic	1.44	1.51	1.40	1.59	1.35	1.33

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.103P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.1074		0.6221		0.3878	
GENDER						
Male	0.1109		0.4361		0.3830	
Female	0.4328		0.9557		0.6856	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1165		0.4518		0.4811	
White	0.0624		0.4507		0.6753	
Black or African American	0.5003		0.7987		0.9587	
American Indian or Alaska Native	0.8276		0.4508		0.5191	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.6554		0.6550		0.6782	
Two or More Races	0.3665		0.9440		0.4685	
Hispanic or Latino	0.6058		0.6435		0.6175	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.2795		0.4471		0.3338	
Female, White, Not Hispanic	0.0816		0.7037		0.6946	
Male, Black, Not Hispanic	0.9642		0.9513		0.9700	
Female, Black, Not Hispanic	0.2663		0.7401		0.9324	
Male, Hispanic	0.6672		0.5430		0.6176	
Female, Hispanic	0.7812		0.9911		0.1909	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.104C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	1,988	1,904	1,688	1,662	1,513	1,452
GENDER						
Male	1,279	1,263	1,116	1,109	1,008	1,006
Female	1,188	1,196	1,011	1,025	913	856
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	1,891	1,820	1,612	1,590	1,439	1,399
White	1,666	1,666	1,440	1,456	1,304	1,291
Black or African American	643	654	491	520	414	451
American Indian or Alaska Native	138	*	*	82	58	68
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	389	358	355	325	289	276
Two or More Races	172	154	158	133	126	111
Hispanic or Latino	648	658	564	569	424	448
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	1,025	1,067	909	949	832	877
Female, White, Not Hispanic	1,047	1,063	917	922	841	780
Male, Black, Not Hispanic	385	449	306	352	277	323
Female, Black, Not Hispanic	421	403	310	325	246	260
Male, Hispanic	469	493	410	427	320	362
Female, Hispanic	345	368	299	317	239	223

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.104D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2004 and 2005

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.33	0.30	0.43	0.44	0.46	0.47
GENDER						
Male	0.42	0.32	0.57	0.57	0.63	0.62
Female	0.47	0.47	0.60	0.58	0.62	0.59
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.35	0.32	0.47	0.47	0.50	0.50
White	0.33	0.30	0.50	0.48	0.54	0.53
Black or African American	1.13	1.12	1.24	1.44	1.34	1.41
American Indian or Alaska Native	3.13	*	*	5.17	5.89	5.15
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	3.05	2.42	3.18	2.60	2.73	2.45
Two or More Races	1.54	2.76	2.94	3.74	3.32	3.70
Hispanic or Latino	1.14	1.01	1.29	1.23	1.15	1.28
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.42	0.30	0.64	0.61	0.71	0.67
Female, White, Not Hispanic	0.49	0.48	0.68	0.66	0.74	0.70
Male, Black, Not Hispanic	1.66	1.23	1.90	2.27	1.94	2.26
Female, Black, Not Hispanic	1.53	1.63	1.74	1.73	1.64	1.56
Male, Hispanic	1.29	1.10	1.78	1.66	1.70	1.78
Female, Hispanic	1.59	1.54	1.66	1.72	1.60	1.55

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.104P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.0785		0.0086		0.0034	
GENDER						
Male	0.1605		0.4337		0.0881	
Female	0.2106		0.0026		0.0074	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.3398		0.0867		0.0110	
White	0.2736		0.2083		0.0437	
Black or African American	0.1819		0.0307		0.0198	
American Indian or Alaska Native	*		*		0.2749	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4372		0.9346		0.7669	
Two or More Races	0.0732		0.0496		0.2022	
Hispanic or Latino	0.0237		0.0033		0.0639	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.3381		0.9307		0.4453	
Female, White, Not Hispanic	0.4845		0.0668		0.0296	
Male, Black, Not Hispanic	0.2135		0.4016		0.1292	
Female, Black, Not Hispanic	0.4538		0.0314		0.0523	
Male, Hispanic	0.0490		0.0171		0.0409	
Female, Hispanic	0.1123		0.0523		0.5794	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.105C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

Demographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2004	2005	2004	2005	2004	2005
TOTAL	191	208	160	170	98	86
GENDER						
Male	130	137	115	123	70	67
Female	119	128	93	92	50	44
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	171	189	143	152	91	78
White	154	175	130	142	82	76
Black or African American	56	58	37	37	17	15
American Indian or Alaska Native	7	8	5	8	4	5
Native Hawaiian or Other Pacific Islander	*	4	*	3	1	1
Asian	27	29	22	18	9	5
Two or More Races	17	15	14	13	10	9
Hispanic or Latino	76	79	64	66	32	32
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	104	115	93	101	58	58
Female, White, Not Hispanic	94	106	76	80	45	40
Male, Black, Not Hispanic	38	40	29	30	16	13
Female, Black, Not Hispanic	36	36	21	20	5	8
Male, Hispanic	54	59	48	53	27	27
Female, Hispanic	51	51	40	37	18	16

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.105D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Percentages, 2004 and 2005

Demographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2004	2005	2004	2005	2004	2005
TOTAL	0.39	0.41	0.36	0.37	0.24	0.21
GENDER						
Male	0.55	0.53	0.52	0.51	0.34	0.32
Female	0.51	0.54	0.44	0.43	0.26	0.22
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.43	0.44	0.39	0.40	0.27	0.23
White	0.50	0.52	0.47	0.47	0.33	0.29
Black or African American	0.81	0.83	0.58	0.60	0.30	0.26
American Indian or Alaska Native	2.92	2.85	2.31	2.60	1.75	1.68
Native Hawaiian or Other Pacific Islander	*	3.54	*	2.65	0.65	0.90
Asian	1.57	1.59	1.35	1.06	0.60	0.34
Two or More Races	2.28	2.24	2.00	2.06	1.49	1.46
Hispanic or Latino	0.97	0.97	0.88	0.86	0.49	0.46
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.68	0.69	0.64	0.65	0.45	0.43
Female, White, Not Hispanic	0.65	0.68	0.58	0.57	0.37	0.32
Male, Black, Not Hispanic	1.10	1.21	0.94	0.99	0.55	0.45
Female, Black, Not Hispanic	1.10	1.03	0.71	0.65	0.16	0.27
Male, Hispanic	1.41	1.37	1.30	1.31	0.78	0.76
Female, Hispanic	1.35	1.33	1.16	1.01	0.58	0.47

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.105P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Demographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2004	2005	2004	2005	2004	2005
TOTAL	0.3878		0.1124		0.3447	
GENDER						
Male	0.3830		0.3121		0.2350	
Female	0.6856		0.1486		0.9377	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4811		0.2182		0.4745	
White	0.6753		0.4017		0.5330	
Black or African American	0.9587		0.3169		0.5400	
American Indian or Alaska Native	0.5191		0.8129		0.5831	
Native Hawaiian or Other Pacific Islander	*		*		0.7930	
Asian	0.6782		0.7296		0.5522	
Two or More Races	0.4685		0.6298		0.6742	
Hispanic or Latino	0.6175		0.2356		0.4777	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.3338		0.3094		0.3496	
Female, White, Not Hispanic	0.6946		0.8799		0.8659	
Male, Black, Not Hispanic	0.9700		0.8472		0.2045	
Female, Black, Not Hispanic	0.9324		0.1625		0.1297	
Male, Hispanic	0.6176		0.7949		0.7448	
Female, Hispanic	0.1909		0.0213		0.4098	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.106C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

Demographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2004	2005	2004	2005	2004	2005
TOTAL	1,513	1,452	787	795	384	379
GENDER						
Male	1,008	1,006	632	629	329	330
Female	913	856	393	413	165	157
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	1,439	1,399	724	741	358	355
White	1,304	1,291	646	669	337	332
Black or African American	414	451	253	281	113	116
American Indian or Alaska Native	58	68	49	62	34	46
Native Hawaiian or Other Pacific Islander	*	*	*	*	11	11
Asian	289	276	127	117	53	39
Two or More Races	126	111	58	61	28	24
Hispanic or Latino	424	448	286	293	125	133
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	832	877	513	525	286	290
Female, White, Not Hispanic	841	780	333	344	159	137
Male, Black, Not Hispanic	277	323	188	209	101	102
Female, Black, Not Hispanic	246	260	145	171	52	52
Male, Hispanic	320	362	253	255	113	125
Female, Hispanic	239	223	127	128	43	42

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.106D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2004 and 2005

Demographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2004	2005	2004	2005	2004	2005
TOTAL	0.46	0.47	0.34	0.33	0.18	0.18
GENDER						
Male	0.63	0.62	0.52	0.53	0.32	0.32
Female	0.62	0.59	0.36	0.37	0.16	0.15
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.50	0.50	0.36	0.35	0.20	0.19
White	0.54	0.53	0.41	0.38	0.23	0.22
Black or African American	1.34	1.41	1.01	1.06	0.48	0.49
American Indian or Alaska Native	5.89	5.15	4.93	4.86	3.13	4.05
Native Hawaiian or Other Pacific Islander	*	*	*	*	2.26	1.92
Asian	2.73	2.45	1.32	1.36	0.62	0.45
Two or More Races	3.32	3.70	2.98	2.80	1.43	1.18
Hispanic or Latino	1.15	1.28	0.97	1.00	0.48	0.52
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.71	0.67	0.63	0.59	0.40	0.38
Female, White, Not Hispanic	0.74	0.70	0.43	0.42	0.22	0.18
Male, Black, Not Hispanic	1.94	2.26	1.63	1.80	0.95	0.99
Female, Black, Not Hispanic	1.64	1.56	1.11	1.17	0.41	0.41
Male, Hispanic	1.70	1.78	1.58	1.60	0.83	0.93
Female, Hispanic	1.60	1.55	1.04	0.99	0.35	0.34

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.106P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Demographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2004	2005	2004	2005	2004	2005
TOTAL	0.0034		0.9425		0.1654	
GENDER						
Male	0.0881		0.5139		0.5564	
Female	0.0074		0.3562		0.0380	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0110		0.9485		0.0811	
White	0.0437		0.4497		0.1033	
Black or African American	0.0198		0.0623		0.8234	
American Indian or Alaska Native	0.2749		0.1967		0.3882	
Native Hawaiian or Other Pacific Islander	*		*		0.9078	
Asian	0.7669		0.7681		0.3454	
Two or More Races	0.2022		0.4301		0.2866	
Hispanic or Latino	0.0639		1.0000		0.4434	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.4453		0.3108		0.4852	
Female, White, Not Hispanic	0.0296		0.9501		0.0202	
Male, Black, Not Hispanic	0.1292		0.2991		0.6708	
Female, Black, Not Hispanic	0.0523		0.0875		0.8149	
Male, Hispanic	0.0409		0.7341		0.5236	
Female, Hispanic	0.5794		0.5826		0.5879	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.107C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	216	197	190	182	146	141
NOT HISPANIC OR LATINO¹	196	178	174	164	136	127
White	166	156	151	147	121	117
Black or African American	80	73	65	60	41	40
American Indian or Alaska Native	18	11	16	9	12	5
Native Hawaiian or Other Pacific Islander	12	11	11	8	8	5
Chinese	17	18	13	17	8	9
Filipino	13	14	11	12	8	8
Japanese	*	*	*	*	3	*
Asian Indian	11	12	10	12	9	6
Korean	11	13	11	12	8	8
Vietnamese	17	12	15	9	9	6
HISPANIC OR LATINO¹	90	83	71	73	48	55
Mexican	79	72	61	62	40	46
Puerto Rican	28	22	24	20	15	14
Central or South American	26	24	22	21	17	18
Cuban	12	12	11	11	9	8

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.107D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.31	0.29	0.32	0.30	0.31	0.28
NOT HISPANIC OR LATINO¹	0.34	0.31	0.35	0.31	0.34	0.31
White	0.37	0.35	0.37	0.37	0.37	0.36
Black or African American	0.77	0.74	0.72	0.69	0.58	0.58
American Indian or Alaska Native	3.01	2.89	3.01	2.59	2.84	2.03
Native Hawaiian or Other Pacific Islander	4.71	5.30	5.26	5.20	4.38	3.92
Chinese	4.12	3.79	3.54	3.80	2.15	2.68
Filipino	3.00	4.23	2.99	3.90	2.23	2.79
Japanese	*	*	*	*	4.01	*
Asian Indian	3.30	3.05	3.12	3.02	2.86	1.77
Korean	4.76	4.82	4.82	4.74	4.23	3.98
Vietnamese	4.87	5.89	5.12	5.12	4.10	3.51
HISPANIC OR LATINO¹	0.75	0.73	0.78	0.72	0.64	0.69
Mexican	0.92	0.91	0.95	0.90	0.76	0.85
Puerto Rican	2.13	2.06	2.23	2.07	1.77	1.76
Central or South American	2.30	2.07	2.32	2.25	2.19	2.12
Cuban	3.58	3.65	3.83	3.47	4.36	3.21

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.107P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.0002		0.2456		0.3104	
NOT HISPANIC OR LATINO¹	0.0006		0.2051		0.1485	
White	0.0117		0.5039		0.2707	
Black or African American	0.0381		0.4225		0.7255	
American Indian or Alaska Native	0.0580		0.1158		0.0709	
Native Hawaiian or Other Pacific Islander	0.7790		0.4456		0.0587	
Chinese	0.0740		0.0333		0.3008	
Filipino	0.0528		0.2955		0.1245	
Japanese	*		*		*	
Asian Indian	0.2703		0.3922		0.1827	
Korean	0.5741		0.9760		0.9529	
Vietnamese	0.2617		0.1969		0.4034	
HISPANIC OR LATINO¹	0.0855		0.9551		0.3449	
Mexican	0.1314		0.6692		0.1539	
Puerto Rican	0.0722		0.1731		0.1814	
Central or South American	0.6428		0.8451		0.6153	
Cuban	0.1723		0.4053		0.2883	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.108C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	1,553	1,376	1,333	1,184	1,156	1,048
NOT HISPANIC OR LATINO¹	1,492	1,312	1,286	1,132	1,107	1,003
White	1,344	1,178	1,171	1,024	1,029	918
Black or African American	469	458	387	357	311	306
American Indian or Alaska Native	75	85	63	56	59	45
Native Hawaiian or Other Pacific Islander	*	*	41	43	40	35
Chinese	126	148	111	140	93	118
Filipino	156	125	125	111	108	90
Japanese	*	*	*	*	63	*
Asian Indian	118	101	107	86	89	67
Korean	*	*	*	46	54	41
Vietnamese	*	57	*	51	*	45
HISPANIC OR LATINO¹	502	462	437	401	357	308
Mexican	382	357	331	313	266	238
Puerto Rican	144	147	124	135	96	101
Central or South American	159	177	149	143	121	116
Cuban	131	88	94	78	78	71

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.108D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.22	0.23	0.32	0.31	0.34	0.33
NOT HISPANIC OR LATINO¹	0.23	0.24	0.35	0.33	0.36	0.35
White	0.21	0.22	0.37	0.35	0.40	0.38
Black or African American	0.74	0.79	0.97	0.95	0.87	0.97
American Indian or Alaska Native	3.34	2.62	3.57	4.85	4.22	4.04
Native Hawaiian or Other Pacific Islander	*	*	5.67	5.92	5.52	5.23
Chinese	3.02	4.20	4.00	4.38	3.78	4.21
Filipino	3.51	4.35	4.40	4.63	4.08	4.01
Japanese	*	*	*	*	6.03	*
Asian Indian	3.54	3.26	3.73	3.24	3.46	2.73
Korean	*	*	*	5.44	5.99	5.37
Vietnamese	*	5.78	*	5.96	*	5.53
HISPANIC OR LATINO¹	0.75	0.76	1.02	0.89	0.99	0.86
Mexican	0.94	0.93	1.23	1.13	1.16	1.05
Puerto Rican	2.28	2.19	3.03	2.81	2.99	2.90
Central or South American	2.04	2.05	2.57	2.39	2.53	2.43
Cuban	2.58	2.61	3.48	3.89	3.82	3.46

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.108P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.3989		0.4233		0.1364	
NOT HISPANIC OR LATINO¹	0.6265		0.2557		0.0432	
White	0.4789		0.0535		0.0103	
Black or African American	0.1655		0.7328		1.0000	
American Indian or Alaska Native	0.4283		0.2816		0.5321	
Native Hawaiian or Other Pacific Islander	*		0.5339		0.6065	
Chinese	0.3440		0.9905		0.9932	
Filipino	0.6488		0.7259		0.6306	
Japanese	*		*		*	
Asian Indian	0.5440		0.0674		0.0215	
Korean	*		*		0.7973	
Vietnamese	*		*		*	
HISPANIC OR LATINO¹	0.5784		0.6374		0.3818	
Mexican	0.5196		0.3391		0.1786	
Puerto Rican	0.9291		0.8169		0.7361	
Central or South American	0.4912		0.6697		0.3547	
Cuban	0.4564		0.0706		0.1689	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 2.109C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Racial/Ethnic Subgroups:
Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005**

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	146	141	119	117	57	65
NOT HISPANIC OR LATINO¹	136	127	109	104	55	60
White	121	117	100	96	53	56
Black or African American	41	40	27	26	10	11
American Indian or Alaska Native	12	5	7	5	2	3
Native Hawaiian or Other Pacific Islander	8	5	7	4	2	1
Chinese	8	9	6	6	3	3
Filipino	8	8	6	5	2	2
Japanese	3	*	3	*	*	1
Asian Indian	9	6	5	6	2	2
Korean	8	8	5	6	*	2
Vietnamese	9	6	5	3	*	1
HISPANIC OR LATINO¹	48	55	39	46	18	23
Mexican	40	46	32	39	16	19
Puerto Rican	15	14	12	12	5	6
Central or South American	17	18	14	16	5	6
Cuban	9	8	8	6	3	4

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.109D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.31	0.28	0.27	0.26	0.15	0.16
NOT HISPANIC OR LATINO¹	0.34	0.31	0.29	0.28	0.17	0.18
White	0.37	0.36	0.34	0.33	0.21	0.22
Black or African American	0.58	0.58	0.44	0.42	0.18	0.20
American Indian or Alaska Native	2.84	2.03	2.19	1.74	0.84	1.22
Native Hawaiian or Other Pacific Islander	4.38	3.92	4.04	3.30	1.50	0.56
Chinese	2.15	2.68	1.87	1.85	1.09	0.91
Filipino	2.23	2.79	2.00	1.94	0.88	0.77
Japanese	4.01	*	3.62	*	*	1.60
Asian Indian	2.86	1.77	1.78	1.53	0.76	0.43
Korean	4.23	3.98	3.23	2.95	*	0.82
Vietnamese	4.10	3.51	2.27	1.80	*	0.58
HISPANIC OR LATINO¹	0.64	0.69	0.59	0.61	0.29	0.34
Mexican	0.76	0.85	0.69	0.75	0.37	0.40
Puerto Rican	1.77	1.76	1.48	1.57	0.70	0.81
Central or South American	2.19	2.12	1.89	1.99	0.67	0.89
Cuban	4.36	3.21	4.34	2.44	1.73	1.86

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.109P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.3104		0.9232		0.9537	
NOT HISPANIC OR LATINO¹	0.1485		0.3522		0.8388	
White	0.2707		0.6777		1.0000	
Black or African American	0.7255		0.8913		0.6684	
American Indian or Alaska Native	0.0709		0.2559		0.1739	
Native Hawaiian or Other Pacific Islander	0.0587		0.0593		0.1407	
Chinese	0.3008		0.7186		0.7322	
Filipino	0.1245		0.3810		0.9855	
Japanese	*		*		*	
Asian Indian	0.1827		0.6548		0.4263	
Korean	0.9529		0.9407		*	
Vietnamese	0.4034		0.5891		*	
HISPANIC OR LATINO¹	0.3449		0.0401		0.5379	
Mexican	0.1539		0.0301		0.9414	
Puerto Rican	0.1814		0.5631		0.9509	
Central or South American	0.6153		0.4994		0.5296	
Cuban	0.2883		0.0550		0.7409	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.110C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	1,156	1,048	591	559	289	270
NOT HISPANIC OR LATINO¹	1,107	1,003	561	518	263	252
White	1,029	918	488	465	241	236
Black or African American	311	306	201	189	91	81
American Indian or Alaska Native	59	45	33	40	20	29
Native Hawaiian or Other Pacific Islander	40	35	22	29	9	8
Chinese	93	118	34	41	8	21
Filipino	108	90	39	46	14	15
Japanese	63	*	18	28	4	4
Asian Indian	89	67	43	38	15	12
Korean	54	41	31	27	*	11
Vietnamese	*	45	18	21	11	12
HISPANIC OR LATINO¹	357	308	237	205	99	91
Mexican	266	238	189	165	75	74
Puerto Rican	96	101	71	71	37	33
Central or South American	121	116	81	65	24	30
Cuban	78	71	50	43	*	15

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.110D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.34	0.33	0.25	0.24	0.14	0.13
NOT HISPANIC OR LATINO¹	0.36	0.35	0.27	0.25	0.14	0.14
White	0.40	0.38	0.29	0.28	0.16	0.16
Black or African American	0.87	0.97	0.73	0.73	0.40	0.34
American Indian or Alaska Native	4.22	4.04	2.69	3.58	1.72	2.58
Native Hawaiian or Other Pacific Islander	5.52	5.23	3.91	4.68	1.94	1.47
Chinese	3.78	4.21	1.98	1.88	0.46	0.94
Filipino	4.08	4.01	2.12	2.72	0.78	0.98
Japanese	6.03	*	2.86	3.29	0.58	0.58
Asian Indian	3.46	2.73	2.14	1.73	0.84	0.54
Korean	5.99	5.37	4.16	4.19	*	1.92
Vietnamese	*	5.53	2.97	3.16	1.78	1.74
HISPANIC OR LATINO¹	0.99	0.86	0.78	0.70	0.40	0.35
Mexican	1.16	1.05	0.99	0.88	0.48	0.46
Puerto Rican	2.99	2.90	2.43	2.36	1.45	1.25
Central or South American	2.53	2.43	1.92	1.65	0.69	0.82
Cuban	3.82	3.46	3.75	3.23	*	1.28

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.110P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.1364		0.8786		0.9877	
NOT HISPANIC OR LATINO¹	0.0432		0.7880		0.8487	
White	0.0103		0.6709		0.5594	
Black or African American	1.0000		0.4262		0.7268	
American Indian or Alaska Native	0.5321		0.7494		0.9759	
Native Hawaiian or Other Pacific Islander	0.6065		0.6769		0.0516	
Chinese	0.9932		0.3385		0.0750	
Filipino	0.6306		0.3843		0.9792	
Japanese	*		0.9780		0.3729	
Asian Indian	0.0215		0.8325		0.5787	
Korean	0.7973		0.3354		*	
Vietnamese	*		0.2733		0.9750	
HISPANIC OR LATINO¹	0.3818		0.2247		0.6568	
Mexican	0.1786		0.2390		0.8513	
Puerto Rican	0.7361		0.9524		0.5129	
Central or South American	0.3547		0.2536		0.5701	
Cuban	0.1689		0.9062		*	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.111C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	272	286	246	268	191	208
GEOGRAPHIC DIVISION						
Northeast	131	129	116	122	97	108
New England	55	85	57	83	56	78
Middle Atlantic	119	96	101	89	79	74
Midwest	126	120	118	115	99	91
East North Central	110	101	101	98	81	78
West North Central	62	69	62	65	56	51
South	146	188	133	176	91	127
South Atlantic	98	129	88	122	62	87
East South Central	48	60	48	54	35	36
West South Central	96	113	88	106	57	83
West	140	137	123	128	97	93
Mountain	67	63	58	57	39	45
Pacific	123	122	108	114	89	81
COUNTY TYPE						
Large Metro	224	218	200	202	149	151
Small Metro	178	171	162	160	127	126
250K – 1 Mil. Pop.	140	140	124	130	97	106
< 250K Pop.	124	105	115	97	89	72
Nonmetro	114	123	101	110	71	77
Urbanized	89	79	78	72	58	51
Less Urbanized	85	91	76	81	52	55
Completely Rural	34	38	29	34	20	23

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.111D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.41	0.41	0.41	0.43	0.39	0.41
GEOGRAPHIC DIVISION						
Northeast	1.04	0.99	1.04	1.04	1.00	1.15
New England	1.45	2.46	1.77	2.50	2.27	2.91
Middle Atlantic	1.32	1.03	1.25	1.08	1.08	1.14
Midwest	0.83	0.80	0.84	0.85	0.85	0.83
East North Central	0.99	0.93	0.98	1.03	0.95	1.00
West North Central	1.54	1.56	1.66	1.52	1.79	1.50
South	0.62	0.74	0.62	0.75	0.57	0.68
South Atlantic	0.90	1.06	0.85	1.06	0.79	0.95
East South Central	1.57	1.58	1.69	1.59	1.42	1.23
West South Central	1.02	1.27	1.08	1.34	1.00	1.31
West	0.88	0.95	0.86	0.95	0.83	0.86
Mountain	1.34	1.57	1.37	1.46	1.18	1.40
Pacific	1.11	1.18	1.07	1.20	1.06	1.06
COUNTY TYPE						
Large Metro	0.63	0.58	0.62	0.60	0.57	0.57
Small Metro	0.72	0.76	0.76	0.80	0.76	0.79
250K – 1 Mil. Pop.	0.86	0.97	0.91	0.98	0.93	1.00
< 250K Pop.	1.24	1.29	1.32	1.39	1.33	1.32
Nonmetro	1.01	0.93	1.03	0.91	0.85	0.81
Urbanized	1.47	1.63	1.47	1.66	1.46	1.42
Less Urbanized	1.53	1.20	1.48	1.18	1.19	1.05
Completely Rural	3.49	2.64	3.31	2.51	2.60	2.35

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.111P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.1074		0.6221		0.3878	
GEOGRAPHIC DIVISION						
Northeast	0.1721		0.2133		0.5477	
New England	0.0994		0.3841		0.7905	
Middle Atlantic	0.5449		0.3511		0.5656	
Midwest	0.1675		0.3743		0.7381	
East North Central	0.3627		0.6441		0.7590	
West North Central	0.2703		0.3765		0.9110	
South	0.5043		0.2087		0.7832	
South Atlantic	0.8968		0.4104		0.8705	
East South Central	0.4526		0.9179		0.3083	
West South Central	0.1227		0.2133		0.2274	
West	0.1441		0.6113		0.2658	
Mountain	0.9615		0.4847		0.8517	
Pacific	0.1036		0.3503		0.1773	
COUNTY TYPE						
Large Metro	0.7988		0.6532		0.4566	
Small Metro	0.0782		0.3334		0.0667	
250K – 1 Mil. Pop.	0.2575		0.8623		0.4678	
< 250K Pop.	0.1663		0.1823		0.0362	
Nonmetro	0.3478		0.5366		0.4701	
Urbanized	0.5258		0.8948		0.9530	
Less Urbanized	0.7165		0.8634		0.7935	
Completely Rural	0.5990		0.3202		0.1135	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.112C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	1,988	1,904	1,688	1,662	1,513	1,452
GEOGRAPHIC DIVISION						
Northeast	896	870	771	771	683	680
New England	425	395	364	367	346	327
Middle Atlantic	788	768	680	674	589	594
Midwest	722	761	616	634	529	559
East North Central	555	629	498	535	433	468
West North Central	462	421	364	344	303	305
South	1,253	1,169	1,040	990	929	882
South Atlantic	1,040	947	843	828	759	743
East South Central	441	433	320	332	246	305
West South Central	541	560	519	468	476	408
West	1,030	1,009	890	893	824	784
Mountain	419	456	385	416	365	381
Pacific	941	883	802	775	739	679
COUNTY TYPE						
Large Metro	1,659	1,542	1,420	1,373	1,237	1,216
Small Metro	1,256	1,092	1,016	921	888	792
250K – 1 Mil. Pop.	1,009	886	845	769	734	660
< 250K Pop.	767	675	587	524	503	449
Nonmetro	767	861	605	646	526	543
Urbanized	510	555	399	432	346	381
Less Urbanized	590	637	462	465	382	370
Completely Rural	317	406	253	308	231	264

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.112D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2004 and 2005

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.33	0.30	0.43	0.44	0.46	0.47
GEOGRAPHIC DIVISION						
Northeast	0.70	0.65	0.97	0.96	1.07	1.10
New England	1.37	1.04	1.40	1.71	1.60	1.90
Middle Atlantic	0.81	0.80	1.22	1.16	1.32	1.33
Midwest	0.44	0.43	0.75	0.73	0.81	0.82
East North Central	0.48	0.50	0.91	0.87	0.93	0.91
West North Central	0.91	0.87	1.30	1.32	1.61	1.64
South	0.62	0.56	0.79	0.78	0.84	0.80
South Atlantic	0.95	0.82	1.15	1.14	1.25	1.19
East South Central	1.29	1.24	1.49	1.76	1.55	1.79
West South Central	0.96	0.92	1.42	1.24	1.43	1.26
West	0.83	0.73	0.92	0.93	1.00	1.02
Mountain	1.16	0.90	1.46	1.32	1.59	1.52
Pacific	1.07	0.95	1.15	1.19	1.26	1.32
COUNTY TYPE						
Large Metro	0.46	0.43	0.59	0.61	0.62	0.68
Small Metro	0.56	0.54	0.80	0.79	0.86	0.81
250K – 1 Mil. Pop.	0.69	0.68	0.91	0.93	0.98	0.92
< 250K Pop.	0.93	0.93	1.42	1.38	1.53	1.51
Nonmetro	0.86	0.71	1.05	0.95	1.07	0.97
Urbanized	0.99	1.10	1.53	1.60	1.58	1.68
Less Urbanized	1.39	0.99	1.51	1.27	1.45	1.32
Completely Rural	2.21	2.57	3.20	3.98	3.79	3.78

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.112P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOTAL	0.0785		0.0086		0.0034	
GEOGRAPHIC DIVISION						
Northeast	0.3662		0.5808		0.8071	
New England	0.6464		0.8186		0.2580	
Middle Atlantic	0.2018		0.4708		0.7879	
Midwest	0.1086		0.2588		0.1542	
East North Central	0.5720		0.7092		0.2720	
West North Central	0.0612		0.1266		0.3485	
South	0.2359		0.0574		0.0063	
South Atlantic	0.2581		0.0590		0.0048	
East South Central	0.5505		0.8966		0.2277	
West South Central	0.8812		0.4518		0.8577	
West	0.7920		0.1085		0.1147	
Mountain	0.3577		0.0601		0.0280	
Pacific	0.9111		0.3894		0.5134	
COUNTY TYPE						
Large Metro	0.2446		0.0114		0.0299	
Small Metro	0.4273		0.0968		0.0466	
250K – 1 Mil. Pop.	0.1444		0.0488		0.0042	
< 250K Pop.	0.4766		0.8348		0.7442	
Nonmetro	0.2557		0.6475		0.5752	
Urbanized	0.6173		0.5223		0.4843	
Less Urbanized	0.3038		0.7303		0.3107	
Completely Rural	0.5839		0.1724		0.2133	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 2.113C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2004 and 2005**

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2004	2005	2004	2005	2004	2005
TOTAL	191	208	160	170	98	86
GEOGRAPHIC DIVISION						
Northeast	97	108	80	86	56	49
New England	56	78	49	63	40	37
Middle Atlantic	79	74	63	58	39	32
Midwest	99	91	85	79	44	38
East North Central	81	78	65	70	31	33
West North Central	56	51	54	39	32	20
South	91	127	77	99	50	46
South Atlantic	62	87	47	65	33	31
East South Central	35	36	31	27	18	17
West South Central	57	83	52	66	34	28
West	97	93	77	73	43	37
Mountain	39	45	33	37	16	20
Pacific	89	81	70	64	40	31
COUNTY TYPE						
Large Metro	149	151	114	122	61	60
Small Metro	127	126	107	102	70	54
250K – 1 Mil. Pop.	97	106	82	86	52	49
< 250K Pop.	89	72	71	56	46	23
Nonmetro	71	77	59	56	30	28
Urbanized	58	51	44	39	23	21
Less Urbanized	52	55	44	38	18	17
Completely Rural	20	23	16	19	9	10

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 2.113D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics:
Standard Errors of Percentages, 2004 and 2005**

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2004	2005	2004	2005	2004	2005
TOTAL	0.39	0.41	0.36	0.37	0.24	0.21
GEOGRAPHIC DIVISION						
Northeast	1.00	1.15	0.93	1.01	0.77	0.64
New England	2.27	2.91	2.15	2.65	2.10	1.78
Middle Atlantic	1.08	1.14	0.99	0.98	0.72	0.59
Midwest	0.85	0.83	0.81	0.78	0.48	0.41
East North Central	0.95	1.00	0.83	0.98	0.45	0.49
West North Central	1.79	1.50	1.93	1.26	1.21	0.80
South	0.57	0.68	0.52	0.58	0.36	0.31
South Atlantic	0.79	0.95	0.65	0.77	0.48	0.43
East South Central	1.42	1.23	1.31	1.02	0.78	0.72
West South Central	1.00	1.31	1.03	1.18	0.71	0.55
West	0.83	0.86	0.77	0.72	0.46	0.39
Mountain	1.18	1.40	1.10	1.22	0.60	0.69
Pacific	1.06	1.06	0.98	0.88	0.59	0.47
COUNTY TYPE						
Large Metro	0.57	0.57	0.51	0.50	0.30	0.27
Small Metro	0.76	0.79	0.70	0.72	0.53	0.43
250K – 1 Mil. Pop.	0.93	1.00	0.84	0.90	0.62	0.58
< 250K Pop.	1.33	1.32	1.29	1.18	0.97	0.57
Nonmetro	0.85	0.81	0.75	0.66	0.44	0.41
Urbanized	1.46	1.42	1.18	1.19	0.73	0.78
Less Urbanized	1.19	1.05	1.11	0.88	0.55	0.49
Completely Rural	2.60	2.35	2.38	1.88	1.64	1.21

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 2.113P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2005 Versus 2004**

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2004	2005	2004	2005	2004	2005
TOTAL	0.3878		0.1124		0.3447	
GEOGRAPHIC DIVISION						
Northeast	0.5477		0.4270		0.3055	
New England	0.7905		0.4219		0.6224	
Middle Atlantic	0.5656		0.7095		0.3272	
Midwest	0.7381		0.3914		0.6160	
East North Central	0.7590		0.6513		0.6464	
West North Central	0.9110		0.4305		0.8397	
South	0.7832		0.8268		0.6447	
South Atlantic	0.8705		0.3551		0.7240	
East South Central	0.3083		0.5372		0.7706	
West South Central	0.2274		0.3664		0.5960	
West	0.2658		0.1740		0.3826	
Mountain	0.8517		0.9200		0.2209	
Pacific	0.1773		0.1353		0.1123	
COUNTY TYPE						
Large Metro	0.4566		0.7932		0.7274	
Small Metro	0.0667		0.1015		0.3792	
250K – 1 Mil. Pop.	0.4678		0.4443		0.7698	
< 250K Pop.	0.0362		0.0900		0.0512	
Nonmetro	0.4701		0.0270		0.1010	
Urbanized	0.9530		0.8740		0.2499	
Less Urbanized	0.7935		0.0170		0.2584	
Completely Rural	0.1135		0.2057		0.8068	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.114C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2004	2005	2004	2005	2004	2005
TOTAL	1,513	1,452	787	795	384	379
GEOGRAPHIC DIVISION						
Northeast	683	680	362	347	166	173
New England	346	327	161	176	75	97
Middle Atlantic	589	594	325	299	148	144
Midwest	529	559	327	324	201	185
East North Central	433	468	284	247	179	135
West North Central	303	305	162	199	92	120
South	929	882	486	481	222	214
South Atlantic	759	743	378	371	155	152
East South Central	246	305	146	189	75	80
West South Central	476	408	267	250	141	136
West	824	784	382	374	174	186
Mountain	365	381	159	183	71	72
Pacific	739	679	348	325	158	169
COUNTY TYPE						
Large Metro	1,237	1,216	624	621	278	287
Small Metro	888	792	467	446	242	215
250K – 1 Mil. Pop.	734	660	364	344	194	177
< 250K Pop.	503	449	291	280	142	121
Nonmetro	526	543	320	294	146	143
Urbanized	346	381	210	202	88	81
Less Urbanized	382	370	224	207	112	113
Completely Rural	231	264	101	110	53	51

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.114D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2004 and 2005

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2004	2005	2004	2005	2004	2005
TOTAL	0.46	0.47	0.34	0.33	0.18	0.18
GEOGRAPHIC DIVISION						
Northeast	1.07	1.10	0.84	0.76	0.42	0.42
New England	1.60	1.90	1.53	1.40	0.75	0.88
Middle Atlantic	1.32	1.33	1.00	0.90	0.50	0.48
Midwest	0.81	0.82	0.65	0.63	0.42	0.39
East North Central	0.93	0.91	0.81	0.69	0.53	0.41
West North Central	1.61	1.64	1.07	1.32	0.62	0.83
South	0.84	0.80	0.58	0.56	0.30	0.28
South Atlantic	1.25	1.19	0.84	0.79	0.38	0.36
East South Central	1.55	1.79	1.04	1.35	0.60	0.63
West South Central	1.43	1.26	1.08	0.97	0.63	0.61
West	1.00	1.02	0.71	0.76	0.37	0.41
Mountain	1.59	1.52	0.98	1.22	0.48	0.53
Pacific	1.26	1.32	0.91	0.94	0.48	0.53
COUNTY TYPE						
Large Metro	0.62	0.68	0.47	0.49	0.24	0.25
Small Metro	0.86	0.81	0.65	0.60	0.39	0.33
250K – 1 Mil. Pop.	0.98	0.92	0.76	0.72	0.47	0.42
< 250K Pop.	1.53	1.51	1.15	1.06	0.65	0.55
Nonmetro	1.07	0.97	0.77	0.68	0.38	0.39
Urbanized	1.58	1.68	1.21	1.20	0.57	0.57
Less Urbanized	1.45	1.32	1.05	0.91	0.57	0.61
Completely Rural	3.79	3.78	2.17	1.82	1.69	1.04

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.114P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2004	2005	2004	2005	2004	2005
TOTAL	0.0034		0.9425		0.1654	
GEOGRAPHIC DIVISION						
Northeast	0.8071		0.1978		0.6509	
New England	0.2580		0.1621		0.8156	
Middle Atlantic	0.7879		0.4872		0.6969	
Midwest	0.1542		0.7780		0.2368	
East North Central	0.2720		0.5530		0.0683	
West North Central	0.3485		0.7232		0.5548	
South	0.0063		0.7080		0.0515	
South Atlantic	0.0048		0.8142		0.0119	
East South Central	0.2277		0.0782		0.6588	
West South Central	0.8577		0.8865		0.9090	
West	0.1147		0.2487		0.7816	
Mountain	0.0280		0.7298		0.8402	
Pacific	0.5134		0.2670		0.7007	
COUNTY TYPE						
Large Metro	0.0299		0.3127		0.2812	
Small Metro	0.0466		0.1887		0.3926	
250K – 1 Mil. Pop.	0.0042		0.3367		0.7686	
< 250K Pop.	0.7442		0.3487		0.2754	
Nonmetro	0.5752		0.7356		0.8381	
Urbanized	0.4843		0.9345		0.7689	
Less Urbanized	0.3107		0.2105		0.4733	
Completely Rural	0.2133		0.1876		0.1266	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.