

Table 2.1C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	812	793	818	885	794	845
Cigarettes	824	858	788	817	768	777
Smokeless Tobacco	656	669	267	295	245	267
Cigars	803	852	448	481	323	354
Pipe Tobacco ²	766	705	--	--	147	146
ALCOHOL	705	649	885	917	957	975
Binge Alcohol Use ³	--	--	--	--	698	701
Heavy Alcohol Use ³	--	--	--	--	382	379

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.1D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.34	0.33	0.34	0.36	0.33	0.35
Cigarettes	0.34	0.35	0.33	0.34	0.32	0.32
Smokeless Tobacco	0.27	0.28	0.11	0.12	0.10	0.11
Cigars	0.33	0.35	0.19	0.20	0.13	0.15
Pipe Tobacco ²	0.32	0.29	--	--	0.06	0.06
ALCOHOL	0.29	0.27	0.37	0.38	0.40	0.40
Binge Alcohol Use ³	--	--	--	--	0.29	0.29
Heavy Alcohol Use ³	--	--	--	--	0.16	0.16

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.1P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.1770		0.4291		0.6856	
Cigarettes	0.1330		0.9044		0.9728	
Smokeless Tobacco	0.7226		0.1500		0.2195	
Cigars	0.9104		0.3315		0.6155	
Pipe Tobacco ²	0.1576		--		0.1095	
ALCOHOL	0.2069		0.0133		0.0067	
Binge Alcohol Use ³	--		--		0.8010	
Heavy Alcohol Use ³	--		--		0.1202	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.2C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	106	113	96	99	80	78
Cigarettes	105	108	89	90	76	71
Smokeless Tobacco	58	58	45	46	32	29
Cigars	80	79	67	67	44	46
Pipe Tobacco ²	34	38	--	--	19	17
ALCOHOL	108	111	103	105	81	81
Binge Alcohol Use ³	--	--	--	--	73	65
Heavy Alcohol Use ³	--	--	--	--	36	32

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.2D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.42	0.44	0.38	0.39	0.32	0.31
Cigarettes	0.42	0.42	0.35	0.36	0.30	0.28
Smokeless Tobacco	0.23	0.23	0.18	0.18	0.13	0.12
Cigars	0.32	0.31	0.27	0.27	0.18	0.18
Pipe Tobacco ²	0.13	0.15	--	--	0.08	0.07
ALCOHOL	0.43	0.44	0.41	0.41	0.32	0.32
Binge Alcohol Use ³	--	--	--	--	0.29	0.26
Heavy Alcohol Use ³	--	--	--	--	0.14	0.13

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.2P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.0014		0.1000		0.0031	
Cigarettes	0.0000		0.0259		0.0069	
Smokeless Tobacco	1.0000		0.8252		0.2407	
Cigars	0.0887		0.3034		0.0242	
Pipe Tobacco ²	0.1714		--		0.4228	
ALCOHOL	0.0250		0.3763		0.0190	
Binge Alcohol Use ³	--		--		0.0027	
Heavy Alcohol Use ³	--		--		0.1856	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.3C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	143	148	162	155	161	155
Cigarettes	153	155	166	156	159	152
Smokeless Tobacco	119	120	74	82	60	64
Cigars	152	152	122	123	97	92
Pipe Tobacco ²	83	82	--	--	30	37
ALCOHOL	117	114	142	139	164	166
Binge Alcohol Use ³	--	--	--	--	166	171
Heavy Alcohol Use ³	--	--	--	--	120	120

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.3D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.44	0.46	0.50	0.48	0.50	0.48
Cigarettes	0.48	0.48	0.52	0.48	0.49	0.47
Smokeless Tobacco	0.37	0.37	0.23	0.25	0.19	0.20
Cigars	0.47	0.47	0.38	0.38	0.30	0.28
Pipe Tobacco ²	0.26	0.25	--	--	0.09	0.11
ALCOHOL	0.36	0.35	0.44	0.43	0.51	0.51
Binge Alcohol Use ³	--	--	--	--	0.52	0.53
Heavy Alcohol Use ³	--	--	--	--	0.37	0.37

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.3P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.0695		0.9894		0.6334	
Cigarettes	0.0450		0.7243		0.4147	
Smokeless Tobacco	0.3100		0.1060		0.4928	
Cigars	0.1569		0.7218		0.0685	
Pipe Tobacco ²	0.9455		--		0.1339	
ALCOHOL	0.3085		0.7943		0.5735	
Binge Alcohol Use ³	--		--		0.3902	
Heavy Alcohol Use ³	--		--		0.6611	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.4C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	768	747	768	829	752	798
Cigarettes	783	807	733	758	718	726
Smokeless Tobacco	641	645	248	277	236	256
Cigars	769	822	420	450	311	335
Pipe Tobacco ²	735	687	--	--	142	141
ALCOHOL	670	617	863	889	933	936
Binge Alcohol Use ³	--	--	--	--	654	650
Heavy Alcohol Use ³	--	--	--	--	356	347

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.4D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.42	0.40	0.42	0.45	0.41	0.43
Cigarettes	0.43	0.44	0.40	0.41	0.39	0.39
Smokeless Tobacco	0.35	0.35	0.14	0.15	0.13	0.14
Cigars	0.42	0.44	0.23	0.24	0.17	0.18
Pipe Tobacco ²	0.40	0.37	--	--	0.08	0.08
ALCOHOL	0.37	0.33	0.47	0.48	0.51	0.51
Binge Alcohol Use ³	--	--	--	--	0.36	0.35
Heavy Alcohol Use ³	--	--	--	--	0.19	0.19

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.4P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.4971		0.2959		0.4141	
Cigarettes	0.4796		0.6353		0.6842	
Smokeless Tobacco	0.8452		0.3006		0.2108	
Cigars	0.7663		0.2647		0.7390	
Pipe Tobacco ²	0.1519		--		0.1582	
ALCOHOL	0.0692		0.0079		0.0041	
Binge Alcohol Use ³	--		--		0.8626	
Heavy Alcohol Use ³	--		--		0.0895	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.5C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	44	44	32	35	24	20
Cigarettes	40	40	30	33	21	18
Smokeless Tobacco	19	17	14	14	9	9
Cigars	25	22	19	16	13	9
Pipe Tobacco ²	11	11	--	--	6	6
ALCOHOL	55	49	42	41	26	26
Binge Alcohol Use ³	--	--	--	--	21	18
Heavy Alcohol Use ³	--	--	--	--	7	7

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.5D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.50	0.49	0.37	0.41	0.28	0.24
Cigarettes	0.46	0.46	0.35	0.38	0.25	0.22
Smokeless Tobacco	0.22	0.21	0.17	0.17	0.11	0.10
Cigars	0.29	0.27	0.22	0.19	0.15	0.11
Pipe Tobacco ²	0.14	0.14	--	--	0.07	0.08
ALCOHOL	0.58	0.53	0.47	0.46	0.31	0.31
Binge Alcohol Use ³	--	--	--	--	0.25	0.22
Heavy Alcohol Use ³	--	--	--	--	0.08	0.08

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.5P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.0470		0.9512		0.3538	
Cigarettes	0.0872		1.0000		0.2163	
Smokeless Tobacco	0.3044		0.3346		0.5290	
Cigars	0.1266		0.1541		0.1859	
Pipe Tobacco ²	0.5794		--		0.9862	
ALCOHOL	0.1491		0.6228		0.7236	
Binge Alcohol Use ³	--		--		0.8193	
Heavy Alcohol Use ³	--		--		0.9587	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.6C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	76	75	61	57	43	40
Cigarettes	71	67	54	50	41	37
Smokeless Tobacco	33	34	27	26	17	15
Cigars	46	47	36	39	23	23
Pipe Tobacco ²	23	21	--	--	12	11
ALCOHOL	89	87	81	77	49	50
Binge Alcohol Use ³	--	--	--	--	38	36
Heavy Alcohol Use ³	--	--	--	--	16	17

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.6D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.68	0.74	0.60	0.61	0.47	0.45
Cigarettes	0.67	0.68	0.55	0.55	0.45	0.42
Smokeless Tobacco	0.37	0.38	0.31	0.30	0.20	0.17
Cigars	0.49	0.51	0.39	0.43	0.26	0.26
Pipe Tobacco ²	0.25	0.24	--	--	0.14	0.13
ALCOHOL	0.73	0.76	0.68	0.70	0.49	0.53
Binge Alcohol Use ³	--	--	--	--	0.40	0.40
Heavy Alcohol Use ³	--	--	--	--	0.18	0.20

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.6P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.0050		0.0339		0.0030	
Cigarettes	0.0001		0.0048		0.0059	
Smokeless Tobacco	0.6539		0.3361		0.5125	
Cigars	0.2080		0.8374		0.1562	
Pipe Tobacco ²	0.2439		--		0.5277	
ALCOHOL	0.0846		0.3319		0.0868	
Binge Alcohol Use ³	--		--		0.0477	
Heavy Alcohol Use ³	--		--		0.7397	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.7C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	96	95	82	80	69	63
Cigarettes	92	91	75	73	63	57
Smokeless Tobacco	44	43	34	33	26	24
Cigars	69	65	57	55	37	38
Pipe Tobacco ²	26	29	--	--	14	11
ALCOHOL	112	109	102	101	76	71
Binge Alcohol Use ³	--	--	--	--	62	54
Heavy Alcohol Use ³	--	--	--	--	32	27

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.7D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.79	0.80	0.80	0.79	0.71	0.66
Cigarettes	0.79	0.82	0.76	0.75	0.67	0.61
Smokeless Tobacco	0.50	0.48	0.39	0.38	0.31	0.28
Cigars	0.71	0.68	0.61	0.59	0.42	0.43
Pipe Tobacco ²	0.31	0.33	--	--	0.17	0.13
ALCOHOL	0.72	0.78	0.76	0.80	0.73	0.70
Binge Alcohol Use ³	--	--	--	--	0.66	0.58
Heavy Alcohol Use ³	--	--	--	--	0.39	0.31

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.7P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.0243		0.1649		0.0278	
Cigarettes	0.0041		0.1168		0.0696	
Smokeless Tobacco	0.8282		0.7516		0.3103	
Cigars	0.1926		0.2246		0.0663	
Pipe Tobacco ²	0.3858		--		0.5070	
ALCOHOL	0.0332		0.2576		0.0176	
Binge Alcohol Use ³	--		--		0.0018	
Heavy Alcohol Use ³	--		--		0.0469	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.8C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	182	188	159	164	134	138
Cigarettes	172	175	145	142	119	122
Smokeless Tobacco	77	82	52	57	37	41
Cigars	133	139	106	109	74	74
Pipe Tobacco ²	48	53	--	--	22	27
ALCOHOL	210	222	197	212	162	180
Binge Alcohol Use ³	--	--	--	--	138	151
Heavy Alcohol Use ³	--	--	--	--	84	78

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.8D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.69	0.69	0.75	0.73	0.75	0.71
Cigarettes	0.73	0.74	0.77	0.75	0.72	0.71
Smokeless Tobacco	0.53	0.54	0.37	0.39	0.28	0.30
Cigars	0.72	0.69	0.65	0.61	0.52	0.49
Pipe Tobacco ²	0.36	0.38	--	--	0.17	0.20
ALCOHOL	0.62	0.59	0.71	0.70	0.78	0.84
Binge Alcohol Use ³	--	--	--	--	0.79	0.80
Heavy Alcohol Use ³	--	--	--	--	0.59	0.51

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.8P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.2048		0.4661		0.3274	
Cigarettes	0.1983		0.2461		0.1182	
Smokeless Tobacco	0.7071		0.4613		0.7967	
Cigars	0.2240		0.4046		0.4183	
Pipe Tobacco ²	0.9871		--		0.3117	
ALCOHOL	0.3329		0.5519		0.9858	
Binge Alcohol Use ³	--		--		0.5672	
Heavy Alcohol Use ³	--		--		0.4308	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.9C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	261	252	214	207	189	184
Cigarettes	252	242	197	191	176	170
Smokeless Tobacco	115	116	60	66	49	51
Cigars	193	183	121	118	83	72
Pipe Tobacco ²	72	71	--	--	21	27
ALCOHOL	290	285	270	277	239	250
Binge Alcohol Use ³	--	--	--	--	184	196
Heavy Alcohol Use ³	--	--	--	--	102	109

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.9D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.55	0.57	0.65	0.61	0.62	0.60
Cigarettes	0.59	0.57	0.65	0.60	0.61	0.58
Smokeless Tobacco	0.46	0.48	0.29	0.32	0.24	0.25
Cigars	0.62	0.62	0.48	0.47	0.37	0.33
Pipe Tobacco ²	0.33	0.33	--	--	0.11	0.13
ALCOHOL	0.42	0.40	0.52	0.51	0.60	0.60
Binge Alcohol Use ³	--	--	--	--	0.60	0.63
Heavy Alcohol Use ³	--	--	--	--	0.43	0.47

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.9P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.2146		0.5701		0.8382	
Cigarettes	0.1348		0.6111		0.8190	
Smokeless Tobacco	0.3640		0.1426		0.4977	
Cigars	0.4378		0.2995		0.0593	
Pipe Tobacco ²	0.8959		--		0.3021	
ALCOHOL	0.7899		0.6885		0.2645	
Binge Alcohol Use ³	--		--		0.0662	
Heavy Alcohol Use ³	--		--		0.1928	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.10C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	248	269	297	295	287	285
Cigarettes	268	281	293	290	281	274
Smokeless Tobacco	234	246	119	122	102	110
Cigars	276	279	180	189	121	129
Pipe Tobacco ²	158	178	--	--	56	49
ALCOHOL	182	192	244	253	274	293
Binge Alcohol Use ³	--	--	--	--	264	281
Heavy Alcohol Use ³	--	--	--	--	171	165

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.10D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.71	0.77	0.85	0.85	0.82	0.82
Cigarettes	0.77	0.80	0.84	0.83	0.80	0.79
Smokeless Tobacco	0.67	0.71	0.34	0.35	0.29	0.32
Cigars	0.79	0.80	0.52	0.54	0.35	0.37
Pipe Tobacco ²	0.45	0.51	--	--	0.16	0.14
ALCOHOL	0.52	0.55	0.70	0.73	0.78	0.84
Binge Alcohol Use ³	--	--	--	--	0.75	0.81
Heavy Alcohol Use ³	--	--	--	--	0.49	0.47

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.10P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.8711		0.3229		0.6772	
Cigarettes	0.8623		0.4301		0.5923	
Smokeless Tobacco	0.8468		0.9858		0.6155	
Cigars	0.8375		0.2612		0.5850	
Pipe Tobacco ²	0.9306		--		0.5570	
ALCOHOL	0.5880		0.0165		0.0770	
Binge Alcohol Use ³	--		--		0.5210	
Heavy Alcohol Use ³	--		--		0.6966	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.11C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	723	693	712	746	693	718
Cigarettes	744	742	670	670	650	645
Smokeless Tobacco	594	593	227	245	213	226
Cigars	732	764	376	410	297	305
Pipe Tobacco ²	684	652	--	--	133	129
ALCOHOL	631	580	835	828	895	865
Binge Alcohol Use ³	--	--	--	--	579	562
Heavy Alcohol Use ³	--	--	--	--	305	305

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.11D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.49	0.46	0.48	0.50	0.47	0.48
Cigarettes	0.50	0.49	0.45	0.44	0.44	0.43
Smokeless Tobacco	0.40	0.39	0.15	0.16	0.14	0.15
Cigars	0.49	0.51	0.25	0.27	0.20	0.20
Pipe Tobacco ²	0.46	0.43	--	--	0.09	0.09
ALCOHOL	0.43	0.39	0.56	0.55	0.60	0.57
Binge Alcohol Use ³	--	--	--	--	0.39	0.37
Heavy Alcohol Use ³	--	--	--	--	0.21	0.20

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.11P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.5005		0.4116		0.4309	
Cigarettes	0.4070		0.7923		0.7742	
Smokeless Tobacco	0.8131		0.2285		0.2338	
Cigars	0.7892		0.4398		0.8711	
Pipe Tobacco ²	0.1060		--		0.1973	
ALCOHOL	0.0746		0.0322		0.0118	
Binge Alcohol Use ³	--		--		0.6835	
Heavy Alcohol Use ³	--		--		0.0342	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.12C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	795	768	816	878	793	839
Cigarettes	812	833	784	811	765	772
Smokeless Tobacco	647	667	260	289	243	264
Cigars	801	847	442	478	323	351
Pipe Tobacco ²	758	696	--	--	145	145
ALCOHOL	685	625	880	908	950	969
Binge Alcohol Use ³	--	--	--	--	699	701
Heavy Alcohol Use ³	--	--	--	--	383	378

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.12D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.37	0.35	0.38	0.40	0.37	0.39
Cigarettes	0.38	0.38	0.36	0.37	0.36	0.35
Smokeless Tobacco	0.30	0.31	0.12	0.13	0.11	0.12
Cigars	0.37	0.39	0.21	0.22	0.15	0.16
Pipe Tobacco ²	0.35	0.32	--	--	0.07	0.07
ALCOHOL	0.32	0.29	0.41	0.42	0.44	0.44
Binge Alcohol Use ³	--	--	--	--	0.32	0.32
Heavy Alcohol Use ³	--	--	--	--	0.18	0.17

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.12P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.3206		0.3338		0.5022	
Cigarettes	0.2907		0.7220		0.8377	
Smokeless Tobacco	0.7058		0.1504		0.1681	
Cigars	0.9831		0.2591		0.8356	
Pipe Tobacco ²	0.1667		--		0.0864	
ALCOHOL	0.1082		0.0106		0.0039	
Binge Alcohol Use ³	--		--		0.9785	
Heavy Alcohol Use ³	--		--		0.1477	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.13C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	465	468	610	598	570	576
Cigarettes	500	534	549	539	537	515
Smokeless Tobacco	564	570	259	263	234	241
Cigars	622	614	411	421	290	324
Pipe Tobacco ²	655	624	--	--	139	141
ALCOHOL	420	348	568	575	626	621
Binge Alcohol Use ³	--	--	--	--	513	529
Heavy Alcohol Use ³	--	--	--	--	319	321

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.13D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.40	0.40	0.52	0.51	0.49	0.49
Cigarettes	0.43	0.45	0.47	0.46	0.46	0.44
Smokeless Tobacco	0.48	0.48	0.22	0.22	0.20	0.20
Cigars	0.53	0.52	0.35	0.36	0.25	0.27
Pipe Tobacco ²	0.56	0.53	--	--	0.12	0.12
ALCOHOL	0.36	0.30	0.49	0.49	0.54	0.53
Binge Alcohol Use ³	--	--	--	--	0.44	0.45
Heavy Alcohol Use ³	--	--	--	--	0.27	0.27

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.13P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.0645		0.9450		0.9103	
Cigarettes	0.0703		0.4297		0.6257	
Smokeless Tobacco	0.9135		0.2834		0.2867	
Cigars	0.4621		0.5215		0.7124	
Pipe Tobacco ²	0.1129		--		0.1633	
ALCOHOL	0.3736		0.5301		0.1259	
Binge Alcohol Use ³	--		--		0.4053	
Heavy Alcohol Use ³	--		--		0.4174	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.14C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	607	598	519	565	497	543
Cigarettes	611	617	518	558	491	538
Smokeless Tobacco	273	237	73	106	65	87
Cigars	434	434	173	182	122	112
Pipe Tobacco ²	227	217	--	--	43	42
ALCOHOL	508	515	632	624	664	637
Binge Alcohol Use ³	--	--	--	--	390	402
Heavy Alcohol Use ³	--	--	--	--	174	163

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.14D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.49	0.48	0.42	0.45	0.40	0.43
Cigarettes	0.49	0.49	0.42	0.45	0.40	0.43
Smokeless Tobacco	0.22	0.19	0.06	0.08	0.05	0.07
Cigars	0.35	0.35	0.14	0.15	0.10	0.09
Pipe Tobacco ²	0.18	0.17	--	--	0.03	0.03
ALCOHOL	0.41	0.41	0.51	0.50	0.54	0.51
Binge Alcohol Use ³	--	--	--	--	0.31	0.32
Heavy Alcohol Use ³	--	--	--	--	0.14	0.13

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.14P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.6970		0.1960		0.6229	
Cigarettes	0.5842		0.3339		0.6607	
Smokeless Tobacco	0.2092		0.1913		0.4675	
Cigars	0.4515		0.3311		0.5840	
Pipe Tobacco ²	0.8244		--		0.3667	
ALCOHOL	0.3226		0.0030		0.0110	
Binge Alcohol Use ³	--		--		0.5398	
Heavy Alcohol Use ³	--		--		0.0450	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.15C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 to 17: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	74	79	63	74	55	59
Cigarettes	70	75	58	65	52	53
Smokeless Tobacco	48	49	40	40	30	28
Cigars	59	64	49	55	37	38
Pipe Tobacco ²	29	28	--	--	15	14
ALCOHOL	77	77	72	74	59	57
Binge Alcohol Use ³	--	--	--	--	52	48
Heavy Alcohol Use ³	--	--	--	--	27	27

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.15D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 to 17: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.57	0.61	0.49	0.57	0.43	0.46
Cigarettes	0.55	0.58	0.45	0.51	0.40	0.41
Smokeless Tobacco	0.37	0.38	0.31	0.31	0.23	0.22
Cigars	0.46	0.49	0.38	0.43	0.28	0.29
Pipe Tobacco ²	0.23	0.21	--	--	0.11	0.11
ALCOHOL	0.59	0.60	0.56	0.58	0.45	0.44
Binge Alcohol Use ³	--	--	--	--	0.40	0.37
Heavy Alcohol Use ³	--	--	--	--	0.21	0.21

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.15P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.0315		0.8341		0.0851	
Cigarettes	0.0039		0.5304		0.3191	
Smokeless Tobacco	0.9264		0.8340		0.2401	
Cigars	0.2932		0.8751		0.0394	
Pipe Tobacco ²	0.0520		--		0.9199	
ALCOHOL	0.1550		0.4139		0.0361	
Binge Alcohol Use ³	--		--		0.0212	
Heavy Alcohol Use ³	--		--		0.4808	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.16C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 to 17: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	75	75	66	63	53	50
Cigarettes	74	73	63	61	52	48
Smokeless Tobacco	25	26	18	18	11	8
Cigars	52	45	42	36	25	24
Pipe Tobacco ²	20	24	--	--	13	11
ALCOHOL	74	78	72	76	59	58
Binge Alcohol Use ³	--	--	--	--	50	43
Heavy Alcohol Use ³	--	--	--	--	23	20

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.16D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 to 17: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.61	0.60	0.53	0.51	0.43	0.40
Cigarettes	0.60	0.59	0.51	0.49	0.42	0.39
Smokeless Tobacco	0.20	0.21	0.15	0.14	0.09	0.07
Cigars	0.42	0.36	0.34	0.29	0.20	0.19
Pipe Tobacco ²	0.16	0.19	--	--	0.10	0.09
ALCOHOL	0.60	0.63	0.58	0.61	0.48	0.47
Binge Alcohol Use ³	--	--	--	--	0.41	0.35
Heavy Alcohol Use ³	--	--	--	--	0.19	0.16

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.16P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.0126		0.0230		0.0092	
Cigarettes	0.0020		0.0094		0.0033	
Smokeless Tobacco	0.8621		0.9329		0.8330	
Cigars	0.1458		0.1232		0.3111	
Pipe Tobacco ²	0.7811		--		0.2684	
ALCOHOL	0.0728		0.6656		0.2402	
Binge Alcohol Use ³	--		--		0.0461	
Heavy Alcohol Use ³	--		--		0.2305	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.17C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 18 to 25: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	92	95	108	105	106	110
Cigarettes	101	103	112	108	105	107
Smokeless Tobacco	98	101	70	75	58	60
Cigars	117	111	104	99	83	79
Pipe Tobacco ²	75	71	--	--	27	33
ALCOHOL	76	75	93	88	110	101
Binge Alcohol Use ³	--	--	--	--	113	111
Heavy Alcohol Use ³	--	--	--	--	88	96

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.17D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 18 to 25: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.57	0.58	0.67	0.64	0.66	0.67
Cigarettes	0.62	0.63	0.69	0.66	0.65	0.66
Smokeless Tobacco	0.61	0.62	0.43	0.46	0.36	0.37
Cigars	0.72	0.68	0.64	0.60	0.51	0.48
Pipe Tobacco ²	0.46	0.44	--	--	0.17	0.20
ALCOHOL	0.47	0.46	0.58	0.54	0.68	0.62
Binge Alcohol Use ³	--	--	--	--	0.70	0.68
Heavy Alcohol Use ³	--	--	--	--	0.54	0.59

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.17P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 18 to 25: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.0388		0.8065		0.8589	
Cigarettes	0.0298		0.6008		0.4885	
Smokeless Tobacco	0.1976		0.2174		0.6311	
Cigars	0.3843		0.8166		0.0637	
Pipe Tobacco ²	0.6812		--		0.4444	
ALCOHOL	0.2212		0.7584		0.1370	
Binge Alcohol Use ³	--		--		0.1040	
Heavy Alcohol Use ³	--		--		0.5586	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.18C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 18 to 25: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	100	101	107	106	102	101
Cigarettes	104	104	108	105	101	101
Smokeless Tobacco	50	53	18	23	10	12
Cigars	89	92	63	65	44	41
Pipe Tobacco ²	34	36	--	--	12	16
ALCOHOL	78	80	94	98	104	112
Binge Alcohol Use ³	--	--	--	--	96	104
Heavy Alcohol Use ³	--	--	--	--	60	56

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.18D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 18 to 25: Standard Errors of Percentages, 2004 and 2005

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.62	0.63	0.67	0.66	0.64	0.63
Cigarettes	0.65	0.64	0.67	0.65	0.63	0.63
Smokeless Tobacco	0.31	0.33	0.11	0.14	0.06	0.08
Cigars	0.56	0.57	0.39	0.40	0.27	0.26
Pipe Tobacco ²	0.21	0.22	--	--	0.08	0.10
ALCOHOL	0.49	0.50	0.59	0.61	0.65	0.69
Binge Alcohol Use ³	--	--	--	--	0.60	0.65
Heavy Alcohol Use ³	--	--	--	--	0.37	0.35

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 2.18P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 18 to 25: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS¹	0.4726		0.8355		0.5705	
Cigarettes	0.3940		1.0000		0.5941	
Smokeless Tobacco	0.9378		0.0872		0.2438	
Cigars	0.1910		0.7827		0.5755	
Pipe Tobacco ²	0.3242		--		0.0602	
ALCOHOL	0.7445		0.4993		0.5435	
Binge Alcohol Use ³	--		--		0.6909	
Heavy Alcohol Use ³	--		--		0.9622	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.