

Table 2.57C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	595	595	596	568	563	546
NOT HISPANIC OR LATINO¹	548	547	564	538	534	518
White	447	451	509	479	479	459
Black or African American	250	210	189	188	181	180
American Indian or Alaska Native	94	86	50	47	41	42
Native Hawaiian or Other Pacific Islander	59	49	44	33	32	28
Chinese	94	132	34	41	30	33
Filipino	98	117	49	46	43	42
Japanese	85	83	26	16	24	17
Asian Indian	93	71	63	35	58	28
Korean	72	40	40	26	39	24
Vietnamese	47	48	34	36	33	35
HISPANIC OR LATINO¹	206	229	193	196	185	181
Mexican	311	298	188	184	171	162
Puerto Rican	135	131	83	89	81	82
Central or South American	153	136	90	76	75	70
Cuban	106	75	77	51	61	45

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.57D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.25	0.25	0.25	0.23	0.24	0.23
NOT HISPANIC OR LATINO¹	0.26	0.26	0.27	0.26	0.26	0.25
White	0.27	0.27	0.31	0.29	0.29	0.27
Black or African American	0.92	0.74	0.69	0.66	0.67	0.64
American Indian or Alaska Native	2.38	2.76	2.56	3.23	2.54	2.97
Native Hawaiian or Other Pacific Islander	4.44	4.75	5.31	4.32	4.09	3.77
Chinese	3.75	4.02	1.71	1.69	1.46	1.36
Filipino	3.58	3.92	2.67	2.28	2.24	2.10
Japanese	4.43	5.77	2.89	1.76	2.67	1.88
Asian Indian	3.44	2.52	2.60	1.40	2.48	1.14
Korean	5.23	3.95	4.17	2.97	4.05	2.74
Vietnamese	4.47	4.07	3.51	3.76	3.41	3.70
HISPANIC OR LATINO¹	0.70	0.73	0.65	0.62	0.63	0.57
Mexican	0.85	0.87	0.72	0.73	0.71	0.67
Puerto Rican	1.80	2.23	2.12	2.20	2.10	2.16
Central or South American	2.26	1.88	1.86	1.54	1.69	1.45
Cuban	3.52	3.21	3.75	3.43	3.05	3.10

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.57P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.0000		0.0234		0.0108	
NOT HISPANIC OR LATINO¹	0.0000		0.0206		0.0153	
White	0.0002		0.2151		0.1542	
Black or African American	0.0265		0.0816		0.0884	
American Indian or Alaska Native	0.4458		0.3506		0.4137	
Native Hawaiian or Other Pacific Islander	0.6227		0.7276		0.7935	
Chinese	0.5055		0.4291		0.6088	
Filipino	0.5574		0.6877		0.8031	
Japanese	0.6649		0.0411		0.0577	
Asian Indian	0.0720		0.0126		0.0243	
Korean	0.0358		0.2087		0.1543	
Vietnamese	0.8872		0.6314		0.5157	
HISPANIC OR LATINO¹	0.0048		0.8903		0.5734	
Mexican	0.0233		0.9745		0.4636	
Puerto Rican	0.4164		0.7423		0.8512	
Central or South American	0.0109		0.0487		0.2607	
Cuban	0.6568		0.8255		0.8056	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.58C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	80	75	64	63	54	52
NOT HISPANIC OR LATINO¹	73	66	59	57	50	47
White	86	72	61	59	48	46
Black or African American	34	33	22	22	16	17
American Indian or Alaska Native	11	8	9	6	7	5
Native Hawaiian or Other Pacific Islander	*	*	*	3	2	2
Chinese	5	7	4	5	3	3
Filipino	6	*	5	3	3	2
Japanese	*	*	*	*	*	*
Asian Indian	7	8	5	5	3	4
Korean	5	5	*	4	1	4
Vietnamese	6	*	2	*	1	*
HISPANIC OR LATINO¹	33	33	26	27	19	22
Mexican	41	36	27	29	18	21
Puerto Rican	16	13	11	10	9	7
Central or South American	14	11	10	7	6	6
Cuban	8	6	6	6	5	4

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.58D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.32	0.30	0.26	0.25	0.22	0.21
NOT HISPANIC OR LATINO¹	0.35	0.31	0.28	0.27	0.24	0.22
White	0.41	0.36	0.34	0.32	0.29	0.27
Black or African American	0.72	0.70	0.52	0.51	0.40	0.41
American Indian or Alaska Native	3.43	3.86	3.19	3.05	2.82	2.60
Native Hawaiian or Other Pacific Islander	*	*	*	3.57	2.85	2.75
Chinese	2.33	3.25	1.94	2.66	1.30	1.62
Filipino	3.74	*	2.93	1.84	2.04	1.29
Japanese	*	*	*	*	*	*
Asian Indian	2.99	2.73	2.58	2.03	1.66	1.43
Korean	4.72	4.04	*	3.24	0.95	3.23
Vietnamese	4.28	*	2.01	*	1.00	*
HISPANIC OR LATINO¹	0.83	0.77	0.64	0.63	0.48	0.51
Mexican	1.04	0.96	0.78	0.81	0.58	0.64
Puerto Rican	1.98	2.18	1.77	1.81	1.59	1.33
Central or South American	2.41	2.23	1.93	1.59	1.33	1.33
Cuban	4.76	3.72	4.19	3.52	3.95	2.54

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.58P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.0000		0.0000		0.0001	
NOT HISPANIC OR LATINO¹	0.0000		0.0000		0.0000	
White	0.0000		0.0001		0.0001	
Black or African American	0.0048		0.0624		0.4267	
American Indian or Alaska Native	0.0845		0.0958		0.0399	
Native Hawaiian or Other Pacific Islander	*		*		0.6605	
Chinese	0.6076		0.7681		0.6442	
Filipino	*		0.1509		0.3185	
Japanese	*		*		*	
Asian Indian	0.1742		0.1157		0.2884	
Korean	0.5802		*		0.0249	
Vietnamese	*		*		*	
HISPANIC OR LATINO¹	0.0002		0.2436		0.6928	
Mexican	0.0089		0.4408		0.9771	
Puerto Rican	0.0684		0.4343		0.2394	
Central or South American	0.0323		0.1225		0.7638	
Cuban	0.0045		0.4454		0.1960	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.59C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	106	109	110	114	107	110
NOT HISPANIC OR LATINO¹	92	97	92	101	91	97
White	204	180	152	145	133	129
Black or African American	70	69	46	52	40	47
American Indian or Alaska Native	19	14	17	10	14	10
Native Hawaiian or Other Pacific Islander	14	12	12	9	11	8
Chinese	17	19	12	16	11	13
Filipino	15	19	13	15	12	12
Japanese	*	*	*	*	*	*
Asian Indian	17	17	14	13	10	12
Korean	22	14	17	12	16	11
Vietnamese	14	10	10	8	8	8
HISPANIC OR LATINO¹	48	50	51	52	48	49
Mexican	85	78	55	59	47	50
Puerto Rican	26	22	21	18	19	17
Central or South American	30	30	20	24	18	20
Cuban	13	12	11	9	10	7

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.59D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.34	0.34	0.35	0.35	0.34	0.34
NOT HISPANIC OR LATINO¹	0.36	0.36	0.36	0.38	0.35	0.37
White	0.36	0.38	0.39	0.44	0.40	0.43
Black or African American	0.92	0.88	0.86	0.86	0.79	0.83
American Indian or Alaska Native	2.54	3.15	3.49	3.41	3.60	3.29
Native Hawaiian or Other Pacific Islander	3.93	4.48	5.17	5.57	4.89	5.19
Chinese	4.22	4.31	3.25	3.73	3.07	3.39
Filipino	3.74	4.06	3.86	4.58	3.95	4.03
Japanese	*	*	*	*	*	*
Asian Indian	3.38	3.28	3.47	2.85	3.06	2.80
Korean	5.20	4.76	4.96	4.99	5.08	4.91
Vietnamese	5.39	5.11	5.11	4.56	4.31	4.47
HISPANIC OR LATINO¹	0.85	0.87	0.91	0.91	0.85	0.86
Mexican	1.05	1.08	1.13	1.10	1.04	1.02
Puerto Rican	2.50	2.63	2.90	2.61	2.66	2.55
Central or South American	2.29	2.34	2.04	2.62	2.06	2.37
Cuban	4.58	4.28	4.94	4.50	4.75	4.24

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.59P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.0000		0.0564		0.0110	
NOT HISPANIC OR LATINO¹	0.0000		0.0339		0.0055	
White	0.0000		0.0376		0.0074	
Black or African American	0.2018		0.5656		0.6019	
American Indian or Alaska Native	0.7806		0.1916		0.6816	
Native Hawaiian or Other Pacific Islander	0.9269		0.6911		0.6020	
Chinese	0.8685		0.5153		0.8252	
Filipino	0.0538		0.1438		0.9041	
Japanese	*		*		*	
Asian Indian	0.2369		0.6286		0.6439	
Korean	0.0351		0.0810		0.0783	
Vietnamese	0.0292		0.5178		0.7611	
HISPANIC OR LATINO¹	0.0060		0.8790		0.7359	
Mexican	0.0409		0.5846		0.8581	
Puerto Rican	0.2102		0.6346		0.4521	
Central or South American	0.0275		0.0550		0.0806	
Cuban	0.8808		0.6903		0.3679	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.60C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	570	562	549	527	521	511
NOT HISPANIC OR LATINO¹	522	524	521	504	496	488
White	1,178	1,028	552	496	518	467
Black or African American	424	377	236	225	222	213
American Indian or Alaska Native	80	80	38	43	35	39
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	27
Chinese	90	127	31	38	27	31
Filipino	96	112	45	42	41	41
Japanese	85	*	26	14	24	16
Asian Indian	91	67	60	30	56	24
Korean	*	*	*	23	*	21
Vietnamese	*	47	*	35	*	*
HISPANIC OR LATINO¹	201	218	181	185	172	171
Mexican	268	273	160	164	151	148
Puerto Rican	127	127	79	84	77	78
Central or South American	139	129	83	71	72	66
Cuban	103	72	76	48	60	43

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.60D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.32	0.31	0.30	0.29	0.29	0.28
NOT HISPANIC OR LATINO¹	0.33	0.32	0.33	0.31	0.31	0.30
White	0.33	0.33	0.35	0.34	0.34	0.32
Black or African American	1.25	0.98	0.95	0.91	0.91	0.87
American Indian or Alaska Native	3.01	3.79	3.39	4.34	3.39	3.98
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	5.17
Chinese	4.75	5.07	1.99	2.02	1.70	1.61
Filipino	4.41	5.05	3.19	2.71	2.69	2.63
Japanese	4.95	*	3.27	1.71	3.05	1.92
Asian Indian	4.46	3.30	3.31	1.65	3.20	1.34
Korean	*	*	*	4.41	*	4.01
Vietnamese	*	5.69	*	5.34	*	*
HISPANIC OR LATINO¹	1.01	1.01	0.91	0.86	0.87	0.79
Mexican	1.29	1.24	1.02	1.04	1.00	0.96
Puerto Rican	2.56	3.10	3.05	3.02	3.02	2.96
Central or South American	3.05	2.49	2.49	2.00	2.31	1.89
Cuban	4.66	3.91	5.02	4.19	4.10	3.77

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.60P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.0002		0.1282		0.0672	
NOT HISPANIC OR LATINO¹	0.0022		0.0995		0.0799	
White	0.0333		0.4965		0.4332	
Black or African American	0.0868		0.0839		0.0735	
American Indian or Alaska Native	0.6298		0.6074		0.5833	
Native Hawaiian or Other Pacific Islander	*		*		*	
Chinese	0.5740		0.5638		0.6093	
Filipino	0.5178		0.5829		0.8957	
Japanese	*		0.0276		0.0491	
Asian Indian	0.1313		0.0174		0.0168	
Korean	*		*		*	
Vietnamese	*		*		*	
HISPANIC OR LATINO¹	0.1004		0.8418		0.7259	
Mexican	0.1911		0.9201		0.4472	
Puerto Rican	0.7381		0.5460		0.8855	
Central or South American	0.0408		0.1518		0.4599	
Cuban	0.9895		0.9709		0.9381	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.61C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	585	582	583	564	554	543
NOT HISPANIC OR LATINO¹	536	538	552	537	524	517
White	1,220	1,058	602	538	563	504
Black or African American	449	395	251	236	233	222
American Indian or Alaska Native	88	84	45	46	39	42
Native Hawaiian or Other Pacific Islander	57	47	41	33	32	28
Chinese	93	132	34	40	29	33
Filipino	97	116	49	46	44	42
Japanese	85	*	26	16	24	17
Asian Indian	92	71	62	35	57	28
Korean	*	39	39	26	38	24
Vietnamese	47	48	34	36	33	35
HISPANIC OR LATINO¹	210	225	193	196	183	181
Mexican	299	292	181	180	167	160
Puerto Rican	132	131	82	88	80	81
Central or South American	151	134	89	76	74	69
Cuban	104	73	76	50	60	43

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.61D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.28	0.27	0.28	0.26	0.26	0.25
NOT HISPANIC OR LATINO¹	0.29	0.28	0.30	0.28	0.28	0.27
White	0.29	0.29	0.33	0.31	0.31	0.30
Black or African American	1.05	0.83	0.80	0.76	0.77	0.73
American Indian or Alaska Native	2.52	3.13	2.90	3.79	2.85	3.52
Native Hawaiian or Other Pacific Islander	4.66	5.33	5.73	4.79	4.70	4.27
Chinese	4.14	4.37	1.88	1.81	1.61	1.50
Filipino	3.93	4.31	2.98	2.53	2.50	2.32
Japanese	4.64	*	3.08	1.82	2.83	1.94
Asian Indian	3.79	2.82	2.84	1.55	2.73	1.27
Korean	*	4.66	4.70	3.54	4.54	3.26
Vietnamese	5.29	4.58	4.08	4.26	3.90	4.15
HISPANIC OR LATINO¹	0.82	0.82	0.76	0.72	0.72	0.66
Mexican	1.01	0.99	0.85	0.86	0.82	0.78
Puerto Rican	2.05	2.58	2.45	2.56	2.43	2.53
Central or South American	2.51	2.07	2.06	1.70	1.89	1.60
Cuban	3.95	3.46	4.20	3.79	3.41	3.40

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.61P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.0000		0.0807		0.0290	
NOT HISPANIC OR LATINO¹	0.0001		0.0659		0.0371	
White	0.0043		0.3864		0.2695	
Black or African American	0.0568		0.1190		0.1017	
American Indian or Alaska Native	0.6033		0.4878		0.5682	
Native Hawaiian or Other Pacific Islander	0.5251		0.7692		0.7790	
Chinese	0.5974		0.4704		0.6823	
Filipino	0.3964		0.8214		0.9040	
Japanese	*		0.0358		0.0553	
Asian Indian	0.0964		0.0180		0.0281	
Korean	*		0.2622		0.1275	
Vietnamese	0.8894		0.7477		0.5967	
HISPANIC OR LATINO¹	0.0264		0.9740		0.6123	
Mexican	0.0740		0.8667		0.4747	
Puerto Rican	0.5538		0.6785		0.9351	
Central or South American	0.0160		0.0648		0.2472	
Cuban	0.9891		0.9020		0.9327	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.62C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	457	479	655	637	688	683
NOT HISPANIC OR LATINO¹	416	436	613	596	651	635
White	318	335	526	500	578	547
Black or African American	168	190	214	217	195	222
American Indian or Alaska Native	85	89	70	59	62	46
Native Hawaiian or Other Pacific Islander	55	50	43	44	40	36
Chinese	126	151	110	142	93	117
Filipino	157	127	128	113	108	91
Japanese	98	*	72	87	63	85
Asian Indian	119	104	108	88	89	69
Korean	80	55	74	48	55	42
Vietnamese	55	58	53	52	42	45
HISPANIC OR LATINO¹	176	201	240	230	233	223
Mexican	421	378	358	328	279	247
Puerto Rican	152	153	131	139	100	103
Central or South American	165	183	155	147	123	119
Cuban	135	93	97	81	80	73

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.62D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.19	0.20	0.28	0.26	0.29	0.28
NOT HISPANIC OR LATINO¹	0.20	0.21	0.30	0.28	0.31	0.30
White	0.19	0.20	0.32	0.30	0.35	0.33
Black or African American	0.62	0.67	0.79	0.77	0.72	0.79
American Indian or Alaska Native	2.93	2.12	3.05	3.85	3.53	3.17
Native Hawaiian or Other Pacific Islander	3.85	4.83	4.37	4.97	4.26	4.30
Chinese	2.76	3.64	3.49	3.85	3.28	3.70
Filipino	3.13	3.87	3.96	4.07	3.60	3.50
Japanese	4.07	*	5.23	5.64	5.35	5.82
Asian Indian	3.15	2.79	3.31	2.79	3.02	2.35
Korean	4.82	3.81	5.08	4.21	5.04	4.16
Vietnamese	5.44	4.91	5.17	5.02	4.39	4.55
HISPANIC OR LATINO¹	0.60	0.64	0.82	0.73	0.79	0.71
Mexican	0.71	0.78	0.96	0.91	0.92	0.84
Puerto Rican	1.79	1.84	2.42	2.28	2.31	2.34
Central or South American	1.74	1.80	2.17	2.07	2.06	2.10
Cuban	2.22	2.33	2.94	3.30	3.36	3.00

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.62P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.1016		0.5331		0.1790	
NOT HISPANIC OR LATINO¹	0.2536		0.3316		0.0668	
White	0.8343		0.0624		0.0150	
Black or African American	0.0728		0.6379		0.9522	
American Indian or Alaska Native	0.6711		0.1985		0.3974	
Native Hawaiian or Other Pacific Islander	0.3356		0.6143		0.4708	
Chinese	0.6100		0.7136		0.8229	
Filipino	0.4254		0.8822		0.8042	
Japanese	*		0.4734		0.3373	
Asian Indian	0.4854		0.0620		0.0178	
Korean	0.5800		0.9972		1.0000	
Vietnamese	0.5493		0.8630		0.4767	
HISPANIC OR LATINO¹	0.3388		0.6570		0.5031	
Mexican	0.2774		0.3926		0.3150	
Puerto Rican	0.7207		0.9668		0.8200	
Central or South American	0.6282		0.6897		0.3425	
Cuban	0.7257		0.0916		0.2664	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.63C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	80	78	76	73	60	57
NOT HISPANIC OR LATINO¹	72	68	69	64	54	51
White	97	92	84	82	58	54
Black or African American	44	45	33	36	22	21
American Indian or Alaska Native	10	8	9	6	6	3
Native Hawaiian or Other Pacific Islander	6	*	6	5	3	3
Chinese	8	10	8	9	5	5
Filipino	7	6	6	6	5	4
Japanese	*	*	*	*	2	*
Asian Indian	8	10	7	10	5	5
Korean	*	7	*	5	3	3
Vietnamese	9	*	*	*	3	*
HISPANIC OR LATINO¹	36	37	33	35	25	29
Mexican	53	47	41	41	23	28
Puerto Rican	20	16	16	14	8	9
Central or South American	16	15	14	14	10	10
Cuban	9	8	8	8	6	6

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.63D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.32	0.31	0.30	0.29	0.24	0.23
NOT HISPANIC OR LATINO¹	0.35	0.32	0.33	0.31	0.26	0.24
White	0.40	0.37	0.37	0.36	0.30	0.29
Black or African American	0.80	0.78	0.69	0.68	0.53	0.48
American Indian or Alaska Native	3.85	3.42	3.17	2.78	2.73	1.92
Native Hawaiian or Other Pacific Islander	5.80	*	5.85	5.74	4.09	3.76
Chinese	3.72	4.07	3.35	3.78	2.07	2.63
Filipino	3.08	3.78	3.08	3.33	2.90	2.11
Japanese	*	*	*	*	3.19	*
Asian Indian	3.30	3.34	3.15	3.33	2.22	1.87
Korean	*	5.21	*	4.33	3.27	2.76
Vietnamese	5.48	*	*	*	2.69	*
HISPANIC OR LATINO¹	0.91	0.87	0.83	0.82	0.64	0.67
Mexican	1.16	1.10	1.02	1.03	0.73	0.84
Puerto Rican	2.26	2.29	2.25	2.18	1.69	1.71
Central or South American	2.52	2.61	2.38	2.60	2.13	2.11
Cuban	4.45	4.44	4.57	4.46	4.38	3.71

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.63P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.0001		0.0540		0.0715	
NOT HISPANIC OR LATINO¹	0.0004		0.0588		0.0131	
White	0.0014		0.0764		0.0098	
Black or African American	0.2446		0.8541		0.8143	
American Indian or Alaska Native	0.2460		0.1596		0.1986	
Native Hawaiian or Other Pacific Islander	*		0.5152		0.5737	
Chinese	0.0761		0.1098		0.3924	
Filipino	0.0012		0.0865		0.0550	
Japanese	*		*		*	
Asian Indian	0.7096		0.6872		0.6628	
Korean	*		*		0.8704	
Vietnamese	*		*		*	
HISPANIC OR LATINO¹	0.0450		0.5848		0.3191	
Mexican	0.1305		0.7223		0.3164	
Puerto Rican	0.1376		0.3190		0.7186	
Central or South American	0.5678		0.6834		0.7197	
Cuban	0.2599		0.6688		0.7774	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.64C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	77	82	99	99	124	117
NOT HISPANIC OR LATINO¹	68	67	84	82	109	100
White	233	213	223	204	204	184
Black or African American	100	87	91	78	65	63
American Indian or Alaska Native	20	14	16	13	14	9
Native Hawaiian or Other Pacific Islander	15	14	14	12	12	10
Chinese	30	28	23	26	19	18
Filipino	18	20	16	18	14	15
Japanese	*	*	*	*	*	*
Asian Indian	18	20	17	19	15	16
Korean	22	18	22	17	20	14
Vietnamese	21	*	19	*	*	10
HISPANIC OR LATINO¹	35	43	45	50	47	52
Mexican	106	92	87	81	71	66
Puerto Rican	29	25	27	23	25	20
Central or South American	34	38	29	33	26	27
Cuban	14	14	14	13	12	12

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.64D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.25	0.25	0.31	0.31	0.40	0.36
NOT HISPANIC OR LATINO¹	0.26	0.25	0.32	0.31	0.42	0.37
White	0.27	0.25	0.33	0.31	0.45	0.41
Black or African American	0.68	0.69	0.80	0.79	0.85	0.92
American Indian or Alaska Native	2.10	3.44	3.66	3.31	3.51	3.13
Native Hawaiian or Other Pacific Islander	4.20	3.23	4.73	4.39	4.87	4.85
Chinese	3.28	3.27	3.12	4.17	4.11	4.05
Filipino	2.59	3.47	4.34	3.97	3.80	4.23
Japanese	*	*	*	*	*	*
Asian Indian	3.80	3.58	3.90	3.50	3.89	3.62
Korean	4.36	3.60	4.55	4.55	5.61	5.20
Vietnamese	4.20	*	5.40	*	*	5.43
HISPANIC OR LATINO¹	0.63	0.74	0.81	0.87	0.84	0.90
Mexican	0.78	0.94	0.99	1.06	1.02	1.10
Puerto Rican	1.60	2.19	1.94	2.44	2.50	2.68
Central or South American	1.85	1.84	2.10	2.38	2.28	2.40
Cuban	2.06	2.93	2.57	3.30	3.49	4.76

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.64P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.0101		0.9911		0.6806	
NOT HISPANIC OR LATINO¹	0.0859		0.3883		0.8880	
White	0.4794		0.3455		0.5223	
Black or African American	0.1379		0.7621		0.7233	
American Indian or Alaska Native	0.2366		0.9061		0.2104	
Native Hawaiian or Other Pacific Islander	0.4192		0.5934		0.4928	
Chinese	0.8261		0.3567		0.6150	
Filipino	0.9728		0.1925		0.7385	
Japanese	*		*		*	
Asian Indian	0.7542		0.8761		1.0000	
Korean	0.5030		0.8623		0.3923	
Vietnamese	*		*		*	
HISPANIC OR LATINO¹	0.0226		0.1231		0.1778	
Mexican	0.0490		0.3538		0.5009	
Puerto Rican	0.0364		0.0267		0.0986	
Central or South American	0.9867		0.3773		0.1029	
Cuban	0.7116		0.9055		0.5149	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.65C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	434	455	636	619	661	661
NOT HISPANIC OR LATINO¹	393	420	598	588	627	620
White	1,315	1,157	1,148	1,003	1,010	895
Black or African American	442	440	363	338	292	289
American Indian or Alaska Native	72	83	60	53	57	43
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	34
Chinese	122	143	107	136	90	115
Filipino	154	120	124	108	110	88
Japanese	*	*	*	*	*	*
Asian Indian	117	97	106	82	88	64
Korean	*	*	*	*	*	*
Vietnamese	*	*	*	*	*	*
HISPANIC OR LATINO¹	175	187	234	218	228	213
Mexican	355	341	307	303	249	232
Puerto Rican	139	143	120	131	93	99
Central or South American	148	172	139	137	111	111
Cuban	127	86	92	76	78	69

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.65D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.24	0.25	0.35	0.34	0.37	0.36
NOT HISPANIC OR LATINO¹	0.25	0.26	0.37	0.36	0.39	0.38
White	0.23	0.24	0.40	0.37	0.43	0.41
Black or African American	0.83	0.89	1.07	1.05	0.96	1.06
American Indian or Alaska Native	3.57	2.99	3.80	5.24	4.64	4.40
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	5.89
Chinese	3.33	4.64	4.38	4.82	4.12	4.70
Filipino	3.89	4.72	4.85	5.09	4.66	4.42
Japanese	*	*	*	*	*	*
Asian Indian	3.91	3.66	4.11	3.61	3.83	3.02
Korean	*	*	*	*	*	*
Vietnamese	*	*	*	*	*	*
HISPANIC OR LATINO¹	0.88	0.87	1.17	1.01	1.15	0.99
Mexican	1.12	1.10	1.44	1.33	1.35	1.24
Puerto Rican	2.53	2.49	3.38	3.21	3.34	3.30
Central or South American	2.38	2.29	2.94	2.69	2.81	2.71
Cuban	2.76	2.80	3.80	4.19	4.20	3.69

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.65P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.5251		0.4270		0.1194	
NOT HISPANIC OR LATINO¹	0.6328		0.3473		0.0508	
White	0.4829		0.0720		0.0135	
Black or African American	0.1541		0.5952		0.9665	
American Indian or Alaska Native	0.4345		0.2158		0.5402	
Native Hawaiian or Other Pacific Islander	*		*		*	
Chinese	0.3289		0.9554		0.9164	
Filipino	0.6725		0.8346		0.6853	
Japanese	*		*		*	
Asian Indian	0.5386		0.0606		0.0151	
Korean	*		*		*	
Vietnamese	*		*		*	
HISPANIC OR LATINO¹	0.9640		0.9770		0.6560	
Mexican	0.8534		0.5929		0.3416	
Puerto Rican	0.8278		0.5452		0.5339	
Central or South American	0.5665		0.7938		0.4792	
Cuban	0.4567		0.0676		0.1754	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.66C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	442	464	651	632	679	678
NOT HISPANIC OR LATINO¹	404	427	612	596	643	636
White	1,367	1,188	1,190	1,034	1,042	929
Black or African American	489	469	406	365	320	310
American Indian or Alaska Native	80	87	66	57	61	45
Native Hawaiian or Other Pacific Islander	54	*	42	44	40	35
Chinese	128	149	111	141	94	117
Filipino	154	126	126	113	108	91
Japanese	*	*	72	*	63	*
Asian Indian	118	102	108	87	90	68
Korean	*	54	*	47	54	42
Vietnamese	*	57	*	52	42	45
HISPANIC OR LATINO¹	174	191	238	225	231	219
Mexican	404	366	343	321	272	244
Puerto Rican	146	150	127	137	99	102
Central or South American	162	180	151	145	122	117
Cuban	132	90	95	80	79	72

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.66D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.21	0.21	0.31	0.29	0.32	0.31
NOT HISPANIC OR LATINO¹	0.22	0.23	0.33	0.32	0.35	0.34
White	0.20	0.21	0.35	0.33	0.38	0.36
Black or African American	0.70	0.75	0.91	0.89	0.81	0.91
American Indian or Alaska Native	3.07	2.38	3.36	4.53	3.95	3.75
Native Hawaiian or Other Pacific Islander	4.45	*	5.01	5.49	4.84	4.83
Chinese	2.95	4.00	3.82	4.16	3.62	4.02
Filipino	3.31	4.15	4.30	4.45	3.90	3.81
Japanese	*	*	5.62	*	5.73	*
Asian Indian	3.39	3.16	3.58	3.13	3.28	2.65
Korean	*	4.26	*	4.85	5.62	4.82
Vietnamese	*	5.32	*	5.48	5.31	5.05
HISPANIC OR LATINO¹	0.68	0.70	0.93	0.83	0.91	0.80
Mexican	0.85	0.86	1.12	1.05	1.05	0.96
Puerto Rican	2.07	2.06	2.78	2.66	2.74	2.77
Central or South American	1.91	1.92	2.39	2.25	2.33	2.29
Cuban	2.45	2.53	3.38	3.71	3.79	3.33

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.66P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.2773		0.4259		0.1464	
NOT HISPANIC OR LATINO¹	0.4492		0.2761		0.0512	
White	0.5738		0.0502		0.0107	
Black or African American	0.1026		0.6515		0.9721	
American Indian or Alaska Native	0.5731		0.2414		0.4454	
Native Hawaiian or Other Pacific Islander	*		0.5751		0.4587	
Chinese	0.3601		0.9366		0.9746	
Filipino	0.6709		0.7079		0.6542	
Japanese	*		*		*	
Asian Indian	0.5116		0.0661		0.0180	
Korean	*		*		0.8276	
Vietnamese	*		*		0.5597	
HISPANIC OR LATINO¹	0.5827		0.7295		0.4408	
Mexican	0.4865		0.4404		0.2574	
Puerto Rican	0.8861		0.8744		0.8150	
Central or South American	0.5779		0.6585		0.3169	
Cuban	0.4978		0.0829		0.2277	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.67C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	688	683	512	495	286	269
NOT HISPANIC OR LATINO¹	651	635	493	459	265	252
White	578	547	430	408	239	238
Black or African American	195	222	161	168	89	78
American Indian or Alaska Native	62	46	35	41	20	29
Native Hawaiian or Other Pacific Islander	40	36	22	30	9	8
Chinese	93	117	35	42	9	21
Filipino	108	91	40	47	14	15
Japanese	63	85	19	28	4	5
Asian Indian	89	69	43	38	15	12
Korean	55	42	32	28	24	11
Vietnamese	42	45	19	21	11	12
HISPANIC OR LATINO¹	233	223	184	179	94	90
Mexican	279	247	198	169	77	76
Puerto Rican	100	103	73	73	37	34
Central or South American	123	119	81	69	24	31
Cuban	80	73	52	43	35	15

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.67D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.29	0.28	0.22	0.20	0.12	0.11
NOT HISPANIC OR LATINO¹	0.31	0.30	0.24	0.22	0.13	0.12
White	0.35	0.33	0.26	0.24	0.14	0.14
Black or African American	0.72	0.79	0.59	0.59	0.33	0.28
American Indian or Alaska Native	3.53	3.17	2.30	2.84	1.44	2.11
Native Hawaiian or Other Pacific Islander	4.26	4.30	3.01	3.80	1.53	1.20
Chinese	3.28	3.70	1.74	1.68	0.43	0.83
Filipino	3.60	3.50	1.89	2.34	0.67	0.84
Japanese	5.35	5.82	2.63	3.08	0.57	0.57
Asian Indian	3.02	2.35	1.82	1.50	0.72	0.47
Korean	5.04	4.16	3.47	3.18	2.74	1.44
Vietnamese	4.39	4.55	2.29	2.58	1.38	1.42
HISPANIC OR LATINO¹	0.79	0.71	0.63	0.57	0.32	0.28
Mexican	0.92	0.84	0.79	0.70	0.38	0.36
Puerto Rican	2.31	2.34	1.90	1.89	1.14	1.02
Central or South American	2.06	2.10	1.59	1.46	0.58	0.70
Cuban	3.36	3.00	3.31	2.73	2.80	1.07

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 2.67P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups:
P Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003**

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.1790		0.8760		0.9816	
NOT HISPANIC OR LATINO¹	0.0668		0.9195		0.8856	
White	0.0150		0.7337		0.5814	
Black or African American	0.9522		0.4439		0.7735	
American Indian or Alaska Native	0.3974		0.8652		0.9062	
Native Hawaiian or Other Pacific Islander	0.4708		0.9050		0.0372	
Chinese	0.8229		0.3722		0.0996	
Filipino	0.8042		0.4689		0.9880	
Japanese	0.3373		0.8990		0.4385	
Asian Indian	0.0178		0.7986		0.5146	
Korean	1.0000		0.4117		0.4904	
Vietnamese	0.4767		0.2867		0.9934	
HISPANIC OR LATINO¹	0.5031		0.4382		0.7574	
Mexican	0.3150		0.4936		0.8385	
Puerto Rican	0.8200		0.9831		0.5481	
Central or South American	0.3425		0.3233		0.4928	
Cuban	0.2664		0.8230		0.6728	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 2.68C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups:
Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005**

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	60	57	49	49	23	24
NOT HISPANIC OR LATINO¹	54	51	44	42	21	21
White	58	54	46	41	21	21
Black or African American	22	21	15	14	4	5
American Indian or Alaska Native	6	3	5	3	2	2
Native Hawaiian or Other Pacific Islander	3	3	2	2	*	0
Chinese	5	5	3	4	*	*
Filipino	5	4	3	2	*	0
Japanese	2	*	0	*	0	*
Asian Indian	5	5	3	4	*	*
Korean	3	3	2	2	*	*
Vietnamese	3	*	*	*	*	*
HISPANIC OR LATINO¹	25	29	19	26	8	12
Mexican	23	28	16	24	8	10
Puerto Rican	8	9	6	7	1	3
Central or South American	10	10	8	9	3	4
Cuban	6	6	*	4	1	2

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 2.68D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups:
Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005**

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.24	0.23	0.20	0.19	0.09	0.10
NOT HISPANIC OR LATINO¹	0.26	0.24	0.21	0.20	0.10	0.10
White	0.30	0.29	0.26	0.24	0.13	0.13
Black or African American	0.53	0.48	0.38	0.35	0.11	0.14
American Indian or Alaska Native	2.73	1.92	2.69	1.72	0.84	1.15
Native Hawaiian or Other Pacific Islander	4.09	3.76	3.12	2.84	*	0.36
Chinese	2.07	2.63	1.19	2.16	*	*
Filipino	2.90	2.11	1.88	1.26	*	0.16
Japanese	3.19	*	0.93	*	0.64	*
Asian Indian	2.22	1.87	1.34	1.61	*	*
Korean	3.27	2.76	2.46	1.60	*	*
Vietnamese	2.69	*	*	*	*	*
HISPANIC OR LATINO¹	0.64	0.67	0.48	0.61	0.20	0.28
Mexican	0.73	0.84	0.55	0.76	0.29	0.32
Puerto Rican	1.69	1.71	1.19	1.34	0.27	0.68
Central or South American	2.13	2.11	1.66	2.00	0.63	0.97
Cuban	4.38	3.71	*	2.59	0.97	1.31

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.68P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.0715		0.5699		0.7617	
NOT HISPANIC OR LATINO¹	0.0131		0.0889		0.3828	
White	0.0098		0.0424		0.2937	
Black or African American	0.8143		0.4664		0.5099	
American Indian or Alaska Native	0.1986		0.3498		0.3408	
Native Hawaiian or Other Pacific Islander	0.5737		0.6360		*	
Chinese	0.3924		0.1561		*	
Filipino	0.0550		0.2988		*	
Japanese	*		*		*	
Asian Indian	0.6628		0.9308		*	
Korean	0.8704		0.7182		*	
Vietnamese	*		*		*	
HISPANIC OR LATINO¹	0.3191		0.0468		0.2420	
Mexican	0.3164		0.0718		0.9483	
Puerto Rican	0.7186		0.9169		0.0799	
Central or South American	0.7197		0.6578		0.4444	
Cuban	0.7774		*		0.8140	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 2.69C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups:
Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005**

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	124	117	119	119	84	85
NOT HISPANIC OR LATINO¹	109	100	105	107	75	77
White	204	184	158	151	87	88
Black or African American	65	63	40	43	16	18
American Indian or Alaska Native	14	9	11	8	4	5
Native Hawaiian or Other Pacific Islander	12	10	11	7	7	4
Chinese	19	18	11	13	4	7
Filipino	14	15	10	11	6	4
Japanese	*	*	*	*	3	1
Asian Indian	15	16	9	11	5	8
Korean	20	14	13	10	*	6
Vietnamese	*	10	8	7	4	3
HISPANIC OR LATINO¹	47	52	48	50	29	29
Mexican	71	66	61	56	25	25
Puerto Rican	25	20	20	18	10	9
Central or South American	26	27	18	24	9	11
Cuban	12	12	10	9	4	6

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.69D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.40	0.36	0.38	0.37	0.27	0.26
NOT HISPANIC OR LATINO¹	0.42	0.37	0.41	0.40	0.29	0.29
White	0.45	0.41	0.44	0.45	0.34	0.35
Black or African American	0.85	0.92	0.74	0.77	0.39	0.39
American Indian or Alaska Native	3.51	3.13	3.39	3.00	1.86	2.31
Native Hawaiian or Other Pacific Islander	4.87	4.85	4.83	4.55	3.89	2.93
Chinese	4.11	4.05	3.09	3.40	1.25	1.91
Filipino	3.80	4.23	3.42	3.99	2.46	1.56
Japanese	*	*	*	*	3.71	1.64
Asian Indian	3.89	3.62	2.90	2.69	1.59	2.22
Korean	5.61	5.20	5.15	4.61	*	3.17
Vietnamese	*	5.43	3.87	3.85	2.52	2.26
HISPANIC OR LATINO¹	0.84	0.90	0.86	0.88	0.52	0.51
Mexican	1.02	1.10	1.02	1.07	0.59	0.61
Puerto Rican	2.50	2.68	2.63	2.68	1.58	1.65
Central or South American	2.28	2.40	2.10	2.40	1.20	1.40
Cuban	3.49	4.76	4.36	4.69	2.71	4.09

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.69P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.6806		0.5781		0.6334	
NOT HISPANIC OR LATINO¹	0.8880		0.5826		0.4596	
White	0.5223		0.3200		0.2864	
Black or African American	0.7233		0.8049		0.7138	
American Indian or Alaska Native	0.2104		0.4975		0.5967	
Native Hawaiian or Other Pacific Islander	0.4928		0.2297		0.9717	
Chinese	0.6150		0.9568		0.2884	
Filipino	0.7385		0.8102		0.5288	
Japanese	*		*		0.0586	
Asian Indian	1.0000		0.6190		0.8076	
Korean	0.3923		0.9179		*	
Vietnamese	*		0.7339		0.8195	
HISPANIC OR LATINO¹	0.1778		0.8819		0.5485	
Mexican	0.5009		0.9199		0.1948	
Puerto Rican	0.0986		0.8520		0.9446	
Central or South American	0.1029		0.9886		0.7804	
Cuban	0.5149		0.0660		0.1029	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 2.70C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups:
Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005**

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	661	661	478	461	257	249
NOT HISPANIC OR LATINO¹	627	620	459	427	235	233
White	1,010	895	468	438	225	226
Black or African American	292	289	193	181	89	79
American Indian or Alaska Native	57	43	31	38	19	28
Native Hawaiian or Other Pacific Islander	*	34	18	28	6	7
Chinese	90	115	31	38	*	19
Filipino	110	88	39	44	13	15
Japanese	*	*	18	28	3	4
Asian Indian	88	64	43	36	14	8
Korean	*	*	*	25	*	9
Vietnamese	*	*	*	19	*	*
HISPANIC OR LATINO¹	228	213	177	170	90	85
Mexican	249	232	173	158	74	72
Puerto Rican	93	99	68	69	36	31
Central or South American	111	111	75	59	22	29
Cuban	78	69	53	41	*	15

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.70D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.37	0.36	0.27	0.25	0.14	0.13
NOT HISPANIC OR LATINO¹	0.39	0.38	0.29	0.26	0.15	0.14
White	0.43	0.41	0.31	0.29	0.16	0.17
Black or African American	0.96	1.06	0.80	0.81	0.45	0.38
American Indian or Alaska Native	4.64	4.40	2.91	3.89	1.91	2.91
Native Hawaiian or Other Pacific Islander	*	5.89	4.27	5.28	1.88	1.42
Chinese	4.12	4.70	2.05	1.95	*	1.01
Filipino	4.66	4.42	2.33	2.86	0.78	1.03
Japanese	*	*	2.83	3.41	0.48	0.61
Asian Indian	3.83	3.02	2.45	1.88	0.89	0.42
Korean	*	*	*	4.75	*	1.88
Vietnamese	*	*	*	3.40	*	*
HISPANIC OR LATINO¹	1.15	0.99	0.89	0.79	0.45	0.39
Mexican	1.35	1.24	1.13	1.03	0.57	0.52
Puerto Rican	3.34	3.30	2.70	2.64	1.63	1.35
Central or South American	2.81	2.71	2.15	1.75	0.76	0.89
Cuban	4.20	3.69	4.33	3.37	*	1.31

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 2.70P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups:
P Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003**

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.1194		0.7286		0.8186	
NOT HISPANIC OR LATINO¹	0.0508		0.9771		0.8860	
White	0.0135		0.8979		0.8460	
Black or African American	0.9665		0.3560		0.6814	
American Indian or Alaska Native	0.5402		0.7059		0.9523	
Native Hawaiian or Other Pacific Islander	*		0.4666		0.0399	
Chinese	0.9164		0.4451		*	
Filipino	0.6853		0.3921		0.7097	
Japanese	*		0.8514		0.9844	
Asian Indian	0.0151		0.6931		0.4433	
Korean	*		*		*	
Vietnamese	*		*		*	
HISPANIC OR LATINO¹	0.6560		0.3279		0.8574	
Mexican	0.3416		0.3661		0.8110	
Puerto Rican	0.5339		0.8680		0.4987	
Central or South American	0.4792		0.2748		0.5761	
Cuban	0.1754		0.8179		*	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.71C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	679	678	507	495	285	269
NOT HISPANIC OR LATINO¹	643	636	489	461	262	253
White	1,042	929	503	479	249	246
Black or African American	320	310	204	192	91	81
American Indian or Alaska Native	61	45	35	40	20	29
Native Hawaiian or Other Pacific Islander	40	35	22	30	9	8
Chinese	94	117	35	41	9	21
Filipino	108	91	40	47	15	15
Japanese	63	*	19	28	4	5
Asian Indian	90	68	43	38	15	12
Korean	54	42	32	28	24	11
Vietnamese	42	45	18	21	11	12
HISPANIC OR LATINO¹	231	219	182	176	94	89
Mexican	272	244	195	169	77	75
Puerto Rican	99	102	73	73	37	34
Central or South American	122	117	81	67	24	31
Cuban	79	72	51	43	*	15

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 2.71D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.32	0.31	0.24	0.23	0.13	0.12
NOT HISPANIC OR LATINO¹	0.35	0.34	0.26	0.24	0.14	0.13
White	0.38	0.36	0.28	0.27	0.16	0.16
Black or African American	0.81	0.91	0.67	0.69	0.38	0.32
American Indian or Alaska Native	3.95	3.75	2.56	3.33	1.64	2.42
Native Hawaiian or Other Pacific Islander	4.84	4.83	3.55	4.31	1.72	1.36
Chinese	3.62	4.02	1.94	1.80	0.49	0.90
Filipino	3.90	3.81	2.01	2.56	0.74	0.93
Japanese	5.73	*	2.82	3.18	0.60	0.59
Asian Indian	3.28	2.65	2.03	1.67	0.80	0.52
Korean	5.62	4.82	3.86	3.76	3.08	1.70
Vietnamese	5.31	5.05	2.70	2.89	1.61	1.60
HISPANIC OR LATINO¹	0.91	0.80	0.71	0.65	0.37	0.33
Mexican	1.05	0.96	0.91	0.81	0.44	0.42
Puerto Rican	2.74	2.77	2.25	2.23	1.33	1.19
Central or South American	2.33	2.29	1.78	1.58	0.65	0.77
Cuban	3.79	3.33	3.69	3.05	*	1.19

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 2.71P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups:
P Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003**

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL¹	0.1464		0.8950		0.9868	
NOT HISPANIC OR LATINO¹	0.0512		0.8369		0.8531	
White	0.0107		0.6382		0.5419	
Black or African American	0.9721		0.4044		0.7428	
American Indian or Alaska Native	0.4454		0.8051		0.9691	
Native Hawaiian or Other Pacific Islander	0.4587		0.9200		0.0537	
Chinese	0.9746		0.4686		0.1262	
Filipino	0.6542		0.3951		0.9035	
Japanese	*		0.9970		0.4487	
Asian Indian	0.0180		0.7953		0.5635	
Korean	0.8276		0.3240		0.5320	
Vietnamese	0.5597		0.3416		0.9759	
HISPANIC OR LATINO¹	0.4408		0.3136		0.6673	
Mexican	0.2574		0.3649		0.8421	
Puerto Rican	0.8150		0.9912		0.4618	
Central or South American	0.3169		0.2749		0.5709	
Cuban	0.2277		0.9215		*	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.