

Table 3.1C Perceived Risk and Availability of Substances, by Age Group: Standard Errors of Numbers in Thousands, 2004 and 2005

Risk/Availability ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
PERCEPTIONS OF GREAT RISK								
Cigarettes								
Smoke One or More Packs Per Day	720	762	104	110	134	134	694	733
Marijuana								
Smoke Once a Month	897	927	109	107	137	133	863	896
Smoke Once or Twice a Week	905	873	112	117	150	156	858	834
Cocaine								
Use Once a Month	735	748	106	114	138	145	702	708
Use Once or Twice a Week	488	508	92	95	107	106	474	475
Heroin								
Try Once or Twice	619	608	103	110	117	120	589	567
Use Once or Twice a Week	385	384	86	86	75	77	368	358
LSD								
Try Once or Twice	765	753	105	115	140	145	726	700
Use Once or Twice a Week	483	545	88	95	101	105	472	513
Alcohol								
Have Four or Five Drinks Nearly Every Day	792	799	112	108	143	148	773	757
Have Five or More Drinks Once or Twice a Week	855	903	101	118	149	155	834	868
PERCEIVED AVAILABILITY								
Fairly or Very Easy to Obtain								
Marijuana	880	938	108	110	139	132	855	908
Cocaine	738	807	102	99	147	146	718	776
Crack	717	820	100	96	134	141	703	791
Heroin	614	665	80	74	112	115	591	639
LSD	604	655	84	79	113	116	586	635
Approached in the Past Month by Someone Selling Drugs	380	359	85	86	122	116	341	306

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.1D Perceived Risk and Availability of Substances, by Age Group: Standard Errors of Percentages, 2004 and 2005

Risk/Availability ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
PERCEPTIONS OF GREAT RISK								
Cigarettes								
Smoke One or More Packs Per Day	0.30	0.32	0.42	0.44	0.42	0.41	0.38	0.40
Marijuana								
Smoke Once a Month	0.38	0.39	0.44	0.43	0.43	0.41	0.48	0.49
Smoke Once or Twice a Week	0.38	0.37	0.45	0.47	0.47	0.48	0.48	0.46
Cocaine								
Use Once a Month	0.31	0.31	0.43	0.46	0.43	0.45	0.39	0.39
Use Once or Twice a Week	0.21	0.21	0.37	0.38	0.33	0.33	0.26	0.26
Heroin								
Try Once or Twice	0.26	0.25	0.42	0.44	0.37	0.38	0.33	0.31
Use Once or Twice a Week	0.16	0.16	0.35	0.35	0.24	0.24	0.20	0.20
LSD								
Try Once or Twice	0.33	0.32	0.44	0.48	0.44	0.45	0.40	0.39
Use Once or Twice a Week	0.21	0.23	0.36	0.39	0.32	0.33	0.26	0.28
Alcohol								
Have Four or Five Drinks Nearly Every Day	0.33	0.33	0.45	0.43	0.45	0.46	0.43	0.41
Have Five or More Drinks Once or Twice a Week	0.36	0.37	0.40	0.47	0.46	0.48	0.46	0.47
PERCEIVED AVAILABILITY								
Fairly or Very Easy to Obtain								
Marijuana	0.38	0.40	0.44	0.44	0.44	0.41	0.49	0.51
Cocaine	0.33	0.35	0.42	0.41	0.47	0.46	0.42	0.45
Crack	0.32	0.36	0.41	0.39	0.43	0.45	0.41	0.46
Heroin	0.27	0.29	0.33	0.31	0.36	0.37	0.35	0.37
LSD	0.27	0.29	0.35	0.33	0.37	0.37	0.34	0.37
Approached in the Past Month by Someone Selling Drugs	0.16	0.15	0.34	0.34	0.38	0.36	0.19	0.17

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.1P Perceived Risk and Availability of Substances, by Age Group: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Risk/Availability ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
PERCEPTIONS OF GREAT RISK								
Cigarettes								
Smoke One or More Packs Per Day	0.6681		0.1944		0.8937		0.7903	
Marijuana								
Smoke Once a Month	0.0085		0.1024		0.0518		0.0248	
Smoke Once or Twice a Week	0.0659		0.6472		0.0931		0.0909	
Cocaine								
Use Once a Month	0.5147		0.2035		0.7110		0.3671	
Use Once or Twice a Week	0.1843		0.8690		0.9850		0.1832	
Heroin								
Try Once or Twice	0.1217		0.3767		0.9850		0.0849	
Use Once or Twice a Week	0.6574		0.5077		0.1871		0.9823	
LSD								
Try Once or Twice	0.1308		0.1433		0.4752		0.0565	
Use Once or Twice a Week	0.9082		0.5337		0.6578		0.7876	
Alcohol								
Have Four or Five Drinks Nearly Every Day	0.2211		0.0012		0.9026		0.4426	
Have Five or More Drinks Once or Twice a Week	0.2203		0.5955		0.0777		0.3834	
PERCEIVED AVAILABILITY								
Fairly or Very Easy to Obtain								
Marijuana	0.6270		0.0391		0.4442		0.7236	
Cocaine	0.1985		0.3547		0.1140		0.0783	
Crack	0.5088		0.6622		0.3097		0.3523	
Heroin	0.3089		0.9539		0.9411		0.2872	
LSD	0.0794		0.0120		0.0582		0.2631	
Approached in the Past Month by Someone Selling Drugs	0.3009		0.0999		0.9425		0.4762	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.2C Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Smoke One or More Packs of Cigarettes Per Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	104	110	104	110
AGE				
12-13	115	111	70	75
14-15	118	123	83	77
16-17	115	119	70	73
GENDER				
Male	77	79	77	79
Female	68	70	68	70
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	90	95	90	95
White	170	183	104	102
Black or African American	92	104	57	56
American Indian or Alaska Native	12	12	9	12
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	49	56	27	38
Two or More Races	25	22	14	11
Hispanic or Latino	53	52	53	52
COUNTY TYPE				
Large Metro	186	194	108	113
Small Metro	133	127	83	75
Nonmetro	97	93	55	57
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	57	54	49	44
Marijuana	46	45	41	37
Illicit Drugs Other Than Marijuana ²	39	33	35	32
Cigarettes	54	56	56	50
Alcohol	75	75	57	55
Binge Alcohol ³	59	53	49	45
PAST YEAR				
Illicit Drugs ²	84	81	62	64
Marijuana	67	65	53	52
Illicit Drugs Other Than Marijuana ²	62	59	48	49
Cigarettes	74	73	65	65
Alcohol	115	115	78	78
LIFETIME				
Illicit Drugs ²	101	100	76	75
Marijuana	76	75	60	60
Illicit Drugs Other Than Marijuana ²	83	78	63	63
Cigarettes	98	98	80	77
Alcohol	130	130	89	86

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.2D Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Smoke One or More Packs of Cigarettes Per Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.42	0.44	0.42	0.44
AGE				
12-13	0.71	0.73	0.71	0.73
14-15	0.75	0.75	0.75	0.75
16-17	0.72	0.73	0.72	0.73
GENDER				
Male	0.60	0.61	0.60	0.61
Female	0.56	0.57	0.56	0.57
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.43	0.46	0.43	0.46
White	0.51	0.50	0.51	0.50
Black or African American	1.09	1.14	1.09	1.14
American Indian or Alaska Native	4.76	4.68	4.76	4.68
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.15	2.47	2.15	2.47
Two or More Races	2.76	2.59	2.76	2.59
Hispanic or Latino	1.28	1.21	1.28	1.21
COUNTY TYPE				
Large Metro	0.62	0.62	0.62	0.62
Small Metro	0.66	0.75	0.66	0.75
Nonmetro	0.92	0.95	0.92	0.95
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.31	0.30	0.54	0.48
Marijuana	0.25	0.26	0.46	0.43
Illicit Drugs Other Than Marijuana ²	0.22	0.19	0.40	0.39
Cigarettes	0.30	0.32	0.61	0.56
Alcohol	0.39	0.39	0.60	0.60
Binge Alcohol ³	0.33	0.30	0.55	0.50
PAST YEAR				
Illicit Drugs ²	0.43	0.41	0.64	0.66
Marijuana	0.36	0.35	0.57	0.57
Illicit Drugs Other Than Marijuana ²	0.33	0.32	0.53	0.55
Cigarettes	0.39	0.41	0.67	0.67
Alcohol	0.51	0.51	0.74	0.74
LIFETIME				
Illicit Drugs ²	0.46	0.47	0.73	0.72
Marijuana	0.40	0.39	0.63	0.63
Illicit Drugs Other Than Marijuana ²	0.40	0.40	0.65	0.65
Cigarettes	0.47	0.51	0.77	0.73
Alcohol	0.53	0.55	0.79	0.76

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.2P Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day (II), among Persons Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

(I) Demographic Characteristic and (II) Substance Use	Smoke One or More Packs of Cigarettes Per Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.1944		0.1944	
AGE				
12-13	0.2527		0.2527	
14-15	0.5400		0.5400	
16-17	0.6020		0.6020	
GENDER				
Male	0.0623		0.0623	
Female	0.9460		0.9460	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.2523		0.2523	
White	0.3536		0.3536	
Black or African American	0.1534		0.1534	
American Indian or Alaska Native	0.1805		0.1805	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.1031		0.1031	
Two or More Races	0.2357		0.2357	
Hispanic or Latino	0.5034		0.5034	
COUNTY TYPE				
Large Metro	0.3580		0.3580	
Small Metro	0.0461		0.0461	
Nonmetro	0.2474		0.2474	
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.0238		0.9695	
Marijuana	0.0094		0.6174	
Illicit Drugs Other Than Marijuana ²	0.1180		0.9215	
Cigarettes	0.0333		0.1269	
Alcohol	0.0414		0.3434	
Binge Alcohol ³	0.0027		0.3372	
PAST YEAR				
Illicit Drugs ²	0.0052		0.9621	
Marijuana	0.0009		0.9780	
Illicit Drugs Other Than Marijuana ²	0.1269		0.6762	
Cigarettes	0.0079		0.8836	
Alcohol	0.1985		0.7376	
LIFETIME				
Illicit Drugs ²	0.0000		0.4576	
Marijuana	0.0000		0.8130	
Illicit Drugs Other Than Marijuana ²	0.0002		0.3325	
Cigarettes	0.0000		0.4538	
Alcohol	0.0194		0.6780	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.3C Risk Perceptions of Smoking Marijuana Once a Month, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking Marijuana Once a Month (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Smoke Marijuana Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	109	107	109	107
AGE				
12-13	87	83	96	98
14-15	79	77	119	117
16-17	68	68	114	123
GENDER				
Male	73	73	73	73
Female	73	74	73	74
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	92	92	92	92
White	106	108	165	176
Black or African American	57	58	90	97
American Indian or Alaska Native	9	7	11	15
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	36	42	47	53
Two or More Races	14	11	25	21
Hispanic or Latino	56	57	56	57
COUNTY TYPE				
Large Metro	113	116	183	187
Small Metro	86	80	138	120
Nonmetro	62	57	87	88
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	26	25	71	66
Marijuana	16	15	60	57
Illicit Drugs Other Than Marijuana ²	22	20	49	43
Cigarettes	29	28	77	68
Alcohol	41	38	89	83
Binge Alcohol ³	30	28	73	62
PAST YEAR				
Illicit Drugs ²	41	39	97	97
Marijuana	27	23	83	81
Illicit Drugs Other Than Marijuana ²	34	33	74	72
Cigarettes	41	39	91	91
Alcohol	65	59	125	128
LIFETIME				
Illicit Drugs ²	54	52	116	112
Marijuana	33	30	95	91
Illicit Drugs Other Than Marijuana ²	48	47	95	90
Cigarettes	56	53	116	117
Alcohol	74	71	144	141

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.3D Risk Perceptions of Smoking Marijuana Once a Month, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking Marijuana Once a Month (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Smoke Marijuana Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.44	0.43	0.44	0.43
AGE				
12-13	0.74	0.72	0.74	0.72
14-15	0.72	0.71	0.72	0.71
16-17	0.69	0.71	0.69	0.71
GENDER				
Male	0.58	0.57	0.58	0.57
Female	0.60	0.60	0.60	0.60
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.44	0.44	0.44	0.44
White	0.50	0.51	0.50	0.51
Black or African American	1.06	1.05	1.06	1.05
American Indian or Alaska Native	4.50	3.67	4.50	3.67
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.73	2.52	2.73	2.52
Two or More Races	2.72	2.39	2.72	2.39
Hispanic or Latino	1.34	1.32	1.34	1.32
COUNTY TYPE				
Large Metro	0.64	0.61	0.64	0.61
Small Metro	0.74	0.75	0.74	0.75
Nonmetro	0.89	0.87	0.89	0.87
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.29	0.29	0.39	0.35
Marijuana	0.18	0.18	0.33	0.32
Illicit Drugs Other Than Marijuana ²	0.24	0.23	0.28	0.24
Cigarettes	0.33	0.32	0.42	0.38
Alcohol	0.44	0.42	0.45	0.43
Binge Alcohol ³	0.33	0.33	0.41	0.35
PAST YEAR				
Illicit Drugs ²	0.44	0.43	0.47	0.46
Marijuana	0.31	0.26	0.42	0.41
Illicit Drugs Other Than Marijuana ²	0.37	0.37	0.40	0.38
Cigarettes	0.45	0.44	0.46	0.47
Alcohol	0.65	0.59	0.51	0.52
LIFETIME				
Illicit Drugs ²	0.54	0.54	0.50	0.49
Marijuana	0.37	0.34	0.47	0.45
Illicit Drugs Other Than Marijuana ²	0.49	0.50	0.47	0.45
Cigarettes	0.58	0.58	0.53	0.55
Alcohol	0.69	0.71	0.55	0.53

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.3P Risk Perceptions of Smoking Marijuana Once a Month, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking Marijuana Once a Month (II), among Persons Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

(I) Demographic Characteristic and (II) Substance Use	Smoke Marijuana Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.1024		0.1024	
AGE				
12-13	0.0749		0.0749	
14-15	0.2105		0.2105	
16-17	0.6214		0.6214	
GENDER				
Male	0.0662		0.0662	
Female	0.5941		0.5941	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.1382		0.1382	
White	0.2933		0.2933	
Black or African American	0.3863		0.3863	
American Indian or Alaska Native	0.1376		0.1376	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.9390		0.9390	
Two or More Races	0.1039		0.1039	
Hispanic or Latino	0.4532		0.4532	
COUNTY TYPE				
Large Metro	0.0207		0.0207	
Small Metro	0.0834		0.0834	
Nonmetro	0.0360		0.0360	
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.8128		0.0150	
Marijuana	0.9546		0.0056	
Illicit Drugs Other Than Marijuana ²	0.9023		0.1079	
Cigarettes	0.0699		0.0204	
Alcohol	0.9843		0.0043	
Binge Alcohol ³	0.6134		0.0011	
PAST YEAR				
Illicit Drugs ²	0.7440		0.0033	
Marijuana	0.1951		0.0037	
Illicit Drugs Other Than Marijuana ²	0.6352		0.0220	
Cigarettes	0.1849		0.0395	
Alcohol	0.9332		0.2019	
LIFETIME				
Illicit Drugs ²	0.0720		0.0000	
Marijuana	0.3694		0.0001	
Illicit Drugs Other Than Marijuana ²	0.0679		0.0006	
Cigarettes	0.0187		0.0001	
Alcohol	0.4657		0.0105	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.4C Risk Perceptions of Smoking Marijuana Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking Marijuana Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Smoke Marijuana Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	112	117	112	117
AGE				
12-13	109	106	78	79
14-15	100	105	99	94
16-17	89	89	97	104
GENDER				
Male	79	81	79	81
Female	73	81	73	81
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	99	100	99	100
White	145	157	126	126
Black or African American	74	66	80	88
American Indian or Alaska Native	12	11	10	12
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	43	63	40	39
Two or More Races	20	16	21	17
Hispanic or Latino	58	58	58	58
COUNTY TYPE				
Large Metro	149	168	141	146
Small Metro	118	109	110	90
Nonmetro	86	81	71	67
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	31	30	66	65
Marijuana	19	18	59	57
Illicit Drugs Other Than Marijuana ²	25	24	46	41
Cigarettes	37	31	74	66
Alcohol	52	47	83	77
Binge Alcohol ³	37	34	71	61
PAST YEAR				
Illicit Drugs ²	53	51	92	93
Marijuana	35	30	81	79
Illicit Drugs Other Than Marijuana ²	43	41	71	69
Cigarettes	50	45	88	86
Alcohol	83	77	112	114
LIFETIME				
Illicit Drugs ²	67	65	109	107
Marijuana	46	38	89	89
Illicit Drugs Other Than Marijuana ²	59	58	91	84
Cigarettes	70	67	105	105
Alcohol	97	88	122	124

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.4D Risk Perceptions of Smoking Marijuana Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking Marijuana Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Smoke Marijuana Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.45	0.47	0.45	0.47
AGE				
12-13	0.75	0.74	0.75	0.74
14-15	0.75	0.78	0.75	0.78
16-17	0.77	0.76	0.77	0.76
GENDER				
Male	0.63	0.64	0.63	0.64
Female	0.59	0.66	0.59	0.66
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.48	0.49	0.48	0.49
White	0.52	0.53	0.52	0.53
Black or African American	1.26	1.08	1.26	1.08
American Indian or Alaska Native	5.17	4.28	5.17	4.28
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.91	3.03	2.91	3.03
Two or More Races	3.07	2.90	3.07	2.90
Hispanic or Latino	1.40	1.34	1.40	1.34
COUNTY TYPE				
Large Metro	0.64	0.70	0.64	0.70
Small Metro	0.81	0.75	0.81	0.75
Nonmetro	1.06	0.92	1.06	0.92
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.23	0.21	0.48	0.49
Marijuana	0.14	0.13	0.45	0.45
Illicit Drugs Other Than Marijuana ²	0.18	0.17	0.37	0.33
Cigarettes	0.26	0.22	0.57	0.53
Alcohol	0.35	0.33	0.58	0.56
Binge Alcohol ³	0.26	0.24	0.54	0.49
PAST YEAR				
Illicit Drugs ²	0.37	0.34	0.59	0.63
Marijuana	0.25	0.21	0.56	0.55
Illicit Drugs Other Than Marijuana ²	0.30	0.28	0.53	0.53
Cigarettes	0.35	0.32	0.63	0.64
Alcohol	0.52	0.48	0.65	0.66
LIFETIME				
Illicit Drugs ²	0.43	0.43	0.61	0.64
Marijuana	0.33	0.27	0.58	0.59
Illicit Drugs Other Than Marijuana ²	0.39	0.38	0.61	0.60
Cigarettes	0.46	0.44	0.66	0.68
Alcohol	0.57	0.54	0.63	0.64

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.4P Risk Perceptions of Smoking Marijuana Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Smoking Marijuana Once or Twice a Week (II), among Persons Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

(I) Demographic Characteristic and (II) Substance Use	Smoke Marijuana Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.6472		0.6472	
AGE				
12-13	0.6948		0.6948	
14-15	0.3135		0.3135	
16-17	0.5360		0.5360	
GENDER				
Male	0.3909		0.3909	
Female	0.8512		0.8512	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.9428		0.9428	
White	0.2331		0.2331	
Black or African American	0.0221		0.0221	
American Indian or Alaska Native	0.4049		0.4049	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.5725		0.5725	
Two or More Races	0.9804		0.9804	
Hispanic or Latino	0.4047		0.4047	
COUNTY TYPE				
Large Metro	0.4457		0.4457	
Small Metro	0.0220		0.0220	
Nonmetro	0.7914		0.7914	
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.4320		0.0903	
Marijuana	0.6270		0.0191	
Illicit Drugs Other Than Marijuana ²	0.2886		0.4460	
Cigarettes	0.0001		0.3705	
Alcohol	0.3257		0.0446	
Binge Alcohol ³	0.2303		0.0095	
PAST YEAR				
Illicit Drugs ²	0.4042		0.0259	
Marijuana	0.0730		0.0282	
Illicit Drugs Other Than Marijuana ²	0.8314		0.0943	
Cigarettes	0.0047		0.4893	
Alcohol	0.1956		0.6744	
LIFETIME				
Illicit Drugs ²	0.0023		0.0011	
Marijuana	0.0014		0.0235	
Illicit Drugs Other Than Marijuana ²	0.0412		0.0032	
Cigarettes	0.0001		0.0240	
Alcohol	0.0412		0.4292	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.5C Risk Perceptions of Using Cocaine Once a Month, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Cocaine Once a Month (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Use Cocaine Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	106	114	106	114
AGE				
12-13	85	84	102	99
14-15	102	96	104	103
16-17	101	102	82	89
GENDER				
Male	72	79	72	79
Female	74	78	74	78
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	93	100	93	100
White	136	131	135	148
Black or African American	76	84	79	75
American Indian or Alaska Native	9	10	10	14
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	38	41	42	60
Two or More Races	19	17	19	16
Hispanic or Latino	47	55	47	55
COUNTY TYPE				
Large Metro	130	150	154	160
Small Metro	103	99	122	99
Nonmetro	79	72	72	74
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	49	45	53	51
Marijuana	42	38	44	42
Illicit Drugs Other Than Marijuana ²	33	29	40	36
Cigarettes	55	50	53	50
Alcohol	67	61	67	66
Binge Alcohol ³	55	48	54	48
PAST YEAR				
Illicit Drugs ²	72	70	70	75
Marijuana	61	58	58	56
Illicit Drugs Other Than Marijuana ²	52	48	57	61
Cigarettes	71	66	67	67
Alcohol	101	96	93	97
LIFETIME				
Illicit Drugs ²	86	84	89	91
Marijuana	75	68	63	64
Illicit Drugs Other Than Marijuana ²	66	61	78	80
Cigarettes	90	83	85	86
Alcohol	113	106	103	107

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.5D Risk Perceptions of Using Cocaine Once a Month, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Cocaine Once a Month (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Use Cocaine Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.43	0.46	0.43	0.46
AGE				
12-13	0.76	0.74	0.76	0.74
14-15	0.82	0.78	0.82	0.78
16-17	0.74	0.76	0.74	0.76
GENDER				
Male	0.57	0.63	0.57	0.63
Female	0.61	0.64	0.61	0.64
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.45	0.49	0.45	0.49
White	0.52	0.51	0.52	0.51
Black or African American	1.29	1.19	1.29	1.19
American Indian or Alaska Native	4.32	4.15	4.32	4.15
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.52	3.00	2.52	3.00
Two or More Races	2.74	2.96	2.74	2.96
Hispanic or Latino	1.13	1.28	1.13	1.28
COUNTY TYPE				
Large Metro	0.60	0.68	0.60	0.68
Small Metro	0.78	0.75	0.78	0.75
Nonmetro	0.95	0.93	0.95	0.93
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.37	0.35	0.38	0.37
Marijuana	0.32	0.30	0.32	0.31
Illicit Drugs Other Than Marijuana ²	0.26	0.24	0.30	0.27
Cigarettes	0.42	0.39	0.38	0.37
Alcohol	0.48	0.45	0.47	0.46
Binge Alcohol ³	0.42	0.37	0.40	0.36
PAST YEAR				
Illicit Drugs ²	0.52	0.49	0.46	0.51
Marijuana	0.46	0.43	0.40	0.40
Illicit Drugs Other Than Marijuana ²	0.39	0.37	0.41	0.43
Cigarettes	0.51	0.50	0.48	0.47
Alcohol	0.64	0.59	0.58	0.61
LIFETIME				
Illicit Drugs ²	0.58	0.56	0.53	0.56
Marijuana	0.55	0.49	0.44	0.45
Illicit Drugs Other Than Marijuana ²	0.47	0.45	0.49	0.52
Cigarettes	0.60	0.59	0.56	0.55
Alcohol	0.65	0.61	0.60	0.63

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.5P Risk Perceptions of Using Cocaine Once a Month, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Cocaine Once a Month (II), among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2005 Versus 2004

(I) Demographic Characteristic and (II) Substance Use	Use Cocaine Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.2035		0.2035	
AGE				
12-13	0.5149		0.5149	
14-15	0.3281		0.3281	
16-17	0.3966		0.3966	
GENDER				
Male	0.3235		0.3235	
Female	0.3890		0.3890	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.0406		0.0406	
White	0.0156		0.0156	
Black or African American	0.8889		0.8889	
American Indian or Alaska Native	0.1634		0.1634	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.9537		0.9537	
Two or More Races	0.4086		0.4086	
Hispanic or Latino	0.2291		0.2291	
COUNTY TYPE				
Large Metro	0.0611		0.0611	
Small Metro	0.0329		0.0329	
Nonmetro	0.0088		0.0088	
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.0401		0.3135	
Marijuana	0.0198		0.1524	
Illicit Drugs Other Than Marijuana ²	0.3934		0.2994	
Cigarettes	0.0388		0.0285	
Alcohol	0.0053		0.6465	
Binge Alcohol ³	0.0019		0.2665	
PAST YEAR				
Illicit Drugs ²	0.0153		0.2858	
Marijuana	0.0186		0.0999	
Illicit Drugs Other Than Marijuana ²	0.2313		0.1809	
Cigarettes	0.1049		0.0784	
Alcohol	0.3746		0.8221	
LIFETIME				
Illicit Drugs ²	0.0000		0.0569	
Marijuana	0.0010		0.1683	
Illicit Drugs Other Than Marijuana ²	0.0012		0.0246	
Cigarettes	0.0006		0.0064	
Alcohol	0.1106		0.1936	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.6C Risk Perceptions of Using Cocaine Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Cocaine Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Use Cocaine Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	92	95	92	95
AGE				
12-13	117	116	66	67
14-15	127	133	64	60
16-17	124	129	52	52
GENDER				
Male	63	64	63	64
Female	62	65	62	65
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	76	79	76	79
White	193	206	68	69
Black or African American	94	104	53	50
American Indian or Alaska Native	14	13	7	10
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	55	60	30	33
Two or More Races	27	20	10	12
Hispanic or Latino	47	47	47	47
COUNTY TYPE				
Large Metro	198	215	83	91
Small Metro	152	143	76	52
Nonmetro	103	107	43	41
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	64	60	39	36
Marijuana	53	51	31	28
Illicit Drugs Other Than Marijuana ²	43	38	31	28
Cigarettes	72	64	37	33
Alcohol	85	83	42	39
Binge Alcohol ³	68	60	35	33
PAST YEAR				
Illicit Drugs ²	93	91	47	51
Marijuana	77	76	38	37
Illicit Drugs Other Than Marijuana ²	66	64	41	42
Cigarettes	89	85	46	44
Alcohol	125	128	57	58
LIFETIME				
Illicit Drugs ²	113	109	58	65
Marijuana	89	88	42	42
Illicit Drugs Other Than Marijuana ²	89	82	51	58
Cigarettes	112	110	53	57
Alcohol	143	144	63	64

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.6D Risk Perceptions of Using Cocaine Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Cocaine Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Use Cocaine Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.37	0.38	0.37	0.38
AGE				
12-13	0.68	0.69	0.68	0.69
14-15	0.62	0.63	0.62	0.63
16-17	0.58	0.56	0.58	0.56
GENDER				
Male	0.50	0.51	0.50	0.51
Female	0.51	0.53	0.51	0.53
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.37	0.38	0.37	0.38
White	0.38	0.39	0.38	0.39
Black or African American	1.04	0.98	1.04	0.98
American Indian or Alaska Native	4.10	3.84	4.10	3.84
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.67	2.34	2.67	2.34
Two or More Races	2.11	2.66	2.11	2.66
Hispanic or Latino	1.13	1.09	1.13	1.09
COUNTY TYPE				
Large Metro	0.51	0.57	0.51	0.57
Small Metro	0.77	0.59	0.77	0.59
Nonmetro	0.76	0.80	0.76	0.80
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.30	0.28	0.72	0.66
Marijuana	0.25	0.24	0.59	0.52
Illicit Drugs Other Than Marijuana ²	0.21	0.18	0.58	0.53
Cigarettes	0.33	0.31	0.72	0.61
Alcohol	0.37	0.37	0.76	0.71
Binge Alcohol ³	0.32	0.28	0.65	0.62
PAST YEAR				
Illicit Drugs ²	0.40	0.39	0.83	0.84
Marijuana	0.35	0.34	0.69	0.66
Illicit Drugs Other Than Marijuana ²	0.30	0.30	0.74	0.73
Cigarettes	0.40	0.39	0.87	0.75
Alcohol	0.47	0.48	0.93	0.93
LIFETIME				
Illicit Drugs ²	0.45	0.44	0.93	0.95
Marijuana	0.40	0.38	0.76	0.73
Illicit Drugs Other Than Marijuana ²	0.37	0.36	0.87	0.91
Cigarettes	0.46	0.47	0.95	0.91
Alcohol	0.50	0.50	0.97	0.96

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.6P Risk Perceptions of Using Cocaine Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Cocaine Once or Twice a Week (II), among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2005 Versus 2004

(I) Demographic Characteristic and (II) Substance Use	Use Cocaine Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.8690		0.8690	
AGE				
12-13	0.7016		0.7016	
14-15	0.9414		0.9414	
16-17	0.5679		0.5679	
GENDER				
Male	0.5107		0.5107	
Female	0.6770		0.6770	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.5657		0.5657	
White	0.7241		0.7241	
Black or African American	0.7903		0.7903	
American Indian or Alaska Native	0.0776		0.0776	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.6660		0.6660	
Two or More Races	0.3617		0.3617	
Hispanic or Latino	0.1599		0.1599	
COUNTY TYPE				
Large Metro	0.2655		0.2655	
Small Metro	0.0959		0.0959	
Nonmetro	0.7555		0.7555	
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.0655		0.4029	
Marijuana	0.0500		0.0792	
Illicit Drugs Other Than Marijuana ²	0.2762		0.5426	
Cigarettes	0.0187		0.1074	
Alcohol	0.0657		0.2025	
Binge Alcohol ³	0.0068		0.2362	
PAST YEAR				
Illicit Drugs ²	0.0140		0.7355	
Marijuana	0.0196		0.1423	
Illicit Drugs Other Than Marijuana ²	0.1400		0.4765	
Cigarettes	0.0546		0.2360	
Alcohol	0.1429		0.2322	
LIFETIME				
Illicit Drugs ²	0.0000		0.9226	
Marijuana	0.0020		0.2114	
Illicit Drugs Other Than Marijuana ²	0.0000		0.8209	
Cigarettes	0.0001		0.0930	
Alcohol	0.0141		0.7132	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.7C Risk Perceptions of Trying Heroin Once or Twice, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Trying Heroin Once or Twice (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Try Heroin Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	103	110	103	110
AGE				
12-13	86	88	96	95
14-15	107	108	95	92
16-17	113	113	70	73
GENDER				
Male	74	76	74	76
Female	73	73	73	73
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	91	97	91	97
White	140	150	129	131
Black or African American	80	89	69	66
American Indian or Alaska Native	14	13	8	10
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	40	43	42	57
Two or More Races	21	19	18	15
Hispanic or Latino	47	52	47	52
COUNTY TYPE				
Large Metro	142	162	138	144
Small Metro	119	111	101	85
Nonmetro	81	82	64	66
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	59	57	41	39
Marijuana	50	49	34	31
Illicit Drugs Other Than Marijuana ²	40	36	32	29
Cigarettes	65	60	42	38
Alcohol	75	75	53	53
Binge Alcohol ³	62	58	42	37
PAST YEAR				
Illicit Drugs ²	84	84	55	59
Marijuana	70	70	46	41
Illicit Drugs Other Than Marijuana ²	63	60	47	50
Cigarettes	82	79	51	52
Alcohol	110	110	80	79
LIFETIME				
Illicit Drugs ²	99	98	72	73
Marijuana	83	82	50	48
Illicit Drugs Other Than Marijuana ²	81	74	65	66
Cigarettes	101	98	67	71
Alcohol	123	123	89	90

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.7D Risk Perceptions of Trying Heroin Once or Twice, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Trying Heroin Once or Twice (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Try Heroin Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.42	0.44	0.42	0.44
AGE				
12-13	0.75	0.76	0.75	0.76
14-15	0.79	0.79	0.79	0.79
16-17	0.70	0.68	0.70	0.68
GENDER				
Male	0.59	0.60	0.59	0.60
Female	0.60	0.60	0.60	0.60
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.44	0.47	0.44	0.47
White	0.52	0.53	0.52	0.53
Black or African American	1.16	1.11	1.16	1.11
American Indian or Alaska Native	5.25	4.30	5.25	4.30
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.83	3.09	2.83	3.09
Two or More Races	2.68	3.01	2.68	3.01
Hispanic or Latino	1.14	1.22	1.14	1.22
COUNTY TYPE				
Large Metro	0.60	0.67	0.60	0.67
Small Metro	0.74	0.72	0.74	0.72
Nonmetro	0.86	0.96	0.86	0.96
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.39	0.37	0.36	0.35
Marijuana	0.34	0.33	0.30	0.28
Illicit Drugs Other Than Marijuana ²	0.27	0.25	0.29	0.26
Cigarettes	0.41	0.40	0.37	0.34
Alcohol	0.46	0.47	0.45	0.46
Binge Alcohol ³	0.41	0.38	0.37	0.33
PAST YEAR				
Illicit Drugs ²	0.51	0.50	0.46	0.49
Marijuana	0.45	0.44	0.40	0.36
Illicit Drugs Other Than Marijuana ²	0.40	0.39	0.42	0.43
Cigarettes	0.49	0.49	0.44	0.45
Alcohol	0.57	0.58	0.61	0.60
LIFETIME				
Illicit Drugs ²	0.54	0.55	0.54	0.55
Marijuana	0.51	0.50	0.42	0.42
Illicit Drugs Other Than Marijuana ²	0.48	0.46	0.52	0.52
Cigarettes	0.57	0.57	0.53	0.57
Alcohol	0.59	0.61	0.63	0.65

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.7P Risk Perceptions of Trying Heroin Once or Twice, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Trying Heroin Once or Twice (II), among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2005 Versus 2004

(I) Demographic Characteristic and (II) Substance Use	Try Heroin Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.3767		0.3767	
AGE				
12-13	0.2489		0.2489	
14-15	0.6300		0.6300	
16-17	0.7349		0.7349	
GENDER				
Male	0.9201		0.9201	
Female	0.2339		0.2339	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.1844		0.1844	
White	0.0494		0.0494	
Black or African American	0.6942		0.6942	
American Indian or Alaska Native	0.6455		0.6455	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.6802		0.6802	
Two or More Races	0.5665		0.5665	
Hispanic or Latino	0.5509		0.5509	
COUNTY TYPE				
Large Metro	0.1193		0.1193	
Small Metro	0.0771		0.0771	
Nonmetro	0.1092		0.1092	
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.1438		0.1820	
Marijuana	0.0655		0.0868	
Illicit Drugs Other Than Marijuana ²	0.5418		0.2332	
Cigarettes	0.0540		0.0202	
Alcohol	0.0398		0.3487	
Binge Alcohol ³	0.0196		0.0881	
PAST YEAR				
Illicit Drugs ²	0.0683		0.1273	
Marijuana	0.0705		0.0154	
Illicit Drugs Other Than Marijuana ²	0.2462		0.2763	
Cigarettes	0.1752		0.0286	
Alcohol	0.4830		0.6200	
LIFETIME				
Illicit Drugs ²	0.0010		0.0072	
Marijuana	0.0206		0.0092	
Illicit Drugs Other Than Marijuana ²	0.0022		0.0450	
Cigarettes	0.0034		0.0008	
Alcohol	0.1234		0.1417	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.8C Risk Perceptions of Using Heroin Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Heroin Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Use Heroin Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	86	86	86	86
AGE				
12-13	114	114	67	68
14-15	130	133	61	57
16-17	131	136	42	42
GENDER				
Male	64	62	64	62
Female	55	62	55	62
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	75	73	75	73
White	192	206	68	67
Black or African American	94	104	53	47
American Indian or Alaska Native	14	15	7	7
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	52	67	28	29
Two or More Races	27	21	10	10
Hispanic or Latino	43	44	43	44
COUNTY TYPE				
Large Metro	199	215	80	83
Small Metro	151	143	68	54
Nonmetro	102	110	42	33
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	69	62	30	30
Marijuana	57	53	23	21
Illicit Drugs Other Than Marijuana ²	47	38	24	25
Cigarettes	75	68	28	25
Alcohol	87	86	35	31
Binge Alcohol ³	72	64	28	24
PAST YEAR				
Illicit Drugs ²	97	94	40	42
Marijuana	81	79	28	26
Illicit Drugs Other Than Marijuana ²	73	68	34	37
Cigarettes	94	90	36	35
Alcohol	131	131	49	45
LIFETIME				
Illicit Drugs ²	116	115	48	51
Marijuana	92	91	31	31
Illicit Drugs Other Than Marijuana ²	95	89	43	46
Cigarettes	117	115	46	45
Alcohol	149	145	52	54

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.8D Risk Perceptions of Using Heroin Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Heroin Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Use Heroin Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.35	0.35	0.35	0.35
AGE				
12-13	0.69	0.71	0.69	0.71
14-15	0.62	0.59	0.62	0.59
16-17	0.49	0.47	0.49	0.47
GENDER				
Male	0.51	0.49	0.51	0.49
Female	0.45	0.51	0.45	0.51
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.37	0.35	0.37	0.35
White	0.39	0.38	0.39	0.38
Black or African American	1.05	0.91	1.05	0.91
American Indian or Alaska Native	4.20	3.27	4.20	3.27
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.40	2.39	2.40	2.39
Two or More Races	2.15	2.41	2.15	2.41
Hispanic or Latino	1.05	1.03	1.05	1.03
COUNTY TYPE				
Large Metro	0.50	0.52	0.50	0.52
Small Metro	0.68	0.62	0.68	0.62
Nonmetro	0.75	0.69	0.75	0.69
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.31	0.28	0.60	0.62
Marijuana	0.26	0.25	0.47	0.45
Illicit Drugs Other Than Marijuana ²	0.22	0.18	0.51	0.52
Cigarettes	0.33	0.32	0.60	0.53
Alcohol	0.37	0.37	0.69	0.66
Binge Alcohol ³	0.33	0.30	0.58	0.52
PAST YEAR				
Illicit Drugs ²	0.41	0.39	0.77	0.80
Marijuana	0.36	0.34	0.57	0.55
Illicit Drugs Other Than Marijuana ²	0.32	0.31	0.69	0.74
Cigarettes	0.40	0.40	0.74	0.73
Alcohol	0.46	0.48	0.89	0.90
LIFETIME				
Illicit Drugs ²	0.44	0.45	0.85	0.91
Marijuana	0.39	0.39	0.62	0.64
Illicit Drugs Other Than Marijuana ²	0.38	0.38	0.82	0.87
Cigarettes	0.46	0.47	0.90	0.86
Alcohol	0.49	0.50	0.91	1.00

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.8P Risk Perceptions of Using Heroin Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using Heroin Once or Twice a Week (II), among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2005 Versus 2004

(I) Demographic Characteristic and (II) Substance Use	Use Heroin Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.5077		0.5077	
AGE				
12-13	0.7057		0.7057	
14-15	0.3574		0.3574	
16-17	0.8314		0.8314	
GENDER				
Male	0.2856		0.2856	
Female	0.8591		0.8591	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.5931		0.5931	
White	0.9909		0.9909	
Black or African American	0.1829		0.1829	
American Indian or Alaska Native	0.8867		0.8867	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.8147		0.8147	
Two or More Races	0.5698		0.5698	
Hispanic or Latino	0.6173		0.6173	
COUNTY TYPE				
Large Metro	0.6611		0.6611	
Small Metro	0.2076		0.2076	
Nonmetro	0.4165		0.4165	
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.0235		0.9244	
Marijuana	0.0087		0.7539	
Illicit Drugs Other Than Marijuana ²	0.1289		0.9098	
Cigarettes	0.0041		0.4972	
Alcohol	0.0209		0.6144	
Binge Alcohol ³	0.0037		0.5127	
PAST YEAR				
Illicit Drugs ²	0.0075		0.9571	
Marijuana	0.0036		0.7205	
Illicit Drugs Other Than Marijuana ²	0.0637		1.0000	
Cigarettes	0.0242		0.5134	
Alcohol	0.2503		0.7594	
LIFETIME				
Illicit Drugs ²	0.0000		0.4800	
Marijuana	0.0005		0.5815	
Illicit Drugs Other Than Marijuana ²	0.0001		0.6476	
Cigarettes	0.0000		0.3040	
Alcohol	0.0144		0.8925	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.9C Risk Perceptions of Trying LSD Once or Twice, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Trying LSD Once or Twice (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Try LSD Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	105	115	105	115
AGE				
12-13	85	85	95	93
14-15	98	103	92	94
16-17	99	98	81	85
GENDER				
Male	74	80	74	80
Female	76	77	76	77
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	89	101	89	101
White	132	138	134	138
Black or African American	78	83	67	73
American Indian or Alaska Native	11	10	8	14
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	41	48	38	45
Two or More Races	19	18	18	14
Hispanic or Latino	52	51	52	51
COUNTY TYPE				
Large Metro	133	156	146	143
Small Metro	112	105	109	91
Nonmetro	79	74	68	70
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	50	49	49	48
Marijuana	43	40	41	41
Illicit Drugs Other Than Marijuana ²	33	32	38	35
Cigarettes	57	53	51	50
Alcohol	69	67	62	62
Binge Alcohol ³	55	52	53	45
PAST YEAR				
Illicit Drugs ²	74	76	66	70
Marijuana	64	62	54	54
Illicit Drugs Other Than Marijuana ²	54	53	56	56
Cigarettes	72	71	62	63
Alcohol	101	101	89	90
LIFETIME				
Illicit Drugs ²	91	88	80	85
Marijuana	77	70	61	61
Illicit Drugs Other Than Marijuana ²	71	67	71	73
Cigarettes	94	88	78	78
Alcohol	114	114	96	101

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.9D Risk Perceptions of Trying LSD Once or Twice, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Trying LSD Once or Twice (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Try LSD Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.44	0.48	0.44	0.48
AGE				
12-13	0.77	0.77	0.77	0.77
14-15	0.73	0.80	0.73	0.80
16-17	0.73	0.73	0.73	0.73
GENDER				
Male	0.60	0.65	0.60	0.65
Female	0.64	0.65	0.64	0.65
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.44	0.51	0.44	0.51
White	0.52	0.54	0.52	0.54
Black or African American	1.13	1.26	1.13	1.26
American Indian or Alaska Native	3.83	4.77	3.83	4.77
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.73	3.14	2.73	3.14
Two or More Races	2.67	3.01	2.67	3.01
Hispanic or Latino	1.30	1.22	1.30	1.22
COUNTY TYPE				
Large Metro	0.63	0.69	0.63	0.69
Small Metro	0.79	0.79	0.79	0.79
Nonmetro	0.93	0.98	0.93	0.98
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.37	0.37	0.40	0.39
Marijuana	0.32	0.31	0.34	0.33
Illicit Drugs Other Than Marijuana ²	0.25	0.25	0.32	0.29
Cigarettes	0.41	0.40	0.41	0.40
Alcohol	0.48	0.47	0.48	0.48
Binge Alcohol ³	0.40	0.39	0.43	0.36
PAST YEAR				
Illicit Drugs ²	0.50	0.50	0.49	0.52
Marijuana	0.46	0.43	0.42	0.43
Illicit Drugs Other Than Marijuana ²	0.39	0.39	0.44	0.43
Cigarettes	0.49	0.50	0.49	0.48
Alcohol	0.60	0.61	0.61	0.59
LIFETIME				
Illicit Drugs ²	0.58	0.56	0.55	0.58
Marijuana	0.55	0.48	0.47	0.47
Illicit Drugs Other Than Marijuana ²	0.48	0.47	0.53	0.53
Cigarettes	0.60	0.57	0.57	0.56
Alcohol	0.62	0.65	0.61	0.61

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.9P Risk Perceptions of Trying LSD Once or Twice, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Trying LSD Once or Twice (II), among Persons Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

(I) Demographic Characteristic and (II) Substance Use	Try LSD Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.1433		0.1433	
AGE				
12-13	0.2440		0.2440	
14-15	0.4987		0.4987	
16-17	0.2461		0.2461	
GENDER				
Male	0.4371		0.4371	
Female	0.1845		0.1845	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.0864		0.0864	
White	0.1449		0.1449	
Black or African American	0.3643		0.3643	
American Indian or Alaska Native	0.3355		0.3355	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.9851		0.9851	
Two or More Races	0.4342		0.4342	
Hispanic or Latino	0.9923		0.9923	
COUNTY TYPE				
Large Metro	0.0263		0.0263	
Small Metro	0.1081		0.1081	
Nonmetro	0.0955		0.0955	
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.0321		0.2997	
Marijuana	0.0082		0.1540	
Illicit Drugs Other Than Marijuana ²	0.2478		0.4785	
Cigarettes	0.0299		0.0906	
Alcohol	0.0435		0.3534	
Binge Alcohol ³	0.0378		0.0629	
PAST YEAR				
Illicit Drugs ²	0.0543		0.1475	
Marijuana	0.0548		0.0253	
Illicit Drugs Other Than Marijuana ²	0.1067		0.4745	
Cigarettes	0.1095		0.0850	
Alcohol	0.3184		0.9763	
LIFETIME				
Illicit Drugs ²	0.0001		0.0113	
Marijuana	0.0024		0.0644	
Illicit Drugs Other Than Marijuana ²	0.0017		0.0178	
Cigarettes	0.0014		0.0013	
Alcohol	0.0412		0.5359	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.10C Risk Perceptions of Using LSD Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using LSD Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Use LSD Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	88	95	88	95
AGE				
12-13	104	108	71	68
14-15	124	129	61	62
16-17	123	123	51	56
GENDER				
Male	67	66	67	66
Female	59	67	59	67
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	78	82	78	82
White	184	194	76	77
Black or African American	90	91	53	62
American Indian or Alaska Native	12	14	9	10
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	54	59	30	27
Two or More Races	26	21	13	10
Hispanic or Latino	43	46	43	46
COUNTY TYPE				
Large Metro	180	203	94	90
Small Metro	149	133	69	60
Nonmetro	99	96	44	42
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	64	58	39	38
Marijuana	55	48	31	32
Illicit Drugs Other Than Marijuana ²	41	37	29	28
Cigarettes	68	62	38	36
Alcohol	84	80	43	42
Binge Alcohol ³	67	57	36	34
PAST YEAR				
Illicit Drugs ²	91	89	50	49
Marijuana	78	74	38	38
Illicit Drugs Other Than Marijuana ²	65	65	41	40
Cigarettes	88	83	45	45
Alcohol	124	124	60	59
LIFETIME				
Illicit Drugs ²	107	104	61	60
Marijuana	88	85	43	43
Illicit Drugs Other Than Marijuana ²	84	81	52	52
Cigarettes	110	107	57	55
Alcohol	142	139	66	66

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.10D Risk Perceptions of Using LSD Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using LSD Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Use LSD Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.36	0.39	0.36	0.39
AGE				
12-13	0.70	0.72	0.70	0.72
14-15	0.61	0.67	0.61	0.67
16-17	0.57	0.61	0.57	0.61
GENDER				
Male	0.55	0.54	0.55	0.54
Female	0.49	0.57	0.49	0.57
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.39	0.41	0.39	0.41
White	0.42	0.43	0.42	0.43
Black or African American	1.03	1.17	1.03	1.17
American Indian or Alaska Native	4.30	4.83	4.30	4.83
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.86	2.26	2.86	2.26
Two or More Races	2.71	2.51	2.71	2.51
Hispanic or Latino	1.07	1.10	1.07	1.10
COUNTY TYPE				
Large Metro	0.55	0.58	0.55	0.58
Small Metro	0.68	0.66	0.68	0.66
Nonmetro	0.74	0.75	0.74	0.75
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.32	0.29	0.61	0.62
Marijuana	0.28	0.25	0.51	0.52
Illicit Drugs Other Than Marijuana ²	0.22	0.19	0.48	0.47
Cigarettes	0.33	0.32	0.62	0.58
Alcohol	0.39	0.38	0.70	0.67
Binge Alcohol ³	0.34	0.29	0.60	0.56
PAST YEAR				
Illicit Drugs ²	0.43	0.41	0.74	0.73
Marijuana	0.38	0.36	0.59	0.61
Illicit Drugs Other Than Marijuana ²	0.32	0.32	0.63	0.63
Cigarettes	0.41	0.41	0.71	0.70
Alcohol	0.49	0.49	0.87	0.86
LIFETIME				
Illicit Drugs ²	0.46	0.46	0.83	0.83
Marijuana	0.42	0.40	0.65	0.68
Illicit Drugs Other Than Marijuana ²	0.39	0.39	0.76	0.78
Cigarettes	0.48	0.49	0.84	0.82
Alcohol	0.52	0.52	0.89	0.90

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.10P Risk Perceptions of Using LSD Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Using LSD Once or Twice a Week (II), among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2005 Versus 2004

(I) Demographic Characteristic and (II) Substance Use	Use LSD Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.5337		0.5337	
AGE				
12-13	0.4868		0.4868	
14-15	0.8206		0.8206	
16-17	0.5592		0.5592	
GENDER				
Male	0.6759		0.6759	
Female	0.1791		0.1791	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.6344		0.6344	
White	0.9277		0.9277	
Black or African American	0.1311		0.1311	
American Indian or Alaska Native	0.9316		0.9316	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.2701		0.2701	
Two or More Races	0.9234		0.9234	
Hispanic or Latino	0.7658		0.7658	
COUNTY TYPE				
Large Metro	0.5453		0.5453	
Small Metro	0.8223		0.8223	
Nonmetro	0.4353		0.4353	
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.0106		0.8864	
Marijuana	0.0013		0.9313	
Illicit Drugs Other Than Marijuana ²	0.0873		0.9882	
Cigarettes	0.0049		0.6011	
Alcohol	0.0373		0.5807	
Binge Alcohol ³	0.0078		0.3658	
PAST YEAR				
Illicit Drugs ²	0.0095		0.9366	
Marijuana	0.0018		0.8828	
Illicit Drugs Other Than Marijuana ²	0.1259		0.6118	
Cigarettes	0.0134		0.7956	
Alcohol	0.1934		0.3991	
LIFETIME				
Illicit Drugs ²	0.0000		0.4058	
Marijuana	0.0002		0.7571	
Illicit Drugs Other Than Marijuana ²	0.0004		0.1624	
Cigarettes	0.0000		0.1920	
Alcohol	0.0303		0.9831	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.11C Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Have Four or Five Drinks of Alcoholic Beverage Nearly Every Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	112	108	112	108
AGE				
12-13	119	108	75	75
14-15	112	117	88	79
16-17	99	108	82	81
GENDER				
Male	81	80	81	80
Female	71	75	71	75
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	97	91	97	91
White	162	168	113	114
Black or African American	92	94	55	58
American Indian or Alaska Native	*	14	*	9
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	52	67	32	28
Two or More Races	22	21	17	11
Hispanic or Latino	55	55	55	55
COUNTY TYPE				
Large Metro	184	189	117	121
Small Metro	131	120	92	78
Nonmetro	83	86	70	62
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	49	48	55	49
Marijuana	39	40	48	42
Illicit Drugs Other Than Marijuana ²	34	30	39	34
Cigarettes	52	49	62	51
Alcohol	65	63	67	64
Binge Alcohol ³	49	42	57	55
PAST YEAR				
Illicit Drugs ²	72	72	76	69
Marijuana	56	57	65	58
Illicit Drugs Other Than Marijuana ²	54	55	59	54
Cigarettes	68	65	73	70
Alcohol	99	99	92	93
LIFETIME				
Illicit Drugs ²	88	90	90	81
Marijuana	65	64	74	68
Illicit Drugs Other Than Marijuana ²	71	73	76	67
Cigarettes	88	86	89	85
Alcohol	117	114	101	100

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.11D Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Have Four or Five Drinks of Alcoholic Beverage Nearly Every Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.45	0.43	0.45	0.43
AGE				
12-13	0.79	0.71	0.79	0.71
14-15	0.75	0.73	0.75	0.73
16-17	0.72	0.72	0.72	0.72
GENDER				
Male	0.64	0.62	0.64	0.62
Female	0.57	0.61	0.57	0.61
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.47	0.44	0.47	0.44
White	0.53	0.50	0.53	0.50
Black or African American	1.03	1.00	1.03	1.00
American Indian or Alaska Native	*	4.40	*	4.40
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.67	2.24	2.67	2.24
Two or More Races	2.76	2.43	2.76	2.43
Hispanic or Latino	1.30	1.26	1.30	1.26
COUNTY TYPE				
Large Metro	0.67	0.65	0.67	0.65
Small Metro	0.74	0.71	0.74	0.71
Nonmetro	0.97	0.92	0.97	0.92
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.30	0.29	0.50	0.48
Marijuana	0.24	0.24	0.45	0.42
Illicit Drugs Other Than Marijuana ²	0.22	0.18	0.38	0.36
Cigarettes	0.33	0.30	0.57	0.51
Alcohol	0.38	0.37	0.60	0.60
Binge Alcohol ³	0.31	0.25	0.53	0.54
PAST YEAR				
Illicit Drugs ²	0.42	0.40	0.65	0.62
Marijuana	0.34	0.33	0.58	0.53
Illicit Drugs Other Than Marijuana ²	0.33	0.32	0.54	0.53
Cigarettes	0.41	0.39	0.64	0.63
Alcohol	0.51	0.51	0.68	0.69
LIFETIME				
Illicit Drugs ²	0.46	0.47	0.69	0.68
Marijuana	0.38	0.37	0.64	0.61
Illicit Drugs Other Than Marijuana ²	0.39	0.40	0.65	0.63
Cigarettes	0.48	0.48	0.71	0.70
Alcohol	0.55	0.55	0.68	0.67

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.11P Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day (II), among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2005 Versus 2004

(I) Demographic Characteristic and (II) Substance Use	Have Four or Five Drinks of Alcoholic Beverage Nearly Every Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.0012		0.0012	
AGE				
12-13	0.5304		0.5304	
14-15	0.0030		0.0030	
16-17	0.0225		0.0225	
GENDER				
Male	0.0015		0.0015	
Female	0.1548		0.1548	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.0008		0.0008	
White	0.0016		0.0016	
Black or African American	0.7682		0.7682	
American Indian or Alaska Native	*		*	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.0716		0.0716	
Two or More Races	0.2004		0.2004	
Hispanic or Latino	0.4594		0.4594	
COUNTY TYPE				
Large Metro	0.0740		0.0740	
Small Metro	0.0047		0.0047	
Nonmetro	0.3185		0.3185	
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.4631		0.1669	
Marijuana	0.3428		0.0592	
Illicit Drugs Other Than Marijuana ²	0.3043		0.7266	
Cigarettes	0.1919		0.0535	
Alcohol	0.7318		0.0316	
Binge Alcohol ³	0.0296		0.1971	
PAST YEAR				
Illicit Drugs ²	0.5169		0.0523	
Marijuana	0.4921		0.0100	
Illicit Drugs Other Than Marijuana ²	0.4733		0.3542	
Cigarettes	0.0183		0.9864	
Alcohol	0.8123		0.8168	
LIFETIME				
Illicit Drugs ²	0.0411		0.0010	
Marijuana	0.0921		0.0128	
Illicit Drugs Other Than Marijuana ²	0.0595		0.0088	
Cigarettes	0.0014		0.0448	
Alcohol	0.1215		0.5589	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.12C Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Have Five or More Drinks of Alcoholic Beverage Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	101	118	101	118
AGE				
12-13	89	84	102	101
14-15	81	91	117	109
16-17	73	88	106	106
GENDER				
Male	78	78	78	78
Female	76	80	76	80
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	94	105	94	105
White	109	114	165	167
Black or African American	74	79	74	82
American Indian or Alaska Native	8	9	10	13
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	43	56	45	45
Two or More Races	16	15	22	18
Hispanic or Latino	52	55	52	55
COUNTY TYPE				
Large Metro	119	146	166	170
Small Metro	88	89	133	114
Nonmetro	65	62	88	88
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	37	35	64	59
Marijuana	28	28	57	51
Illicit Drugs Other Than Marijuana ²	27	23	45	40
Cigarettes	41	36	68	62
Alcohol	49	43	82	79
Binge Alcohol ³	39	30	67	63
PAST YEAR				
Illicit Drugs ²	55	54	90	87
Marijuana	42	42	78	71
Illicit Drugs Other Than Marijuana ²	44	39	67	68
Cigarettes	52	50	85	83
Alcohol	71	69	119	118
LIFETIME				
Illicit Drugs ²	67	68	105	102
Marijuana	52	47	85	82
Illicit Drugs Other Than Marijuana ²	54	55	87	84
Cigarettes	70	67	105	103
Alcohol	82	81	129	130

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.12D Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2004 and 2005

(I) Demographic Characteristic and (II) Substance Use	Have Five or More Drinks of Alcoholic Beverage Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.40	0.47	0.40	0.47
AGE				
12-13	0.79	0.74	0.79	0.74
14-15	0.72	0.78	0.72	0.78
16-17	0.68	0.77	0.68	0.77
GENDER				
Male	0.61	0.61	0.61	0.61
Female	0.62	0.65	0.62	0.65
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.45	0.51	0.45	0.51
White	0.52	0.50	0.52	0.50
Black or African American	1.09	1.21	1.09	1.21
American Indian or Alaska Native	3.82	3.52	3.82	3.52
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	3.04	3.05	3.04	3.05
Two or More Races	2.60	2.88	2.60	2.88
Hispanic or Latino	1.25	1.27	1.25	1.27
COUNTY TYPE				
Large Metro	0.58	0.71	0.58	0.71
Small Metro	0.71	0.78	0.71	0.78
Nonmetro	0.99	0.94	0.99	0.94
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.38	0.35	0.37	0.35
Marijuana	0.28	0.28	0.34	0.31
Illicit Drugs Other Than Marijuana ²	0.28	0.23	0.27	0.24
Cigarettes	0.41	0.36	0.39	0.37
Alcohol	0.48	0.42	0.44	0.44
Binge Alcohol ³	0.40	0.30	0.39	0.37
PAST YEAR				
Illicit Drugs ²	0.54	0.50	0.47	0.47
Marijuana	0.42	0.40	0.43	0.40
Illicit Drugs Other Than Marijuana ²	0.44	0.38	0.38	0.39
Cigarettes	0.51	0.49	0.47	0.46
Alcohol	0.64	0.61	0.52	0.54
LIFETIME				
Illicit Drugs ²	0.61	0.59	0.49	0.50
Marijuana	0.51	0.45	0.46	0.45
Illicit Drugs Other Than Marijuana ²	0.51	0.50	0.44	0.46
Cigarettes	0.65	0.61	0.54	0.53
Alcohol	0.67	0.67	0.53	0.55

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 3.12P Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Demographic Characteristics (I), and Selected Substance Use, by Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week (II), among Persons Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

(I) Demographic Characteristic and (II) Substance Use	Have Five or More Drinks of Alcoholic Beverage Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2004	2005	2004	2005
(I) Demographic Characteristic				
TOTAL	0.5955		0.5955	
AGE				
12-13	0.6458		0.6458	
14-15	0.5215		0.5215	
16-17	0.3706		0.3706	
GENDER				
Male	0.2896		0.2896	
Female	0.7632		0.7632	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.5411		0.5411	
White	0.7813		0.7813	
Black or African American	0.9763		0.9763	
American Indian or Alaska Native	0.4862		0.4862	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.4280		0.4280	
Two or More Races	0.8086		0.8086	
Hispanic or Latino	0.9484		0.9484	
COUNTY TYPE				
Large Metro	0.3964		0.3964	
Small Metro	0.0086		0.0086	
Nonmetro	0.6645		0.6645	
(II) Substance Use				
PAST MONTH				
Illicit Drugs ²	0.2210		0.1300	
Marijuana	0.1595		0.0517	
Illicit Drugs Other Than Marijuana ²	0.2533		0.4337	
Cigarettes	0.0111		0.0839	
Alcohol	0.0206		0.2528	
Binge Alcohol ³	0.0020		0.1221	
PAST YEAR				
Illicit Drugs ²	0.3805		0.0284	
Marijuana	0.3458		0.0080	
Illicit Drugs Other Than Marijuana ²	0.3858		0.2027	
Cigarettes	0.0385		0.1990	
Alcohol	0.4333		0.6888	
LIFETIME				
Illicit Drugs ²	0.0067		0.0007	
Marijuana	0.0073		0.0175	
Illicit Drugs Other Than Marijuana ²	0.0539		0.0015	
Cigarettes	0.0001		0.0185	
Alcohol	0.1062		0.1558	

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.