

Table 5.67C Needed and Received Treatment for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

| Demographic Characteristic | NEEDED TREATMENT FOR AN ALCOHOL PROBLEM IN THE PAST YEAR | | | | | | Percentage Who Received Treatment at a Specialty Facility among Persons Who Needed Treatment | |
|---|--|------|--|------|---|------|--|------|
| | Total | | Received Treatment at a Specialty Facility | | Did Not Receive Treatment at a Specialty Facility | | | |
| | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 |
| TOTAL | 431 | 413 | 135 | 139 | 409 | 386 | 0.66 | 0.68 |
| AGE | | | | | | | | |
| 12-17 | 52 | 52 | 15 | 14 | 49 | 50 | 0.91 | 0.91 |
| 18-25 | 121 | 121 | 28 | 25 | 118 | 118 | 0.48 | 0.43 |
| 26 or Older | 389 | 369 | 131 | 135 | 367 | 345 | 1.03 | 1.05 |
| GENDER | | | | | | | | |
| Male | 331 | 329 | 129 | 116 | 311 | 308 | 0.94 | 0.86 |
| Female | 216 | 218 | 47 | 73 | 209 | 205 | 0.72 | 1.03 |
| HISPANIC ORIGIN AND RACE | | | | | | | | |
| Not Hispanic or Latino | 393 | 366 | 98 | 122 | 380 | 346 | 0.57 | 0.70 |
| White | 361 | 323 | 74 | 91 | 350 | 314 | 0.52 | 0.64 |
| Black or African American | 134 | 149 | 58 | 79 | 121 | 127 | 2.76 | 3.33 |
| American Indian or Alaska Native | 37 | 54 | 14 | 6 | 35 | 53 | * | 2.94 |
| Native Hawaiian or Other Pacific Islander | * | 18 | * | 0 | 13 | 18 | * | * |
| Asian | 61 | 54 | 9 | 3 | 60 | 54 | * | 0.78 |
| Two or More Races | 41 | 25 | 11 | 5 | 40 | 25 | 3.82 | 2.67 |
| Hispanic or Latino | 165 | 168 | 94 | 60 | 137 | 153 | 3.23 | 2.02 |

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 5.67D Needed and Received Treatment for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2004 and 2005

| Demographic Characteristic | NEEDED TREATMENT FOR AN ALCOHOL PROBLEM IN THE PAST YEAR | | | | | | Percentage Who Received Treatment at a Specialty Facility among Persons Who Needed Treatment | |
|---|--|------|--|------|---|------|--|------|
| | Total | | Received Treatment at a Specialty Facility | | Did Not Receive Treatment at a Specialty Facility | | | |
| | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 |
| TOTAL | 0.18 | 0.17 | 0.06 | 0.06 | 0.17 | 0.16 | 0.66 | 0.68 |
| AGE | | | | | | | | |
| 12-17 | 0.20 | 0.21 | 0.06 | 0.05 | 0.20 | 0.20 | 0.91 | 0.91 |
| 18-25 | 0.38 | 0.37 | 0.09 | 0.08 | 0.37 | 0.36 | 0.48 | 0.43 |
| 26 or Older | 0.21 | 0.20 | 0.07 | 0.07 | 0.20 | 0.19 | 1.03 | 1.05 |
| GENDER | | | | | | | | |
| Male | 0.28 | 0.28 | 0.11 | 0.10 | 0.27 | 0.26 | 0.94 | 0.86 |
| Female | 0.17 | 0.17 | 0.04 | 0.06 | 0.17 | 0.16 | 0.72 | 1.03 |
| HISPANIC ORIGIN AND RACE | | | | | | | | |
| Not Hispanic or Latino | 0.19 | 0.17 | 0.05 | 0.06 | 0.18 | 0.16 | 0.57 | 0.70 |
| White | 0.22 | 0.19 | 0.04 | 0.05 | 0.21 | 0.19 | 0.52 | 0.64 |
| Black or African American | 0.48 | 0.52 | 0.21 | 0.28 | 0.43 | 0.44 | 2.76 | 3.33 |
| American Indian or Alaska Native | 2.87 | 3.62 | 1.04 | 0.47 | 2.68 | 3.63 | * | 2.94 |
| Native Hawaiian or Other Pacific Islander | * | 2.60 | * | 0.02 | 2.24 | 2.60 | * | * |
| Asian | 0.62 | 0.54 | 0.09 | 0.03 | 0.61 | 0.54 | * | 0.78 |
| Two or More Races | 1.56 | 1.01 | 0.42 | 0.21 | 1.49 | 0.98 | 3.82 | 2.67 |
| Hispanic or Latino | 0.53 | 0.52 | 0.30 | 0.19 | 0.44 | 0.47 | 3.23 | 2.02 |

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 5.67P Needed and Received Treatment for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

| Demographic Characteristic | NEEDED TREATMENT FOR AN ALCOHOL PROBLEM IN THE PAST YEAR | | | | | | Percentage Who Received Treatment at a Specialty Facility among Persons Who Needed Treatment |
|---|--|------|--|------|---|------|--|
| | Total | | Received Treatment at a Specialty Facility | | Did Not Receive Treatment at a Specialty Facility | | |
| | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 | |
| TOTAL | 0.7398 | | 0.7067 | | 0.6295 | | 0.6267 |
| AGE | | | | | | | |
| 12-17 | 0.1061 | | 0.6841 | | 0.1154 | | 0.9471 |
| 18-25 | 0.9073 | | 0.4040 | | 0.7577 | | 0.3779 |
| 26 or Older | 0.8629 | | 0.5476 | | 0.6830 | | 0.4910 |
| GENDER | | | | | | | |
| Male | 0.2437 | | 0.5060 | | 0.3271 | | 0.6727 |
| Female | 0.2700 | | 0.0296 | | 0.6073 | | 0.0456 |
| HISPANIC ORIGIN AND RACE | | | | | | | |
| Not Hispanic or Latino | 0.7710 | | 0.6899 | | 0.6709 | | 0.6172 |
| White | 0.6176 | | 0.7503 | | 0.5525 | | 0.6356 |
| Black or African American | 0.5371 | | 0.4396 | | 0.7847 | | 0.5192 |
| American Indian or Alaska Native | 0.3828 | | 0.7642 | | 0.3321 | | * |
| Native Hawaiian or Other Pacific Islander | * | | * | | 0.3121 | | * |
| Asian | 0.9968 | | 0.4225 | | 0.9262 | | * |
| Two or More Races | 0.0936 | | 0.1019 | | 0.1892 | | 0.2451 |
| Hispanic or Latino | 0.8426 | | 0.9314 | | 0.7828 | | 0.8929 |

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 5.68C Needed and Received Treatment for an Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

| Demographic Characteristic | NEEDED TREATMENT FOR AN ALCOHOL PROBLEM IN THE PAST YEAR | | | | | | Percentage Who Received Treatment at a Specialty Facility among Persons Who Needed Treatment | |
|---|--|------|--|------|---|------|--|------|
| | Total | | Received Treatment at a Specialty Facility | | Did Not Receive Treatment at a Specialty Facility | | | |
| | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 |
| TOTAL | 425 | 407 | 134 | 138 | 404 | 381 | 0.72 | 0.72 |
| GENDER | | | | | | | | |
| Male | 327 | 327 | 128 | 115 | 307 | 306 | 0.99 | 0.91 |
| Female | 213 | 215 | 46 | 72 | 207 | 203 | 0.81 | 1.14 |
| HISPANIC ORIGIN AND RACE | | | | | | | | |
| Not Hispanic or Latino | 391 | 363 | 96 | 121 | 379 | 345 | 0.60 | 0.74 |
| White | 363 | 346 | 72 | 90 | 352 | 336 | 0.54 | 0.69 |
| Black or African American | 139 | 156 | 58 | 79 | 125 | 132 | 2.93 | 3.45 |
| American Indian or Alaska Native | 36 | 53 | 13 | 6 | 33 | 52 | * | 2.98 |
| Native Hawaiian or Other Pacific Islander | * | 18 | * | 0 | 13 | 18 | * | * |
| Asian | 61 | 53 | 9 | 3 | 60 | 53 | * | 0.81 |
| Two or More Races | 40 | 25 | 11 | 5 | 38 | 24 | * | * |
| Hispanic or Latino | 162 | 164 | 94 | 60 | 137 | 149 | * | 2.23 |
| EDUCATION | | | | | | | | |
| < High School | 174 | 188 | 80 | 74 | 155 | 167 | 2.27 | 2.14 |
| High School Graduate | 201 | 208 | 61 | 71 | 194 | 197 | 1.11 | 1.29 |
| Some College | 219 | 211 | 62 | 61 | 209 | 201 | 1.17 | 1.12 |
| College Graduate | 228 | 236 | 26 | 69 | 226 | 227 | 0.61 | 1.51 |
| CURRENT EMPLOYMENT | | | | | | | | |
| Full-Time | 336 | 335 | 76 | 74 | 326 | 321 | 0.66 | 0.64 |
| Part-Time | 164 | 143 | 34 | 43 | 160 | 136 | 1.17 | 1.51 |
| Unemployed | 91 | 91 | 32 | 37 | 84 | 81 | 2.50 | 3.08 |
| Other ¹ | 158 | 185 | 79 | 95 | 134 | 159 | 2.71 | 2.79 |

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 5.68D Needed and Received Treatment for an Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2004 and 2005

| Demographic Characteristic | NEEDED TREATMENT FOR AN ALCOHOL PROBLEM IN THE PAST YEAR | | | | | | Percentage Who Received Treatment at a Specialty Facility among Persons Who Needed Treatment | |
|---|--|------|--|------|---|------|--|------|
| | Total | | Received Treatment at a Specialty Facility | | Did Not Receive Treatment at a Specialty Facility | | | |
| | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 |
| TOTAL | 0.20 | 0.19 | 0.06 | 0.06 | 0.19 | 0.18 | 0.72 | 0.72 |
| GENDER | | | | | | | | |
| Male | 0.32 | 0.31 | 0.12 | 0.11 | 0.30 | 0.29 | 0.99 | 0.91 |
| Female | 0.19 | 0.19 | 0.04 | 0.06 | 0.19 | 0.18 | 0.81 | 1.14 |
| HISPANIC ORIGIN AND RACE | | | | | | | | |
| Not Hispanic or Latino | 0.21 | 0.19 | 0.05 | 0.06 | 0.20 | 0.18 | 0.60 | 0.74 |
| White | 0.24 | 0.21 | 0.05 | 0.06 | 0.23 | 0.21 | 0.54 | 0.69 |
| Black or African American | 0.55 | 0.60 | 0.24 | 0.32 | 0.49 | 0.51 | 2.93 | 3.45 |
| American Indian or Alaska Native | 3.26 | 4.13 | 1.19 | 0.52 | 3.00 | 4.14 | * | 2.98 |
| Native Hawaiian or Other Pacific Islander | * | 2.90 | * | 0.03 | 2.50 | 2.90 | * | * |
| Asian | 0.68 | 0.61 | 0.10 | 0.03 | 0.67 | 0.60 | * | 0.81 |
| Two or More Races | 1.84 | 1.19 | 0.51 | 0.24 | 1.75 | 1.15 | * | * |
| Hispanic or Latino | 0.60 | 0.59 | 0.35 | 0.22 | 0.51 | 0.54 | * | 2.23 |
| EDUCATION | | | | | | | | |
| < High School | 0.47 | 0.51 | 0.22 | 0.21 | 0.42 | 0.45 | 2.27 | 2.14 |
| High School Graduate | 0.30 | 0.29 | 0.09 | 0.10 | 0.28 | 0.28 | 1.11 | 1.29 |
| Some College | 0.39 | 0.37 | 0.11 | 0.11 | 0.37 | 0.35 | 1.17 | 1.12 |
| College Graduate | 0.39 | 0.38 | 0.05 | 0.12 | 0.39 | 0.37 | 0.61 | 1.51 |
| CURRENT EMPLOYMENT | | | | | | | | |
| Full-Time | 0.28 | 0.26 | 0.06 | 0.06 | 0.27 | 0.25 | 0.66 | 0.64 |
| Part-Time | 0.57 | 0.48 | 0.12 | 0.15 | 0.56 | 0.46 | 1.17 | 1.51 |
| Unemployed | 1.18 | 1.13 | 0.43 | 0.48 | 1.11 | 1.01 | 2.50 | 3.08 |
| Other ¹ | 0.25 | 0.29 | 0.13 | 0.15 | 0.22 | 0.25 | 2.71 | 2.79 |

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 5.68P Needed and Received Treatment for an Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

| Demographic Characteristic | NEEDED TREATMENT FOR AN ALCOHOL PROBLEM IN THE PAST YEAR | | | | | | Percentage Who Received Treatment at a Specialty Facility among Persons Who Needed Treatment |
|---|--|------|--|------|---|------|--|
| | Total | | Received Treatment at a Specialty Facility | | Did Not Receive Treatment at a Specialty Facility | | |
| | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 | |
| TOTAL | 0.8878 | | 0.6725 | | 0.7680 | | 0.6289 |
| GENDER | | | | | | | |
| Male | 0.3445 | | 0.5696 | | 0.4343 | | 0.7094 |
| Female | 0.2542 | | 0.0361 | | 0.5643 | | 0.0565 |
| HISPANIC ORIGIN AND RACE | | | | | | | |
| Not Hispanic or Latino | 0.8897 | | 0.6401 | | 0.7759 | | 0.5985 |
| White | 0.6336 | | 0.6481 | | 0.5465 | | 0.5441 |
| Black or African American | 0.4109 | | 0.4737 | | 0.5898 | | 0.6181 |
| American Indian or Alaska Native | 0.3045 | | 0.7597 | | 0.2564 | | * |
| Native Hawaiian or Other Pacific Islander | * | | * | | 0.4089 | | * |
| Asian | 0.9720 | | 0.4395 | | 0.8974 | | * |
| Two or More Races | 0.1740 | | 0.0940 | | 0.3331 | | * |
| Hispanic or Latino | 0.9528 | | 0.9434 | | 0.9151 | | * |
| EDUCATION | | | | | | | |
| < High School | 0.8586 | | 0.9077 | | 0.8863 | | 0.9479 |
| High School Graduate | 0.7984 | | 0.9453 | | 0.7721 | | 0.8807 |
| Some College | 0.9739 | | 0.7246 | | 0.8865 | | 0.7076 |
| College Graduate | 0.8902 | | 0.0671 | | 0.7716 | | 0.0612 |
| CURRENT EMPLOYMENT | | | | | | | |
| Full-Time | 0.5961 | | 0.5409 | | 0.6876 | | 0.6206 |
| Part-Time | 0.4078 | | 0.6604 | | 0.3298 | | 0.5115 |
| Unemployed | 0.1965 | | 0.6973 | | 0.1171 | | 0.3728 |
| Other ¹ | 0.0612 | | 0.4029 | | 0.0966 | | 0.8130 |

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 5.69C Needed and Received Treatment for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

| Demographic Characteristic | NEEDED TREATMENT FOR AN ALCOHOL PROBLEM IN THE PAST YEAR | | | | | | Percentage Who Received Treatment at a Specialty Facility among Persons Who Needed Treatment | |
|--|---|-----------|---|-----------|---|-----------|--|-----------|
| | Total | | Received Treatment at a Specialty Facility | | Did Not Receive Treatment at a Specialty Facility | | | |
| | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 |
| TOTAL | 298 | 299 | 77 | 97 | 288 | 281 | 0.41 | 0.47 |
| AGE | | | | | | | | |
| 12-17 | 38 | 37 | 10 | 10 | 37 | 35 | 0.67 | 0.64 |
| 18-25 | 79 | 85 | 17 | 19 | 78 | 84 | 0.31 | 0.32 |
| 26 or Older | 276 | 268 | 74 | 94 | 268 | 252 | 0.64 | 0.73 |
| GENDER | | | | | | | | |
| Male | 239 | 234 | 69 | 87 | 232 | 219 | 0.54 | 0.64 |
| Female | 161 | 154 | 34 | 43 | 155 | 147 | 0.54 | 0.64 |
| HISPANIC ORIGIN AND RACE | | | | | | | | |
| Not Hispanic or Latino | 273 | 268 | 74 | 78 | 264 | 257 | 0.45 | 0.45 |
| White | 236 | 242 | 62 | 59 | 230 | 235 | 0.46 | 0.41 |
| Black or African American | 93 | 100 | 37 | 49 | 91 | 88 | 1.91 | 2.19 |
| American Indian or Alaska Native | 24 | 33 | 9 | 7 | 22 | 32 | 4.18 | 3.41 |
| Native Hawaiian or Other Pacific Islander | 10 | 15 | 2 | * | 10 | 11 | * | * |
| Asian | 46 | 41 | 4 | 5 | 46 | 40 | 0.83 | 1.27 |
| Two or More Races | 31 | 24 | 6 | 6 | 30 | 23 | 2.46 | 2.47 |
| Hispanic or Latino | 97 | 118 | 23 | 56 | 95 | 103 | 0.90 | 1.90 |

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 5.69D Needed and Received Treatment for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

| Demographic Characteristic | NEEDED TREATMENT FOR AN ALCOHOL PROBLEM IN THE PAST YEAR | | | | | | Percentage Who Received Treatment at a Specialty Facility among Persons Who Needed Treatment | |
|--|---|-----------|---|-----------|---|-----------|--|-----------|
| | Total | | Received Treatment at a Specialty Facility | | Did Not Receive Treatment at a Specialty Facility | | | |
| | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 |
| TOTAL | 0.13 | 0.12 | 0.03 | 0.04 | 0.12 | 0.12 | 0.41 | 0.47 |
| AGE | | | | | | | | |
| 12-17 | 0.15 | 0.15 | 0.04 | 0.04 | 0.15 | 0.14 | 0.67 | 0.64 |
| 18-25 | 0.25 | 0.26 | 0.06 | 0.06 | 0.25 | 0.26 | 0.31 | 0.32 |
| 26 or Older | 0.15 | 0.15 | 0.04 | 0.05 | 0.15 | 0.14 | 0.64 | 0.73 |
| GENDER | | | | | | | | |
| Male | 0.21 | 0.20 | 0.06 | 0.07 | 0.20 | 0.19 | 0.54 | 0.64 |
| Female | 0.13 | 0.12 | 0.03 | 0.03 | 0.13 | 0.12 | 0.54 | 0.64 |
| HISPANIC ORIGIN AND RACE | | | | | | | | |
| Not Hispanic or Latino | 0.13 | 0.13 | 0.04 | 0.04 | 0.13 | 0.12 | 0.45 | 0.45 |
| White | 0.14 | 0.14 | 0.04 | 0.03 | 0.14 | 0.14 | 0.46 | 0.41 |
| Black or African American | 0.34 | 0.35 | 0.14 | 0.17 | 0.34 | 0.31 | 1.91 | 2.19 |
| American Indian or Alaska Native | 1.47 | 2.34 | 0.64 | 0.58 | 1.36 | 2.29 | 4.18 | 3.41 |
| Native Hawaiian or Other Pacific Islander | 1.65 | 2.24 | 0.30 | * | 1.62 | 1.76 | * | * |
| Asian | 0.50 | 0.41 | 0.04 | 0.05 | 0.49 | 0.41 | 0.83 | 1.27 |
| Two or More Races | 1.14 | 0.93 | 0.23 | 0.23 | 1.11 | 0.89 | 2.46 | 2.47 |
| Hispanic or Latino | 0.33 | 0.37 | 0.08 | 0.18 | 0.32 | 0.32 | 0.90 | 1.90 |

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 5.69P Needed and Received Treatment for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

| Demographic Characteristic | NEEDED TREATMENT FOR AN ALCOHOL PROBLEM IN THE PAST YEAR | | | | | | Percentage Who Received Treatment at a Specialty Facility among Persons Who Needed Treatment | |
|--|---|-----------|---|-----------|---|-----------|--|-----------|
| | Total | | Received Treatment at a Specialty Facility | | Did Not Receive Treatment at a Specialty Facility | | | |
| | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 |
| TOTAL | 0.2321 | | 0.3269 | | 0.3442 | | 0.4927 | |
| AGE | | | | | | | | |
| 12-17 | 0.8813 | | 0.3991 | | 0.6896 | | 0.3547 | |
| 18-25 | 0.8490 | | 0.6261 | | 0.9306 | | 0.6508 | |
| 26 or Older | 0.2268 | | 0.4206 | | 0.3201 | | 0.6422 | |
| GENDER | | | | | | | | |
| Male | 0.6326 | | 0.5124 | | 0.7882 | | 0.5756 | |
| Female | 0.1281 | | 0.3773 | | 0.1714 | | 0.6459 | |
| HISPANIC ORIGIN AND RACE | | | | | | | | |
| Not Hispanic or Latino | 0.2336 | | 0.7697 | | 0.2497 | | 0.9631 | |
| White | 0.1303 | | 0.6720 | | 0.0956 | | 0.3815 | |
| Black or African American | 0.8892 | | 0.2583 | | 0.6844 | | 0.2416 | |
| American Indian or Alaska Native | 0.2922 | | 0.3702 | | 0.1603 | | 0.1464 | |
| Native Hawaiian or Other Pacific Islander | 0.6905 | | * | | 0.9285 | | * | |
| Asian | 0.2060 | | 0.7192 | | 0.1890 | | 0.5824 | |
| Two or More Races | 0.8020 | | 0.9832 | | 0.7987 | | 0.9367 | |
| Hispanic or Latino | 0.7920 | | 0.1278 | | 0.7063 | | 0.1109 | |

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 5.70C Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005

| Demographic Characteristic | Total Needing But Not Receiving Treatment ¹ | | FELT NEED FOR TREATMENT ² | | | | | | Did Not Feel Need for Treatment ² | |
|----------------------------|--|------|--------------------------------------|------|-------------|------|----------------|------|--|------|
| | | | Total | | Made Effort | | Made No Effort | | | |
| | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 |
| TOTAL | 423 | 408 | 82 | 80 | 56 | 37 | 60 | 70 | 413 | 410 |
| AGE | | | | | | | | | | |
| 12-17 | 52 | 53 | 9 | 9 | 5 | 2 | 8 | 9 | 53 | 51 |
| 18-25 | 141 | 148 | 24 | 18 | 13 | 9 | 18 | 15 | 135 | 146 |
| 26 or Older | 372 | 362 | 78 | 78 | 54 | 36 | 57 | 70 | 363 | 364 |
| GENDER | | | | | | | | | | |
| Male | 339 | 326 | 66 | 75 | 38 | 35 | 53 | 66 | 332 | 328 |
| Female | 206 | 215 | 45 | 31 | 33 | 14 | 30 | 28 | 203 | 213 |

*Low precision; no estimate reported.

¹ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

² Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 5.70D Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: Standard Errors of Percentages, 2004 and 2005

| Demographic Characteristic | Total Needing But Not Receiving Treatment ¹ | | FELT NEED FOR TREATMENT ² | | | | | | Did Not Feel Need for Treatment ² | |
|----------------------------|--|------|--------------------------------------|------|-------------|------|----------------|------|--|------|
| | | | Total | | Made Effort | | Made No Effort | | | |
| | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 |
| TOTAL | 0.00 | 0.00 | 0.45 | 0.45 | 0.31 | 0.21 | 0.34 | 0.40 | 0.45 | 0.45 |
| AGE | | | | | | | | | | |
| 12-17 | 0.00 | 0.00 | 0.65 | 0.69 | 0.31 | 0.13 | 0.57 | 0.68 | 0.65 | 0.69 |
| 18-25 | 0.00 | 0.00 | 0.43 | 0.32 | 0.24 | 0.16 | 0.33 | 0.27 | 0.43 | 0.32 |
| 26 or Older | 0.00 | 0.00 | 0.70 | 0.72 | 0.48 | 0.33 | 0.51 | 0.64 | 0.70 | 0.72 |
| GENDER | | | | | | | | | | |
| Male | 0.00 | 0.00 | 0.54 | 0.64 | 0.31 | 0.30 | 0.44 | 0.57 | 0.54 | 0.64 |
| Female | 0.00 | 0.00 | 0.75 | 0.50 | 0.56 | 0.22 | 0.50 | 0.45 | 0.75 | 0.50 |

*Low precision; no estimate reported.

¹ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

² Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 5.70P Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

| Demographic Characteristic | Total Needing But Not Receiving Treatment ¹ | | FELT NEED FOR TREATMENT ² | | | | | | Did Not Feel Need for Treatment ² | |
|----------------------------|--|------|--------------------------------------|------|-------------|------|----------------|------|--|------|
| | | | Total | | Made Effort | | Made No Effort | | | |
| | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 | 2004 | 2005 |
| TOTAL | . | | 0.4811 | | 0.0543 | | 0.6066 | | 0.4811 | |
| AGE | | | | | | | | | | |
| 12-17 | . | | 0.7684 | | 0.1077 | | 0.7626 | | 0.7684 | |
| 18-25 | . | | 0.1332 | | 0.0925 | | 0.4717 | | 0.1332 | |
| 26 or Older | . | | 0.7682 | | 0.1467 | | 0.4966 | | 0.7682 | |
| GENDER | | | | | | | | | | |
| Male | . | | 0.8769 | | 0.5305 | | 0.5766 | | 0.8769 | |
| Female | . | | 0.0794 | | 0.0079 | | 0.9714 | | 0.0794 | |

*Low precision; no estimate reported.

¹ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

² Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 5.71C Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

| Demographic Characteristic | Total Needing But Not Receiving Treatment ¹ | | FELT NEED FOR TREATMENT ² | | | | | | Did Not Feel Need for Treatment ² | |
|-------------------------------|---|-----------|--------------------------------------|-----------|-------------|-----------|----------------|-----------|---|-----------|
| | | | Total | | Made Effort | | Made No Effort | | | |
| | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 |
| TOTAL | 313 | 294 | 59 | 57 | 32 | 33 | 48 | 46 | 303 | 291 |
| AGE | | | | | | | | | | |
| 12-17 | 38 | 37 | 5 | 7 | 4 | 2 | 4 | 6 | 38 | 37 |
| 18-25 | 102 | 102 | 14 | 15 | 6 | 8 | 13 | 12 | 101 | 100 |
| 26 or Older | 288 | 260 | 58 | 55 | 31 | 32 | 47 | 45 | 279 | 257 |
| GENDER | | | | | | | | | | |
| Male | 249 | 235 | 49 | 50 | 27 | 26 | 40 | 42 | 240 | 234 |
| Female | 159 | 149 | 33 | 27 | 18 | 18 | 28 | 20 | 153 | 147 |

*Low precision; no estimate reported.

¹ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

² Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 5.71D Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

| Demographic Characteristic | Total Needing But Not Receiving Treatment ¹ | | FELT NEED FOR TREATMENT ² | | | | | | Did Not Feel Need for Treatment ² | |
|-------------------------------|---|-----------|--------------------------------------|-----------|-------------|-----------|----------------|-----------|---|-----------|
| | | | Total | | Made Effort | | Made No Effort | | | |
| | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 |
| TOTAL | 0.00 | 0.00 | 0.34 | 0.32 | 0.18 | 0.19 | 0.28 | 0.26 | 0.34 | 0.32 |
| AGE | | | | | | | | | | |
| 12-17 | 0.00 | 0.00 | 0.37 | 0.47 | 0.27 | 0.17 | 0.26 | 0.44 | 0.37 | 0.47 |
| 18-25 | 0.00 | 0.00 | 0.27 | 0.27 | 0.11 | 0.15 | 0.24 | 0.21 | 0.27 | 0.27 |
| 26 or Older | 0.00 | 0.00 | 0.54 | 0.50 | 0.29 | 0.29 | 0.45 | 0.41 | 0.54 | 0.50 |
| GENDER | | | | | | | | | | |
| Male | 0.00 | 0.00 | 0.42 | 0.42 | 0.23 | 0.22 | 0.34 | 0.36 | 0.42 | 0.42 |
| Female | 0.00 | 0.00 | 0.57 | 0.45 | 0.32 | 0.30 | 0.49 | 0.34 | 0.57 | 0.45 |

*Low precision; no estimate reported.

¹ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

² Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 5.71P Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

| Demographic Characteristic | Total Needing But Not Receiving Treatment ¹ | | FELT NEED FOR TREATMENT ² | | | | | | Did Not Feel Need for Treatment ² | |
|-------------------------------|---|-----------|--------------------------------------|-----------|-------------|-----------|----------------|-----------|---|-----------|
| | | | Total | | Made Effort | | Made No Effort | | | |
| | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 | 2002-2003 | 2004-2005 |
| TOTAL | . | | 0.5255 | | 0.8089 | | 0.5447 | | 0.5255 | |
| AGE | | | | | | | | | | |
| 12-17 | . | | 0.7926 | | 0.5821 | | 0.5160 | | 0.7926 | |
| 18-25 | . | | 0.8157 | | 0.0367 | | 0.3524 | | 0.8157 | |
| 26 or Older | . | | 0.4357 | | 0.4912 | | 0.6403 | | 0.4357 | |
| GENDER | | | | | | | | | | |
| Male | . | | 0.9061 | | 0.6747 | | 0.6819 | | 0.9061 | |
| Female | . | | 0.1711 | | 0.8717 | | 0.0673 | | 0.1711 | |

*Low precision; no estimate reported.

¹ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

² Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.