

**Table 6.24A Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Numbers in Thousands, 2004 and 2005**

Drug	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
<b>ILLICIT DRUGS<sup>3</sup></b>	29,506	29,999	5,916	5,566	23,504	24,308
Marijuana and Hashish	21,796	21,991	4,010	3,679	17,739	18,212
Cocaine	5,247	5,099	1,140	1,150	4,104	3,877
Crack	1,241	1,327	398	422	841	903
Heroin	350	343	126	134	223	199
Hallucinogens	3,125	3,144	596	581	2,518	2,549
LSD	434	413	77	76	351	333
PCP	138 <sup>a</sup>	79	24	31	113 <sup>a</sup>	48
Ecstasy	1,605	1,703	287	307	1,311	1,389
Inhalants	1,087	1,043	281	252	806	789
Nonmedical Use of Psychotherapeutics <sup>4</sup>	12,418	13,068	3,047	2,908	9,321	10,105
Pain Relievers	9,391	10,074	2,178	2,072	7,190 <sup>a</sup>	7,950
OxyContin <sup>®</sup>	1,019	1,039	305	321	712	708
Tranquilizers	4,536	4,752	1,514	1,308	3,014	3,414
Stimulants	2,423	2,268	637	617	1,782	1,648
Methamphetamine	1,277	1,127	357	311	918	815
Sedatives	604	632	240	217	338	415
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>3</sup></b>	16,389	16,980	3,771	3,637	12,563	13,237

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column represent past year use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

<sup>3</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>4</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.24B Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Percentages, 2004 and 2005**

Drug	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>3</sup></b>	13.7	13.8	21.5	19.8	12.5	12.9
Marijuana and Hashish	10.1	10.1	14.6	13.1	9.5	9.6
Cocaine	2.4	2.3	4.1	4.1	2.2	2.1
Crack	0.6	0.6	1.4	1.5	0.4	0.5
Heroin	0.2	0.2	0.5	0.5	0.1	0.1
Hallucinogens	1.5	1.4	2.2	2.1	1.3	1.3
LSD	0.2	0.2	0.3	0.3	0.2	0.2
PCP	0.1 <sup>a</sup>	0.0	0.1	0.1	0.1 <sup>a</sup>	0.0
Ecstasy	0.7	0.8	1.0	1.1	0.7	0.7
Inhalants	0.5	0.5	1.0	0.9	0.4	0.4
Nonmedical Use of Psychotherapeutics <sup>4</sup>	5.8	6.0	11.1	10.3	5.0	5.3
Pain Relievers	4.4	4.6	7.9	7.4	3.8	4.2
OxyContin <sup>®</sup>	0.5	0.5	1.1	1.1	0.4	0.4
Tranquilizers	2.1	2.2	5.5	4.6	1.6	1.8
Stimulants	1.1	1.0	2.3	2.2	1.0	0.9
Methamphetamine	0.6	0.5	1.3	1.1	0.5	0.4
Sedatives	0.3	0.3	0.9	0.8	0.2	0.2
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>3</sup></b>	7.6	7.8	13.7	12.9	6.7	7.0

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column represent past year use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

<sup>3</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>4</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.25A Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Numbers in Thousands, 2004 and 2005**

Substance	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
<b>PAST YEAR</b>						
Tobacco Products <sup>3</sup>	77,504	79,593	11,594	12,453	65,727	66,904
Cigarettes	65,264	66,446	10,423 <sup>a</sup>	11,400	54,677	54,829
Smokeless Tobacco	8,672	9,338	992	918	7,665	8,383
Cigars	23,405	24,423	2,965	2,923	20,374	21,457
Pipe Tobacco <sup>4</sup>	--	--	--	--	--	--
Alcohol	148,149 <sup>b</sup>	153,172	19,668	20,056	128,252 <sup>a</sup>	132,774
Binge Alcohol Use <sup>5</sup>	--	--	--	--	--	--
Heavy Alcohol Use <sup>5</sup>	--	--	--	--	--	--
<b>PAST MONTH</b>						
Tobacco Products <sup>3</sup>	66,622	68,195	10,009	10,863	56,443	57,113
Cigarettes	56,897	57,799	9,151 <sup>a</sup>	10,083	47,592	47,514
Smokeless Tobacco	6,583	7,158	731	666	5,842	6,465
Cigars	12,526	12,576	1,413	1,460	11,073	11,098
Pipe Tobacco	1,657	2,032	239	313	1,412	1,713
Alcohol	116,498 <sup>b</sup>	121,837	15,479	15,685	100,829 <sup>a</sup>	105,886
Binge Alcohol Use <sup>5</sup>	51,931	52,577	6,579	6,292	45,283	46,133
Heavy Alcohol Use <sup>5</sup>	16,018	15,424	2,068	1,952	13,920	13,446

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.<sup>1</sup> Estimates in the Total column represent past year and past month use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.<sup>2</sup> Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.<sup>3</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.<sup>4</sup> Information about past year use of pipe tobacco was not collected.<sup>5</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.25B Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Percentages, 2004 and 2005**

Substance	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
<b>PAST YEAR</b>						
Tobacco Products <sup>3</sup>	36.0	36.5	42.2	44.2	35.1	35.4
Cigarettes	30.3	30.5	37.9	40.5	29.2	29.0
Smokeless Tobacco	4.0	4.3	3.6	3.3	4.1	4.4
Cigars	10.9	11.2	10.8	10.4	10.9	11.3
Pipe Tobacco <sup>4</sup>	--	--	--	--	--	--
Alcohol	68.8 <sup>a</sup>	70.3	71.5	71.2	68.5 <sup>b</sup>	70.2
Binge Alcohol Use <sup>5</sup>	--	--	--	--	--	--
Heavy Alcohol Use <sup>5</sup>	--	--	--	--	--	--
<b>PAST MONTH</b>						
Tobacco Products <sup>3</sup>	30.9	31.3	36.4	38.6	30.1	30.2
Cigarettes	26.4	26.5	33.3	35.8	25.4	25.1
Smokeless Tobacco	3.1	3.3	2.7	2.4	3.1	3.4
Cigars	5.8	5.8	5.1	5.2	5.9	5.9
Pipe Tobacco	0.8	0.9	0.9	1.1	0.8	0.9
Alcohol	54.1 <sup>b</sup>	55.9	56.3	55.7	53.8 <sup>b</sup>	56.0
Binge Alcohol Use <sup>5</sup>	24.1	24.1	23.9	22.3	24.2	24.4
Heavy Alcohol Use <sup>5</sup>	7.4	7.1	7.5	6.9	7.4	7.1

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.<sup>1</sup> Estimates in the Total column represent past year and past month use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.<sup>2</sup> Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.<sup>3</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.<sup>4</sup> Information about past year use of pipe tobacco was not collected.<sup>5</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.26A Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Numbers in Thousands, 2004 and 2005**

Dependence/Abuse	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
<b>DEPENDENCE OR ABUSE<sup>3</sup></b>						
Illicit Drugs or Alcohol <sup>4</sup>	20,278	20,180	4,549	4,257	15,684	15,863
Illicit Drugs <sup>4</sup>	5,951	5,647	1,737	1,497	4,194	4,115
Alcohol	17,137	17,251	3,630	3,465	13,475	13,733
<b>DEPENDENCE<sup>3</sup></b>						
Illicit Drugs or Alcohol <sup>4</sup>	10,719	10,290	2,970	2,678	7,732	7,583
Illicit Drugs <sup>4</sup>	4,115	4,040	1,339	1,170	2,770	2,847
Alcohol	7,821	7,424	2,112	1,917	5,699	5,495

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column represent past year dependence/abuse for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

<sup>3</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>4</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.26B Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Percentages, 2004 and 2005**

Dependence/Abuse	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
<b>DEPENDENCE OR ABUSE<sup>3</sup></b>						
Illicit Drugs or Alcohol <sup>4</sup>	9.4	9.3	16.5	15.1	8.4	8.4
Illicit Drugs <sup>4</sup>	2.8	2.6	6.3	5.3	2.2	2.2
Alcohol	8.0	7.9	13.2	12.3	7.2	7.3
<b>DEPENDENCE<sup>3</sup></b>						
Illicit Drugs or Alcohol <sup>4</sup>	5.0	4.7	10.8	9.5	4.1	4.0
Illicit Drugs <sup>4</sup>	1.9	1.9	4.9	4.2	1.5	1.5
Alcohol	3.6	3.4	7.7	6.8	3.0	2.9

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column represent past year dependence/abuse for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

<sup>3</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>4</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.