

Table 6.36A Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Demographic Characteristics: Numbers in Thousands, 2004 and 2005

Demographic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2004	2005	2004	2005	2004	2005	2004	2005
TOTAL	5,660	5,503	1,855	1,762	2,011	2,054	1,794	1,688
GENDER								
Male	2,582	2,581	933	879	948	949	702	753
Female	3,077	2,922	922	883	1,063	1,105	1,092 ^a	935
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	4,742	4,600	1,574	1,451	1,639	1,722	1,529	1,427
White	3,569	3,351	1,145 ^a	1,020	1,264	1,226	1,160	1,104
Black or African American	895	931	320	339	292	360	283	232
American Indian or Alaska Native	40	31	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	95	139	39	39	23	*	32	35
Two or More Races	126	129	49	40	39	53	38	36
Hispanic or Latino	917	903	280	311	372	332	265	261

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.36B Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Demographic Characteristics: Percentages, 2004 and 2005

Demographic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2004	2005	2004	2005	2004	2005	2004	2005
TOTAL	22.5	21.8	22.3	21.5	23.3	23.7	22.0 ^a	20.1
GENDER								
Male	20.1	20.0	22.1	21.3	21.4	21.5	16.8	17.3
Female	25.0	23.6	22.4	21.8	25.3	25.9	27.3 ^b	23.1
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	22.7	22.0	22.9	21.8	22.7	23.9	22.3 ^a	20.3
White	23.1 ^a	21.8	23.0	20.9	23.7	23.2	22.5	21.3
Black or African American	23.6	24.2	24.2	27.3	22.8	27.1	24.0 ^a	18.1
American Indian or Alaska Native	24.6	19.2	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	9.5	13.2	12.2	12.3	6.3	*	10.7	9.4
Two or More Races	28.7	32.6	28.6	26.6	29.5	38.0	28.0	34.1
Hispanic or Latino	21.8	20.7	19.1	20.3	26.0	22.8	20.1	18.9

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.37A Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Geographic or Socioeconomic Characteristics: Numbers in Thousands, 2004 and 2005

Geographic/Socioeconomic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2004	2005	2004	2005	2004	2005	2004	2005
TOTAL	5,660	5,503	1,855	1,762	2,011	2,054	1,794	1,688
GEOGRAPHIC REGION								
Northeast	1,111	1,064	358	326	395	394	358	345
Midwest	1,274	1,282	420	393	433	470	422	420
South	1,895	1,877	636	646	669	683	589	547
West	1,380	1,281	440	398	514	507	425	376
COUNTY TYPE								
Large Metro	3,000	3,064	994	958	1,083	1,161	923	944
Small Metro	1,800 ^b	1,527	597	520	627 ^a	534	577 ^a	473
250K - 1 Mil. Pop.	1,157 ^a	1,016	376	338	432	356	349	322
< 250K Pop.	643 ^a	511	221	182	195	177	228 ^b	152
Nonmetro	860	913	264	283	301	359	295	271
Urbanized	350	381	108	113	121	148	120	121
Less Urbanized	429	433	132	138	148	172	148	123
Completely Rural	82	99	23	33	32	39	*	27
FAMILY INCOME								
Less Than \$20,000	1,213	1,133	412	371	431	450	371	312
\$20,000 - \$49,999	1,994	1,845	657	614	718	685	619	546
\$50,000 - \$74,999	1,012	977	339	289	356	366	318	322
\$75,000 or More	1,440	1,548	447	488	506	553	487	508
GOVERNMENT ASSISTANCE¹								
Yes	1,326	1,394	482	461	451 ^a	549	393	384
No	4,334	4,109	1,373	1,301	1,560	1,505	1,401	1,304

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.37B Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Geographic or Socioeconomic Characteristics: Percentages, 2004 and 2005

Geographic/Socioeconomic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2004	2005	2004	2005	2004	2005	2004	2005
TOTAL	22.5	21.8	22.3	21.5	23.3	23.7	22.0 ^a	20.1
GEOGRAPHIC REGION								
Northeast	24.6	23.5	24.0	22.3	25.3	24.8	24.4	23.3
Midwest	22.5	22.7	22.3	21.7	22.7	24.8	22.4	21.6
South	21.1	20.8	21.2	22.1	21.8	22.4	20.3	17.9
West	23.1	21.2	22.4	19.9	24.5	23.6	22.2	19.7
COUNTY TYPE								
Large Metro	22.6	22.5	22.0	21.7	23.7	24.7	21.8	21.1
Small Metro	23.6 ^b	20.5	23.8	21.2	24.1	21.2	22.8 ^a	19.2
250K - 1 Mil. Pop.	22.7 ^a	20.0	23.1	20.2	24.1	20.6	20.9	19.2
< 250K Pop.	25.3 ^a	21.7	25.0	23.4	24.3	22.7	26.7 ^b	19.2
Nonmetro	20.4	21.6	20.1	21.6	20.3 ^a	24.7	20.8	18.6
Urbanized	20.7 ^a	24.2	21.1	23.0	20.0 ^a	27.4	21.0	22.0
Less Urbanized	20.0	20.2	19.3	20.8	20.0	23.3	20.8	16.7
Completely Rural	21.7	19.7	20.6	20.3	23.7	22.7	*	16.0
FAMILY INCOME								
Less Than \$20,000	26.4	25.3	26.0	24.8	27.3	29.0	25.8	21.7
\$20,000 - \$49,999	23.3	22.2	22.7	21.6	24.8	24.6	22.5	20.2
\$50,000 - \$74,999	20.9	20.6	21.7	20.2	20.9	21.2	20.2	20.4
\$75,000 or More	20.1	20.1	19.5	20.2	20.5	21.1	20.3	18.9
GOVERNMENT ASSISTANCE¹								
Yes	28.5	27.6	28.6	26.7	28.1	30.9	28.7	24.7
No	21.2	20.3	20.6	20.1	22.2	21.8	20.6	19.0

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.38A Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Demographic Characteristics: Numbers in Thousands, 2004 and 2005

Demographic Characteristic	Total		ILLCIT DRUG USE ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
TOTAL	5,660	5,503	1,775 ^a	1,602	3,885	3,901
AGE						
12-13	1,855	1,762	296	279	1,559	1,483
14-15	2,011	2,054	635	555	1,376	1,498
16-17	1,794	1,688	845	767	949	920
GENDER						
Male	2,582	2,581	675	672	1,907	1,909
Female	3,077	2,922	1,100 ^b	930	1,978	1,992
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	4,742	4,600	1,498 ^a	1,340	3,244	3,260
White	3,569	3,351	1,182 ^a	1,049	2,388	2,302
Black or African American	895	931	248	219	647	712
American Indian or Alaska Native	40	31	*	14	27	17
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	95	139	*	*	81	122
Two or More Races	126	129	*	33	84	95
Hispanic or Latino	917	903	277	262	641	641

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.38B Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Demographic Characteristics: Percentages, 2004 and 2005

Demographic Characteristic	Total		ILLCIT DRUG USE ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
TOTAL	22.5	21.8	33.6	32.0	19.6	19.3
AGE						
12-13	22.3	21.5	38.5	36.6	20.6	20.0
14-15	23.3	23.7	34.8	34.5	20.2	21.2
16-17	22.0 ^a	20.1	31.4	29.1	17.3	16.0
GENDER						
Male	20.1	20.0	25.6	26.4	18.7	18.4
Female	25.0	23.6	41.6 ^a	37.7	20.4	20.1
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	22.7	22.0	33.8	32.2	19.7	19.5
White	23.1 ^a	21.8	34.4	33.4	19.8	18.8
Black or African American	23.6	24.2	35.7 ^a	27.9	20.9	23.2
American Indian or Alaska Native	24.6	19.2	*	29.7	28.0 ^a	14.7
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	9.5	13.2	*	*	9.4	12.6
Two or More Races	28.7	32.6	*	39.1	25.6	30.8
Hispanic or Latino	21.8	20.7	32.9	30.9	19.0	18.2

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.39A Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Geographic or Socioeconomic Characteristics: Numbers in Thousands, 2004 and 2005

Geographic/Socioeconomic Characteristic	Total		ILLCIT DRUG USE ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
TOTAL	5,660	5,503	1,775 ^a	1,602	3,885	3,901
GEOGRAPHIC REGION						
Northeast	1,111	1,064	360	316	751	747
Midwest	1,274	1,282	403	386	871	896
South	1,895	1,877	564	518	1,331	1,359
West	1,380	1,281	448	382	932	899
COUNTY TYPE						
Large Metro	3,000	3,064	922	844	2,078	2,220
Small Metro	1,800 ^b	1,527	553 ^a	464	1,247 ^b	1,063
250K - 1 Mil. Pop.	1,157 ^a	1,016	327	317	830 ^a	699
< 250K Pop.	643 ^a	511	227 ^b	147	417	364
Nonmetro	860	913	300	294	560	619
Urbanized	350	381	110	124	240	257
Less Urbanized	429	433	158	128	271	305
Completely Rural	82	99	*	42	50	57
FAMILY INCOME						
Less Than \$20,000	1,213	1,133	368	329	845	804
\$20,000 - \$49,999	1,994	1,845	639 ^a	536	1,354	1,310
\$50,000 - \$74,999	1,012	977	343	285	669	692
\$75,000 or More	1,440	1,548	424	452	1,016	1,096
GOVERNMENT ASSISTANCE²						
Yes	1,326	1,394	411	409	915	984
No	4,334	4,109	1,364 ^a	1,193	2,969	2,917

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.39B Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Geographic or Socioeconomic Characteristics: Percentages, 2004 and 2005

Geographic/Socioeconomic Characteristic	Total		ILLCIT DRUG USE ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
TOTAL	22.5	21.8	33.6	32.0	19.6	19.3
GEOGRAPHIC REGION						
Northeast	24.6	23.5	36.6	36.1	21.2	20.5
Midwest	22.5	22.7	33.2	35.0	19.6	19.7
South	21.1	20.8	31.8	28.6	18.5	18.8
West	23.1	21.2	34.2	31.3	20.0	18.6
COUNTY TYPE						
Large Metro	22.6	22.5	34.3	32.0	19.6	20.2
Small Metro	23.6 ^b	20.5	33.2	30.7	20.9 ^b	18.0
250K - 1 Mil. Pop.	22.7 ^a	20.0	29.6	30.2	20.8 ^b	17.3
< 250K Pop.	25.3 ^a	21.7	40.0 ^a	31.9	21.1	19.3
Nonmetro	20.4	21.6	32.5	34.3	17.0	18.4
Urbanized	20.7 ^a	24.2	28.9 ^a	38.7	18.3	20.4
Less Urbanized	20.0	20.2	33.8	29.6	16.2	17.9
Completely Rural	21.7	19.7	*	40.1	16.5	14.3
FAMILY INCOME						
Less Than \$20,000	26.4	25.3	35.3	32.7	23.8	23.1
\$20,000 - \$49,999	23.3	22.2	34.7	33.0	20.2	19.5
\$50,000 - \$74,999	20.9	20.6	34.5	30.1	17.4	18.3
\$75,000 or More	20.1	20.1	30.3	31.5	17.6	17.4
GOVERNMENT ASSISTANCE²						
Yes	28.5	27.6	38.4	34.4	25.5	25.5
No	21.2	20.3	32.4	31.2	18.2	17.8

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.40A Source of Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Numbers in Thousands, 2004 and 2005

Source of Treatment/Counseling	Total		AGE GROUP					
			12-13		14-15		16-17	
	2004	2005	2004	2005	2004	2005	2004	2005
RECEIVED TREATMENT/COUNSELING FROM ANY OF FOLLOWING 10 SOURCES¹	5,660	5,503	1,855	1,762	2,011	2,054	1,794	1,688
Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor	2,523	2,573	761	743	923	968	839	861
School Counselor, School Psychologist, or Having Regular Meetings with a Teacher ²	2,619	2,509	899	886	931	901	789	722
Mental Health Clinic or Center	716	657	204	168	261	231	251	259
In-Home Therapist, Counselor, or Family Preservation Worker	762	731	241	227	303	308	218	196
Pediatrician or Other Family Doctor	840	810	263	239	284	319	292	253
Overnight or Longer Stay in Any Type of Hospital	515	529	149	170	190	164	177	195
Special Education Services While in a Regular Classroom or in a Special Classroom or Placement in a Special Program or Special School ²	1,015	976	347	297	357	366	312	313
Partial Day Hospital or Day Treatment Program	439	449	152	156	149	165	138	129
Overnight or Longer Stay in a Residential Treatment Center	299 ^a	229	96 ^a	57	106	89	98	83
Overnight or Longer Stay in Foster Care or in a Therapeutic Foster Care Home	158	143	43	36	65	46	50	61

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple sources; thus, these response categories are not mutually exclusive.

² Respondents who did not report their school enrollment status or who reported not being enrolled in school in the past 12 months were not asked about receipt of mental health treatment/counseling from this source.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.40B Source of Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Percentages, 2004 and 2005

Source of Treatment/Counseling	Total		AGE GROUP					
			12-13		14-15		16-17	
	2004	2005	2004	2005	2004	2005	2004	2005
RECEIVED TREATMENT/COUNSELING FROM ANY OF FOLLOWING 10 SOURCES¹	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor	45.0	47.1	41.7	42.6	46.1	47.6	47.1	51.3
School Counselor, School Psychologist, or Having Regular Meetings with a Teacher ²	46.7	46.0	49.2	50.8	46.6	44.2	44.2	43.1
Mental Health Clinic or Center	12.8	12.1	11.1	9.6	13.1	11.4	14.1	15.4
In-Home Therapist, Counselor, or Family Preservation Worker	13.6	13.4	13.3	13.1	15.2	15.1	12.2	11.7
Pediatrician or Other Family Doctor	15.0	14.9	14.4	13.8	14.3	15.7	16.4	15.0
Overnight or Longer Stay in Any Type of Hospital	9.1	9.7	8.1	9.7	9.5	8.0	9.9	11.6
Special Education Services While in a Regular Classroom or in a Special Classroom or Placement in a Special Program or Special School ²	18.2	18.0	19.1	17.2	17.9	18.1	17.5	18.7
Partial Day Hospital or Day Treatment Program	7.8	8.2	8.3	8.9	7.5	8.1	7.8	7.7
Overnight or Longer Stay in a Residential Treatment Center	5.3 ^a	4.2	5.2 ^a	3.3	5.3	4.3	5.5	5.0
Overnight or Longer Stay in Foster Care or in a Therapeutic Foster Care Home	2.8	2.6	2.3	2.0	3.2	2.2	2.8	3.6

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple sources; thus, these response categories are not mutually exclusive.

² Respondents who did not report their school enrollment status or who reported not being enrolled in school in the past 12 months were not asked about receipt of mental health treatment/counseling from this source.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.41A Reasons for Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Numbers in Thousands, 2005

Reason for Treatment/Counseling ¹	Total	AGE GROUP		
		12-13	14-15	16-17
RECEIVED TREATMENT/ COUNSELING FROM ANY OF NINE SOURCES	5,157	1,649	1,921	1,588
Thought about Killing Self or Tried to Kill Self	805	173	317	315
Felt Depressed	2,082	555	730	796
Felt Very Afraid or Tense	776	252	248	276
Breaking Rules or "Acting Out"	1,142	359	479	304
Eating Problems	370	116	116	137
Trouble Controlling Anger	617	232	247	138
Got into Physical Fights	255	103	109	44
Problems at Home/Family	1,085	326	395	364
Problems with Friends	701	266	251	185
Problems with Other Than Family/ Friends	467	176	163	127
Problems at School	1,045	314	426	306
Other Diagnosed Mental/Neurological Disorder ²	85	21	43	21
Some Other Reason ³	467	134	170	162

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. Youths who reported they only received treatment/counseling from special education services while in a regular classroom or in a special classroom, a special program, or in a special school were not asked for the reasons they received these services and were therefore excluded from this analysis.

NOTE: Due to questionnaire changes in 2005, these estimates are not comparable with those published in prior NSDUH reports.

¹ Respondents were asked the reasons for the last time they received mental health treatment/counseling from each reported source of treatment and could indicate multiple reasons for the last time they received treatment; thus, these response categories are not mutually exclusive. Respondents who did not indicate receiving mental health treatment/counseling for any of the reasons listed in this table were excluded from the reasons for receiving treatment.

² Respondent specified that he/she has been diagnosed with a mental/neurological disorder as a reason for having received mental health treatment/counseling. This reason is one of the most commonly reported other reasons for having received treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Reason for Receiving Mental Health Treatment/Counseling were classified as not having received treatment for Some Other Reason.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 6.41B Reasons for Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Percentages, 2005

Reason for Treatment/Counseling ¹	Total	AGE GROUP		
		12-13	14-15	16-17
RECEIVED TREATMENT/ COUNSELING FROM ANY OF NINE SOURCES	100.0	100.0	100.0	100.0
Thought about Killing Self or Tried to Kill Self	17.3	11.8	18.2	21.6
Felt Depressed	44.6	38.0	41.9	54.5
Felt Very Afraid or Tense	16.6	17.2	14.2	18.9
Breaking Rules or "Acting Out"	24.5	24.6	27.5	20.8
Eating Problems	7.9	8.0	6.7	9.4
Trouble Controlling Anger	13.2	15.9	14.2	9.4
Got into Physical Fights	5.5	7.0	6.2	3.0
Problems at Home/Family	23.3	22.3	22.7	24.9
Problems with Friends	15.0	18.2	14.4	12.7
Problems with Other Than Family/ Friends	10.0	12.1	9.4	8.7
Problems at School	22.4	21.5	24.4	20.9
Other Diagnosed Mental/Neurological Disorder ²	1.8	1.4	2.5	1.4
Some Other Reason ³	10.0	9.2	9.8	11.1

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. Youths who reported they only received treatment/counseling from special education services while in a regular classroom or in a special classroom, a special program, or in a special school were not asked for the reasons they received these services and were therefore excluded from this analysis.

NOTE: Due to questionnaire changes in 2005, these estimates are not comparable with those published in prior NSDUH reports.

¹ Respondents were asked the reasons for the last time they received mental health treatment/counseling from each reported source of treatment and could indicate multiple reasons for the last time they received treatment; thus, these response categories are not mutually exclusive. Respondents who did not indicate receiving mental health treatment/counseling for any of the reasons listed in this table were excluded from the reasons for receiving treatment.

² Respondent specified that he/she has been diagnosed with a mental/neurological disorder as a reason for having received mental health treatment/counseling. This reason is one of the most commonly reported other reasons for having received treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Reason for Receiving Mental Health Treatment/Counseling were classified as not having received treatment for Some Other Reason.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 6.42A Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Numbers in Thousands, 2004 and 2005

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS³	5,300 ^a	5,041	1,775 ^a	1,602	3,504	3,406
Marijuana and Hashish	3,656 ^a	3,383	1,178	1,085	2,462	2,276
Cocaine	411	423	161	157	246	263
Crack	63	54	30	24	30	29
Heroin	48	37	17	19	30	18
Hallucinogens	753	665	297	282	450	376
LSD	157	150	78	70	78	79
PCP	72	85	34	48	38	36
Ecstasy	310	256	123	119	186 ^a	133
Inhalants	1,168	1,144	449	431	718	709
Nonmedical Use of Psychotherapeutics ⁴	2,225	2,105	846	757	1,369	1,333
Pain Relievers	1,865	1,741	705	623	1,152	1,106
OxyContin [®]	194	187	85	86	107	98
Tranquilizers	532	497	221	211	310	281
Stimulants	496	503	211	233	280	266
Methamphetamine	163	170	81	69	77	100
Sedatives	133	118	67	57	66	58
ILLCIT DRUGS OTHER THAN MARIJUANA³	3,269	3,129	1,193	1,081	2,064	2,029

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent past year use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.42B Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Percentages, 2004 and 2005

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS³	21.0 ^a	19.9	31.4	29.1	18.0	17.2
Marijuana and Hashish	14.5 ^b	13.3	20.8	19.7	12.6 ^a	11.5
Cocaine	1.6	1.7	2.8	2.9	1.3	1.3
Crack	0.3	0.2	0.5	0.4	0.2	0.1
Heroin	0.2	0.1	0.3	0.3	0.2	0.1
Hallucinogens	3.0	2.6	5.3	5.1	2.3 ^a	1.9
LSD	0.6	0.6	1.4	1.3	0.4	0.4
PCP	0.3	0.3	0.6	0.9	0.2	0.2
Ecstasy	1.2	1.0	2.2	2.2	1.0 ^a	0.7
Inhalants	4.6	4.5	7.9	7.8	3.7	3.6
Nonmedical Use of Psychotherapeutics ⁴	8.8	8.3	14.9	13.8	7.0	6.7
Pain Relievers	7.4	6.9	12.5	11.3	5.9	5.6
OxyContin [®]	0.8	0.7	1.5	1.6	0.5	0.5
Tranquilizers	2.1	2.0	3.9	3.8	1.6	1.4
Stimulants	2.0	2.0	3.7	4.2	1.4	1.3
Methamphetamine	0.6	0.7	1.4	1.2	0.4	0.5
Sedatives	0.5	0.5	1.2	1.0	0.3	0.3
ILLCIT DRUGS OTHER THAN MARIJUANA³	13.0	12.3	21.1	19.6	10.6	10.3

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent past year use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.43A Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Numbers in Thousands, 2004 and 2005

Substance	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
PAST YEAR						
Tobacco Products ³	5,562	5,365	1,755	1,622	3,784	3,702
Cigarettes	4,645 ^a	4,387	1,563	1,419	3,059	2,941
Smokeless Tobacco	1,085	1,105	277	267	800	825
Cigars	2,568	2,485	810	746	1,751	1,721
Pipe Tobacco ⁴	--	--	--	--	--	--
Alcohol	8,537	8,454	2,307	2,186	6,209	6,233
Binge Alcohol Use ⁵	--	--	--	--	--	--
Heavy Alcohol Use ⁵	--	--	--	--	--	--
PAST MONTH						
Tobacco Products ³	3,635 ^b	3,324	1,207	1,091	2,407 ^a	2,205
Cigarettes	2,998 ^a	2,733	1,052	963	1,925 ^a	1,747
Smokeless Tobacco	571	523	138	110	430	406
Cigars	1,201 ^a	1,064	378	354	817 ^a	702
Pipe Tobacco	178	158	70	58	105	100
Alcohol	4,436 ^a	4,191	1,203	1,160	3,220	3,012
Binge Alcohol Use ⁵	2,794 ^b	2,513	767	749	2,020 ^b	1,751
Heavy Alcohol Use ⁵	671	611	178	204	489 ^a	404

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent past year and past month use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

⁴ Information about past year use of pipe tobacco was not collected.

⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.43B Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Percentages, 2004 and 2005

Substance	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
PAST YEAR						
Tobacco Products ³	22.1	21.2	31.0	29.5	19.4	18.7
Cigarettes	18.4 ^a	17.3	27.6	25.8	15.7	14.9
Smokeless Tobacco	4.3	4.4	4.9	4.8	4.1	4.2
Cigars	10.2	9.8	14.3	13.5	9.0	8.7
Pipe Tobacco ⁴	--	--	--	--	--	--
Alcohol	33.9	33.3	40.8	39.7	31.9	31.5
Binge Alcohol Use ⁵	--	--	--	--	--	--
Heavy Alcohol Use ⁵	--	--	--	--	--	--
PAST MONTH						
Tobacco Products ³	14.4 ^b	13.1	21.3	19.8	12.4 ^a	11.2
Cigarettes	11.9 ^b	10.8	18.6	17.5	9.9 ^a	8.8
Smokeless Tobacco	2.3	2.1	2.4	2.0	2.2	2.1
Cigars	4.8 ^a	4.2	6.7	6.4	4.2 ^a	3.6
Pipe Tobacco	0.7	0.6	1.2	1.1	0.5	0.5
Alcohol	17.6 ^a	16.5	21.3	21.1	16.5 ^a	15.2
Binge Alcohol Use ⁵	11.1 ^b	9.9	13.6	13.6	10.4 ^b	8.9
Heavy Alcohol Use ⁵	2.7	2.4	3.1	3.7	2.5 ^a	2.0

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.¹ Estimates in the Total column represent past year and past month use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.⁴ Information about past year use of pipe tobacco was not collected.⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.44A Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Numbers in Thousands, 2004 and 2005

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
DEPENDENCE OR ABUSE³						
Illicit Drugs or Alcohol ⁴	2,228 ^a	2,038	844	780	1,379	1,250
Illicit Drugs ⁴	1,347 ^a	1,186	555	512	788 ^a	670
Alcohol	1,517	1,407	545	524	966	875
DEPENDENCE³						
Illicit Drugs or Alcohol ⁴	1,075	1,036	450	432	621	598
Illicit Drugs ⁴	737	678	331	301	404	374
Alcohol	535	549	223	213	308	332

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.44B Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Percentages, 2004 and 2005

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
DEPENDENCE OR ABUSE³						
Illicit Drugs or Alcohol ⁴	8.8 ^a	8.0	14.9	14.2	7.1 ^a	6.3
Illicit Drugs ⁴	5.3 ^a	4.7	9.8	9.3	4.0 ^a	3.4
Alcohol	6.0	5.5	9.6	9.5	5.0	4.4
DEPENDENCE³						
Illicit Drugs or Alcohol ⁴	4.3	4.1	7.9	7.9	3.2	3.0
Illicit Drugs ⁴	2.9	2.7	5.8	5.5	2.1	1.9
Alcohol	2.1	2.2	3.9	3.9	1.6	1.7

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.