

**Table 6.3A Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Use of Illicit Drugs: Numbers in Thousands, 2004 and 2005**

Drug	ILLCIT DRUG USE			
	Yes		No	
	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>1</sup></b>	7,626 <sup>a</sup>	6,611	18,606	18,007
Marijuana and Hashish	5,417 <sup>a</sup>	4,661	20,815	19,957
Cocaine	1,657	1,361	24,576	23,257
Crack	607	512	25,625	24,106
Heroin	*	*	26,017	24,475
Hallucinogens	909	821	25,323	23,797
LSD	*	106	26,064	24,512
PCP	*	*	26,217	24,583
Ecstasy	436	459	25,797	24,159
Inhalants	393	312	25,839	24,306
Nonmedical Use of Psychotherapeutics <sup>2</sup>	3,869	3,494	22,364	21,124
Pain Relievers	2,779	2,654	23,453	21,964
OxyContin <sup>®</sup>	423	387	25,809	24,231
Tranquilizers	1,643	1,520	24,589	23,098
Stimulants	993	831	25,239	23,787
Methamphetamine	551	432	25,681	24,186
Sedatives	*	243	25,949	24,375
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	4,761	4,279	21,471	20,339

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.3B Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Use of Illicit Drugs: Percentages, 2004 and 2005**

Drug	ILLCIT DRUG USE			
	Yes		No	
	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>1</sup></b>	24.8 <sup>a</sup>	22.0	10.1	9.6
Marijuana and Hashish	23.5	21.2	10.8	10.2
Cocaine	31.5	26.7	11.7	10.9
Crack	46.4	38.6	12.0 <sup>a</sup>	11.1
Heroin	*	*	12.1 <sup>a</sup>	11.3
Hallucinogens	28.0	26.1	11.9 <sup>a</sup>	11.1
LSD	*	25.7	12.1 <sup>a</sup>	11.3
PCP	*	*	12.2 <sup>a</sup>	11.3
Ecstasy	26.2	26.9	12.1 <sup>a</sup>	11.2
Inhalants	34.1	29.9	12.1 <sup>a</sup>	11.2
Nonmedical Use of Psychotherapeutics <sup>2</sup>	31.9 <sup>a</sup>	26.7	11.0	10.3
Pain Relievers	30.3	26.3	11.4	10.6
OxyContin <sup>®</sup>	44.4	37.2	12.0 <sup>a</sup>	11.2
Tranquilizers	38.9	32.0	11.6	10.8
Stimulants	38.4	36.6	11.9 <sup>a</sup>	11.0
Methamphetamine	42.3	38.3	12.0 <sup>a</sup>	11.2
Sedatives	*	38.4	12.1 <sup>a</sup>	11.2
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	29.2 <sup>a</sup>	25.2	10.8	10.1

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.4A Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Numbers in Thousands, 2004 and 2005**

Drug	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2004	2005	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>2</sup></b>	30,793	29,999	7,626 <sup>a</sup>	6,611	23,167	23,388
Marijuana and Hashish	23,060	21,991	5,417 <sup>a</sup>	4,661	17,642	17,330
Cocaine	5,267	5,099	1,657	1,361	3,610	3,738
Crack	1,309	1,327	607	512	702	815
Heroin	393	343	215	143	178	200
Hallucinogens	3,244	3,144	909	821	2,335	2,323
LSD	476	413	168	106	308	307
PCP	109	79	16	35	93	44
Ecstasy	1,662	1,703	436	459	1,226	1,245
Inhalants	1,151	1,043	393	312	758	731
Nonmedical Use of Psychotherapeutics <sup>3</sup>	12,124	13,068	3,869	3,494	8,255 <sup>b</sup>	9,574
Pain Relievers	9,161	10,074	2,779	2,654	6,382 <sup>a</sup>	7,420
OxyContin <sup>®</sup>	953	1,039	423	387	530	652
Tranquilizers	4,219	4,752	1,643	1,520	2,576 <sup>a</sup>	3,232
Stimulants	2,582	2,268	993	831	1,589	1,437
Methamphetamine	1,302	1,127	551	432	751	695
Sedatives	600	632	283	243	317	389
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	16,303	16,980	4,761	4,279	11,542	12,701

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.4B Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Percentages, 2004 and 2005**

Drug	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2004	2005	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>2</sup></b>	14.3	13.8	29.1	26.9	12.3	12.1
Marijuana and Hashish	10.7	10.1	20.7	18.9	9.3	9.0
Cocaine	2.4	2.3	6.3	5.5	1.9	1.9
Crack	0.6	0.6	2.3	2.1	0.4	0.4
Heroin	0.2	0.2	0.8	0.6	0.1	0.1
Hallucinogens	1.5	1.4	3.5	3.3	1.2	1.2
LSD	0.2	0.2	0.6	0.4	0.2	0.2
PCP	0.1	0.0	0.1	0.1	0.0	0.0
Ecstasy	0.8	0.8	1.7	1.9	0.6	0.6
Inhalants	0.5	0.5	1.5	1.3	0.4	0.4
Nonmedical Use of Psychotherapeutics <sup>3</sup>	5.6	6.0	14.7	14.2	4.4 <sup>a</sup>	5.0
Pain Relievers	4.3	4.6	10.6	10.8	3.4 <sup>a</sup>	3.8
OxyContin <sup>®</sup>	0.4	0.5	1.6	1.6	0.3	0.3
Tranquilizers	2.0	2.2	6.3	6.2	1.4 <sup>a</sup>	1.7
Stimulants	1.2	1.0	3.8	3.4	0.8	0.7
Methamphetamine	0.6	0.5	2.1	1.8	0.4	0.4
Sedatives	0.3	0.3	1.1	1.0	0.2	0.2
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	7.6	7.8	18.2	17.4	6.1	6.6

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.5A Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year and Past Month Tobacco Product and Alcohol Use: Numbers in Thousands, 2004 and 2005**

Substance	TOBACCO PRODUCT OR ALCOHOL USE			
	Yes		No	
	2004	2005	2004	2005
<b>PAST YEAR</b>				
Tobacco Products <sup>1</sup>	13,223	12,978	13,009 <sup>a</sup>	11,640
Cigarettes	12,135	11,941	14,097 <sup>a</sup>	12,677
Smokeless Tobacco	1,134	1,175	25,098	23,443
Cigars	3,990	3,556	22,243	21,062
Pipe Tobacco <sup>2</sup>	--	--	--	--
Alcohol	19,064 <sup>a</sup>	17,541	7,168	7,077
Binge Alcohol Use <sup>3</sup>	--	--	--	--
Heavy Alcohol Use <sup>3</sup>	--	--	--	--
<b>PAST MONTH</b>				
Tobacco Products <sup>1</sup>	11,691	11,453	14,541	13,165
Cigarettes	10,845	10,545	15,388	14,072
Smokeless Tobacco	808	849	25,424	23,769
Cigars	1,814	1,823	24,418	22,795
Pipe Tobacco	209	291	26,023	24,327
Alcohol	14,814	13,555	11,419	11,063
Binge Alcohol Use <sup>3</sup>	7,965	7,113	18,267	17,505
Heavy Alcohol Use <sup>3</sup>	2,791	2,317	23,441	22,301

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.5B Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year and Past Month Tobacco Product and Alcohol Use: Percentages, 2004 and 2005**

Substance	TOBACCO PRODUCT OR ALCOHOL USE			
	Yes		No	
	2004	2005	2004	2005
<b>PAST YEAR</b>				
Tobacco Products <sup>1</sup>	16.8	16.3	9.5 <sup>a</sup>	8.4
Cigarettes	18.3	18.0	9.5 <sup>a</sup>	8.4
Smokeless Tobacco	12.8	12.6	12.2 <sup>a</sup>	11.2
Cigars	16.6	14.6	11.6	10.9
Pipe Tobacco <sup>2</sup>	--	--	--	--
Alcohol	12.8 <sup>b</sup>	11.5	10.7	10.9
Binge Alcohol Use <sup>3</sup>	--	--	--	--
Heavy Alcohol Use <sup>3</sup>	--	--	--	--
<b>PAST MONTH</b>				
Tobacco Products <sup>1</sup>	17.2	16.8	9.9 <sup>a</sup>	8.8
Cigarettes	18.7	18.2	9.8 <sup>a</sup>	8.8
Smokeless Tobacco	11.7	11.9	12.2 <sup>a</sup>	11.3
Cigars	14.5	14.5	12.0 <sup>a</sup>	11.1
Pipe Tobacco	12.4	14.3	12.2 <sup>a</sup>	11.3
Alcohol	12.6 <sup>b</sup>	11.1	11.7	11.5
Binge Alcohol Use <sup>3</sup>	15.3 <sup>a</sup>	13.5	11.2	10.6
Heavy Alcohol Use <sup>3</sup>	17.1	15.0	11.8	11.0

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.6A Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Numbers in Thousands, 2004 and 2005**

Substance	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2004	2005	2004	2005	2004	2005
<b>PAST YEAR</b>						
Tobacco Products <sup>2</sup>	78,642	79,593	13,223	12,978	65,419	66,616
Cigarettes	66,191	66,446	12,135	11,941	54,056	54,504
Smokeless Tobacco	8,895	9,338	1,134	1,175	7,760	8,162
Cigars	23,963	24,423	3,990	3,556	19,974	20,867
Pipe Tobacco <sup>3</sup>	--	--	--	--	--	--
Alcohol	148,431 <sup>b</sup>	153,172	19,064 <sup>a</sup>	17,541	129,366 <sup>a</sup>	135,631
Binge Alcohol Use <sup>4</sup>	--	--	--	--	--	--
Heavy Alcohol Use <sup>4</sup>	--	--	--	--	--	--
<b>PAST MONTH</b>						
Tobacco Products <sup>2</sup>	68,035	68,195	11,691	11,453	56,344	56,742
Cigarettes	58,092	57,799	10,845	10,545	47,247	47,254
Smokeless Tobacco	6,884	7,158	808	849	6,076	6,309
Cigars	12,531	12,576	1,814	1,823	10,717	10,753
Pipe Tobacco	1,683	2,032	209	291	1,474	1,740
Alcohol	117,758 <sup>a</sup>	121,837	14,814	13,555	102,945 <sup>a</sup>	108,282
Binge Alcohol Use <sup>4</sup>	52,046	52,577	7,965	7,113	44,080	45,464
Heavy Alcohol Use <sup>4</sup>	16,292	15,424	2,791	2,317	13,501	13,107

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>3</sup> Information about past year use of pipe tobacco was not collected.

<sup>4</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.6B Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Percentages, 2004 and 2005**

Substance	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2004	2005	2004	2005	2004	2005
<b>PAST YEAR</b>						
Tobacco Products <sup>2</sup>	36.5	36.5	50.4	52.7	34.6	34.5
Cigarettes	30.7	30.5	46.3	48.5	28.6	28.2
Smokeless Tobacco	4.1	4.3	4.3	4.8	4.1	4.2
Cigars	11.1	11.2	15.2	14.4	10.6	10.8
Pipe Tobacco <sup>3</sup>	--	--	--	--	--	--
Alcohol	68.9	70.3	72.7	71.3	68.4 <sup>a</sup>	70.2
Binge Alcohol Use <sup>4</sup>	--	--	--	--	--	--
Heavy Alcohol Use <sup>4</sup>	--	--	--	--	--	--
<b>PAST MONTH</b>						
Tobacco Products <sup>2</sup>	31.6	31.3	44.6	46.5	29.8	29.4
Cigarettes	27.0	26.5	41.3	42.8	25.0	24.5
Smokeless Tobacco	3.2	3.3	3.1	3.4	3.2	3.3
Cigars	5.8	5.8	6.9	7.4	5.7	5.6
Pipe Tobacco	0.8	0.9	0.8	1.2	0.8	0.9
Alcohol	54.7	55.9	56.5	55.1	54.4	56.0
Binge Alcohol Use <sup>4</sup>	24.2	24.1	30.4	28.9	23.3	23.5
Heavy Alcohol Use <sup>4</sup>	7.6	7.1	10.6	9.4	7.1	6.8

\*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>3</sup> Information about past year use of pipe tobacco was not collected.

<sup>4</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.