

Table 6.66A Severity of Role Impairment as Measured by the Sheehan Disability Scale (SDS) (I), and Mean Number of Days in the Past Year in which Unable to Carry Out Normal Activities Due to Depression, by Overall Severity of Role Impairment (II), among Persons Aged 12 to 17 with Past Year Major Depressive Episode (MDE): Numbers in Thousands, 2005

SDS Role Domains (I)	SEVERITY OF ROLE IMPAIRMENT					
	Total ¹	None	Mild	Moderate	Severe	Very Severe
Chores at Home	2,092	142	437	860	493	160
School or Work	2,101	96	420	747	657	180
Close Relationships with Family	2,114	70	291	683	773	296
Social Life	2,086	59	281	806	708	232
Overall ²	2,176	4	104	563	1,013	493
	OVERALL SEVERITY OF ROLE IMPAIRMENT ²					
	Total	None	Mild	Moderate	Severe	Very Severe
Mean Number of Days Unable to Carry Out Normal Activity in the Past Year (II)³	29.6	N/A	16.6	15.6	25.8	56.5

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: The Sheehan Disability Scale (SDS) measures the impact of a disorder on a person's life. The SDS asks youths to give 0 to 10 ratings of the amount of impairment caused by the disorder in each of four role domains: (1) chores at home, (2) school or work, (3) close relationships with family, and (4) social life.

¹ Respondents with unknown severity of SDS role impairment data were excluded.

² Overall role impairment is defined by the highest level of severity of role impairment reported across all four SDS role domains.

³ Respondents with unknown data on the mean number of days that they were unable to carry out normal activities in the past year were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 6.66B Severity of Role Impairment as Measured by the Sheehan Disability Scale (SDS) (I), and Mean Number of Days in the Past Year in which Unable to Carry Out Normal Activities Due to Depression, by Overall Severity of Role Impairment (II), among Persons Aged 12 to 17 with Past Year Major Depressive Episode (MDE): Percentages, 2005

SDS Role Domains (I)	SEVERITY OF ROLE IMPAIRMENT					
	Total ¹	None	Mild	Moderate	Severe	Very Severe
Chores at Home	100.0	6.8	20.9	41.1	23.6	7.6
School or Work	100.0	4.6	20.0	35.6	31.3	8.6
Close Relationships with Family	100.0	3.3	13.8	32.3	36.6	14.0
Social Life	100.0	2.8	13.5	38.7	33.9	11.1
Overall ²	100.0	0.2	4.8	25.9	46.6	22.6
	OVERALL SEVERITY OF ROLE IMPAIRMENT ²					
	Total	None	Mild	Moderate	Severe	Very Severe
Mean Number of Days Unable to Carry Out Normal Activity in the Past Year (II)³	29.6	N/A	16.6	15.6	25.8	56.5

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: The Sheehan Disability Scale (SDS) measures the impact of a disorder on a person's life. The SDS asks youths to give 0 to 10 ratings of the amount of impairment caused by the disorder in each of four role domains: (1) chores at home, (2) school or work, (3) close relationships with family, and (4) social life.

¹ Respondents with unknown severity of SDS role impairment data were excluded.

² Overall role impairment is defined by the highest level of severity of role impairment reported across all four SDS role domains.

³ Respondents with unknown data on the mean number of days that they were unable to carry out normal activities in the past year were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 6.67A Severity of Role Impairment as Measured by the Sheehan Disability Scale (SDS) (I), and Mean Number of Days in the Past Year in which Unable to Carry Out Normal Activities Due to Depression, by Overall Severity of Role Impairment (II), among Persons Aged 18 or Older with Past Year Major Depressive Episode (MDE): Numbers in Thousands, 2005

SDS Role Domains (I)	SEVERITY OF ROLE IMPAIRMENT					
	Total ¹	None	Mild	Moderate	Severe	Very Severe
Home Management	15,261	408	2,119	5,877	5,117	1,739
Work	14,108	742	3,478	4,883	3,607	1,397
Close Relationships with Others	15,112	603	2,648	5,546	4,914	1,400
Social Life	15,123	502	2,122	5,086	5,520	1,893
Overall ²	15,599	51	988	4,366	7,227	2,966
	OVERALL SEVERITY OF ROLE IMPAIRMENT ²					
	Total	None	Mild	Moderate	Severe	Very Severe
Mean Number of Days Unable to Carry Out Normal Activity in the Past Year (II)³	60.2	N/A	*	22.8	62.7	124.6

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: The Sheehan Disability Scale (SDS) measures the impact of a disorder on a person's life. The SDS asks adults to give 0 to 10 ratings of the amount of impairment caused by the disorder in each of four role domains: (1) home management, (2) work, (3) close relationships with others, and (4) social life.

¹ Respondents with unknown severity of SDS role impairment data were excluded.

² Overall role impairment is defined by the highest level of severity of role impairment reported across all four SDS role domains.

³ Respondents with unknown data on the mean number of days that they were unable to carry out normal activities in the past year were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 6.67B Severity of Role Impairment as Measured by the Sheehan Disability Scale (SDS) (I), and Mean Number of Days in the Past Year in which Unable to Carry Out Normal Activities Due to Depression, by Overall Severity of Role Impairment (II), among Persons Aged 18 or Older with Past Year Major Depressive Episode (MDE): Percentages, 2005

SDS Role Domains (I)	SEVERITY OF ROLE IMPAIRMENT					
	Total ¹	None	Mild	Moderate	Severe	Very Severe
Home Management	100.0	2.7	13.9	38.5	33.5	11.4
Work	100.0	5.3	24.7	34.6	25.6	9.9
Close Relationships with Others	100.0	4.0	17.5	36.7	32.5	9.3
Social Life	100.0	3.3	14.0	33.6	36.5	12.5
Overall ²	100.0	0.3	6.3	28.0	46.3	19.0
	OVERALL SEVERITY OF ROLE IMPAIRMENT ²					
	Total	None	Mild	Moderate	Severe	Very Severe
Mean Number of Days Unable to Carry Out Normal Activity in the Past Year (II)³	60.2	N/A	*	22.8	62.7	124.6

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: The Sheehan Disability Scale (SDS) measures the impact of a disorder on a person's life. The SDS asks adults to give 0 to 10 ratings of the amount of impairment caused by the disorder in each of four role domains: (1) home management, (2) work, (3) close relationships with others, and (4) social life.

¹ Respondents with unknown severity of SDS role impairment data were excluded.

² Overall role impairment is defined by the highest level of severity of role impairment reported across all four SDS role domains.

³ Respondents with unknown data on the mean number of days that they were unable to carry out normal activities in the past year were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.