

**Table 6.17C Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Psychological Distress Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2004 and 2005**

Location of Treatment/Counseling <sup>1</sup>	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Outpatient Mental Health Clinic or Center	213	178	27	32	144	108	*	133
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	282	201	54	39	228	137	*	147
Doctor's Office - Not Part of a Clinic	196	117	34	24	172	83	*	76
Outpatient Medical Clinic	186	84	18	16	102	54	*	62
Partial Day Hospital or Day Treatment Program	79	51	15	10	23	22	*	*
School or University Setting/Clinic/Center <sup>2</sup>	18	13	5	13	17	*	*	*
Some Other Place <sup>3</sup>	37	34	22	6	21	22	*	*

\*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

<sup>2</sup> Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

<sup>3</sup> Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.17D Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Psychological Distress Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2004 and 2005**

Location of Treatment/Counseling <sup>1</sup>	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Outpatient Mental Health Clinic or Center	2.57	2.03	2.45	2.16	2.75	2.39	*	5.43
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	2.97	1.96	3.43	2.22	3.36	2.39	*	5.26
Doctor's Office - Not Part of a Clinic	2.29	1.57	2.79	1.84	3.04	1.97	*	3.93
Outpatient Medical Clinic	2.24	1.15	1.69	1.24	2.07	1.32	*	3.17
Partial Day Hospital or Day Treatment Program	1.05	0.72	1.41	0.80	0.49	0.57	*	*
School or University Setting/Clinic/Center <sup>2</sup>	0.24	0.18	0.52	1.03	0.37	*	*	*
Some Other Place <sup>3</sup>	0.50	0.49	2.06	0.52	0.44	0.57	*	*

\*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

<sup>2</sup> Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

<sup>3</sup> Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.17P Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Psychological Distress Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2005 Versus 2004**

Location of Treatment/Counseling <sup>1</sup>	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Outpatient Mental Health Clinic or Center	0.0486		0.0068		0.0412		*	
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	0.2396		0.8993		0.7541		*	
Doctor's Office - Not Part of a Clinic	0.0686		0.0543		0.0417		*	
Outpatient Medical Clinic	0.1326		0.9070		0.5002		*	
Partial Day Hospital or Day Treatment Program	0.8967		0.7965		0.3633		*	
School or University Setting/Clinic/Center <sup>2</sup>	0.6128		0.0235		*		*	
Some Other Place <sup>3</sup>	0.9579		0.1620		0.2267		*	

\*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

<sup>2</sup> Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

<sup>3</sup> Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.