

**Table 6.18C Source of Payment for Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Psychological Distress Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2004 and 2005**

Source of Payment <sup>1</sup>	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Self or Family Member Living in Household	232	151	40	32	201	101	*	109
Family Member Not Living in Household	55	42	29	19	46	15	*	34
Private Health Insurance	299	169	35	35	222	112	*	121
Medicare	149	102	14	9	75	55	*	83
Medicaid	118	125	24	19	93	75	*	94
Rehabilitation Program	13	7	3	5	13	5	*	*
Employer	132	86	17	9	97	70	*	51
VA or Other Military Program	69	81	12	5	42	25	*	77
Other Public Source	80	53	21	14	62	43	*	29
Other Private Source	48	26	3	9	48	21	*	*
Free Treatment	61	54	23	19	46	39	*	*

\*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Respondents could indicate multiple sources of payment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.18D Source of Payment for Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Psychological Distress Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2004 and 2005**

Source of Payment <sup>1</sup>	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Self or Family Member Living in Household	2.57	1.83	3.10	2.14	3.32	2.20	*	5.03
Family Member Not Living in Household	0.75	0.60	2.51	1.47	1.01	0.39	*	1.76
Private Health Insurance	2.93	2.03	2.86	2.42	3.50	2.40	*	5.32
Medicare	1.95	1.36	1.33	0.75	1.57	1.34	*	4.02
Medicaid	1.53	1.63	2.23	1.51	1.88	1.85	*	4.48
Rehabilitation Program	0.18	0.10	0.26	0.39	0.28	0.13	*	*
Employer	1.70	1.19	1.60	0.73	2.01	1.69	*	2.61
VA or Other Military Program	0.93	1.13	1.10	0.45	0.90	0.65	*	3.78
Other Public Source	1.07	0.75	1.98	1.11	1.32	1.06	*	1.52
Other Private Source	0.65	0.38	0.30	0.77	1.03	0.53	*	*
Free Treatment	0.85	0.76	2.22	1.50	0.99	0.99	*	*

\*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Respondents could indicate multiple sources of payment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.18P Source of Payment for Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Psychological Distress Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Source of Payment <sup>1</sup>	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
Self or Family Member Living in Household	0.9727		0.5583		0.0964		*	
Family Member Not Living in Household	0.7002		0.9517		0.3394		*	
Private Health Insurance	0.0645		0.8924		0.0987		*	
Medicare	0.9569		0.8117		0.2552		*	
Medicaid	0.0156		0.8069		0.0147		*	
Rehabilitation Program	0.8619		0.2205		0.7620		*	
Employer	0.5664		0.0751		0.7868		*	
VA or Other Military Program	0.2248		0.1102		0.8911		*	
Other Public Source	0.3344		0.8217		0.1058		*	
Other Private Source	0.7081		0.0378		0.8592		*	
Free Treatment	0.3418		0.8171		0.3952		*	

\*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Respondents could indicate multiple sources of payment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.