

**Table 6.1C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Numbers in Thousands, 2004 and 2005**

Age Category	Total		GENDER			
			Male		Female	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	750	505	454	301	606	390
18	71	49	44	31	55	36
19	74	46	52	28	53	35
20	68	45	49	31	51	33
21	72	49	49	32	51	33
22	60	42	42	28	42	31
23	54	42	36	27	42	30
24	71	40	37	26	59	30
25	62	40	35	23	50	32
26-29	198	122	132	89	144	88
30-34	239	161	148	99	186	122
35-39	286	139	145	88	239	108
40-44	237	183	155	104	187	147
45-49	283	148	157	86	235	117
50-54	314	178	226	107	217	144
55-59	183	182	104	101	150	156
60-64	196	130	77	97	180	87
65 or Older	288	183	181	87	200	164

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.1D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Percentages, 2004 and 2005**

Age Category	Total		GENDER			
			Male		Female	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.35	0.23	0.44	0.29	0.54	0.35
18	1.34	0.94	1.66	1.15	2.05	1.52
19	1.54	0.90	2.27	1.09	2.38	1.40
20	1.53	0.94	2.10	1.35	2.24	1.37
21	1.57	1.02	2.32	1.40	2.18	1.43
22	1.36	0.89	1.95	1.20	1.97	1.35
23	1.32	0.98	1.77	1.32	1.91	1.35
24	1.52	0.93	1.72	1.28	2.36	1.40
25	1.47	0.94	1.66	1.14	2.41	1.47
26-29	1.21	0.75	1.67	1.07	1.76	1.11
30-34	1.13	0.78	1.42	1.00	1.66	1.12
35-39	1.27	0.66	1.39	0.86	1.91	1.04
40-44	0.97	0.73	1.32	0.90	1.54	1.10
45-49	1.23	0.65	1.41	0.79	1.99	0.97
50-54	1.53	0.89	2.23	1.09	2.05	1.39
55-59	1.09	1.03	1.29	1.22	1.76	1.61
60-64	1.52	0.96	1.32	1.40	2.59	1.30
65 or Older	0.83	0.52	1.23	0.58	1.00	0.81

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.1P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Age Category	Total		GENDER	
			Male	Female
	2004	2005	2004	2005
<b>TOTAL</b>	0.0348		0.0504	0.2467
18	0.1587		0.4757	0.3025
19	0.0082		0.0126	0.2591
20	0.1500		0.3149	0.2115
21	0.1507		0.1540	0.6755
22	0.9226		0.3559	0.4507
23	0.4635		0.7799	0.4261
24	0.2725		0.7020	0.1093
25	0.7733		0.9496	0.9602
26-29	0.2594		0.8360	0.1874
30-34	0.4952		0.9162	0.3892
35-39	0.0268		0.2024	0.0939
40-44	0.4974		0.9246	0.4711
45-49	0.0402		0.3195	0.0502
50-54	0.3756		0.0786	0.6657
55-59	0.0079		0.1760	0.0298
60-64	0.7740		0.2563	0.3372
65 or Older	0.7758		0.2279	0.5686

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.2C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics:  
Standard Errors of Numbers in Thousands, 2004 and 2005**

Demographic Characteristic	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	750	505	159	105	516	334	481	332
<b>GENDER</b>								
Male	454	301	113	70	319	207	310	193
Female	606	390	122	76	410	246	372	275
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	677	451	146	95	501	289	422	317
White	642	423	157	107	452	274	394	294
Black or African American	253	188	59	46	188	114	156	126
American Indian or Alaska Native	31	48	*	8	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	152	88	37	35	113	68	*	*
Two or More Races	79	53	*	13	47	31	*	42
Hispanic or Latino	330	217	77	60	224	163	197	110
<b>EDUCATION</b>								
< High School	372	259	93	54	244	148	255	198
High School Graduate	427	255	104	73	305	174	292	172
Some College	408	258	106	79	292	185	259	147
College Graduate	370	253	69	41	288	175	211	169
<b>CURRENT EMPLOYMENT</b>								
Full-Time	531	362	122	87	451	283	263	185
Part-Time	277	154	95	67	169	105	175	90
Unemployed	179	114	57	39	131	84	*	*
Other <sup>1</sup>	469	332	83	56	253	161	367	275

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.2D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics:  
Standard Errors of Percentages, 2004 and 2005**

Demographic Characteristic	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.35	0.23	0.50	0.32	0.52	0.33	0.58	0.39
<b>GENDER</b>								
Male	0.44	0.29	0.70	0.43	0.65	0.42	0.81	0.49
Female	0.54	0.35	0.76	0.47	0.81	0.49	0.83	0.60
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	0.36	0.24	0.55	0.36	0.59	0.34	0.55	0.40
White	0.40	0.26	0.63	0.42	0.65	0.40	0.59	0.42
Black or African American	1.02	0.69	1.28	0.89	1.49	0.86	1.99	1.47
American Indian or Alaska Native	3.07	3.78	*	3.42	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	1.75	0.95	2.35	1.95	2.41	1.31	*	*
Two or More Races	3.96	2.19	*	3.24	4.42	3.40	*	3.73
Hispanic or Latino	1.15	0.72	1.19	0.86	1.38	0.97	3.09	1.63
<b>EDUCATION</b>								
< High School	0.96	0.68	1.17	0.72	1.54	1.01	1.52	1.21
High School Graduate	0.61	0.37	0.82	0.56	0.97	0.56	1.01	0.59
Some College	0.74	0.44	0.88	0.59	1.06	0.65	1.47	0.76
College Graduate	0.62	0.42	1.38	0.86	0.88	0.54	0.95	0.71
<b>CURRENT EMPLOYMENT</b>								
Full-Time	0.42	0.29	0.70	0.48	0.57	0.38	0.81	0.55
Part-Time	0.92	0.51	1.01	0.68	1.51	0.88	1.75	0.95
Unemployed	2.11	1.36	1.95	1.33	3.24	2.10	*	*
Other <sup>1</sup>	0.76	0.50	1.20	0.74	1.73	1.09	0.89	0.63

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.2P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics: P**  
**Values from Tests of Differences of Percentages, 2005 Versus 2004**

Demographic Characteristic	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.0348		0.0073		0.0139		0.7360	
<b>GENDER</b>								
Male	0.0504		0.0269		0.2866		0.3518	
Female	0.2467		0.1306		0.0210		0.2424	
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	0.0285		0.0032		0.0091		0.5536	
White	0.0870		0.0082		0.0206		0.3852	
Black or African American	0.3280		0.3858		0.3183		0.8723	
American Indian or Alaska Native	0.0352		*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.3242		0.8501		0.6073		*	
Two or More Races	0.2153		*		0.9374		*	
Hispanic or Latino	0.7270		0.9801		0.9891		0.5829	
<b>EDUCATION</b>								
< High School	0.5638		0.0000		0.3030		0.2177	
High School Graduate	0.0230		0.9386		0.0061		0.5037	
Some College	0.1919		0.2989		0.7116		0.3342	
College Graduate	0.8275		0.5365		0.5720		0.1970	
<b>CURRENT EMPLOYMENT</b>								
Full-Time	0.0432		0.1299		0.0282		0.7439	
Part-Time	0.1639		0.4323		0.0609		0.8870	
Unemployed	0.5295		0.2712		0.7055		*	
Other <sup>1</sup>	0.7729		0.0548		0.4754		0.6080	

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.