

Table 6.24C Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2004 and 2005

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS³	519	541	230	225	483	503
Marijuana and Hashish	437	463	179	175	395	433
Cocaine	202	202	96	93	179	174
Crack	115	114	63	63	91	95
Heroin	55	42	29	26	45	35
Hallucinogens	147	140	59	62	137	124
LSD	41	45	18	15	35	40
PCP	25	15	8	10	24	11
Ecstasy	108	111	42	51	99	98
Inhalants	78	79	52	35	63	70
Nonmedical Use of Psychotherapeutics ⁴	341	341	162	155	292	301
Pain Relievers	289	300	130	127	246	273
OxyContin [®]	72	80	40	40	61	68
Tranquilizers	203	206	113	97	154	176
Stimulants	142	115	72	60	115	96
Methamphetamine	109	84	60	44	84	71
Sedatives	70	73	39	40	51	60
ILLCIT DRUGS OTHER THAN MARIJUANA³	399	393	184	178	348	351

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.24D Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2004 and 2005

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
ILLICIT DRUGS³	0.24	0.25	0.74	0.75	0.24	0.25
Marijuana and Hashish	0.20	0.21	0.61	0.60	0.20	0.22
Cocaine	0.09	0.09	0.34	0.33	0.09	0.09
Crack	0.05	0.05	0.23	0.22	0.05	0.05
Heroin	0.03	0.02	0.11	0.09	0.02	0.02
Hallucinogens	0.07	0.06	0.21	0.22	0.07	0.07
LSD	0.02	0.02	0.07	0.05	0.02	0.02
PCP	0.01	0.01	0.03	0.04	0.01	0.01
Ecstasy	0.05	0.05	0.15	0.18	0.05	0.05
Inhalants	0.04	0.04	0.19	0.13	0.03	0.04
Nonmedical Use of Psychotherapeutics ⁴	0.16	0.16	0.56	0.53	0.15	0.15
Pain Relievers	0.13	0.14	0.46	0.44	0.13	0.14
OxyContin [®]	0.03	0.04	0.15	0.14	0.03	0.04
Tranquilizers	0.09	0.09	0.40	0.34	0.08	0.09
Stimulants	0.07	0.05	0.26	0.21	0.06	0.05
Methamphetamine	0.05	0.04	0.22	0.15	0.05	0.04
Sedatives	0.03	0.03	0.14	0.14	0.03	0.03
ILLICIT DRUGS OTHER THAN MARIJUANA³	0.19	0.18	0.62	0.60	0.18	0.18

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.24P Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
ILLICIT DRUGS³	0.8516		0.0972		0.3819	
Marijuana and Hashish	0.9204		0.0752		0.5944	
Cocaine	0.4643		0.8937		0.2811	
Crack	0.6597		0.8785		0.6875	
Heroin	0.8639		0.8934		0.6535	
Hallucinogens	0.9300		0.7320		0.9676	
LSD	0.6668		0.8890		0.6956	
PCP	0.0387		0.5935		0.0123	
Ecstasy	0.6108		0.8402		0.6460	
Inhalants	0.6126		0.5782		0.7919	
Nonmedical Use of Psychotherapeutics ⁴	0.3012		0.3298		0.0903	
Pain Relievers	0.1727		0.3740		0.0548	
OxyContin [®]	0.9411		0.8750		0.9101	
Tranquilizers	0.5775		0.1030		0.1125	
Stimulants	0.3201		0.7117		0.3202	
Methamphetamine	0.2339		0.4679		0.3162	
Sedatives	0.8390		0.6081		0.3486	
ILLICIT DRUGS OTHER THAN MARIJUANA³	0.4834		0.3550		0.2460	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.25C Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
PAST YEAR						
Tobacco Products ³	816	878	365	373	908	939
Cigarettes	784	811	325	354	818	840
Smokeless Tobacco	260	289	107	85	263	277
Cigars	442	478	171	151	447	470
Pipe Tobacco ⁴	--	--	--	--	--	--
Alcohol	880	908	513	490	1,548	1,553
Binge Alcohol Use ⁵	--	--	--	--	--	--
Heavy Alcohol Use ⁵	--	--	--	--	--	--
PAST MONTH						
Tobacco Products ³	793	839	327	356	846	868
Cigarettes	765	772	301	342	777	774
Smokeless Tobacco	243	264	99	75	241	254
Cigars	323	351	103	97	321	340
Pipe Tobacco	145	145	65	52	131	138
Alcohol	950	969	445	439	1,398	1,373
Binge Alcohol Use ⁵	699	701	242	245	751	770
Heavy Alcohol Use ⁵	383	378	135	133	364	372

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year and past month use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.² Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.⁴ Information about past year use of pipe tobacco was not collected.⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.25D Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2004 and 2005

Substance	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
PAST YEAR						
Tobacco Products ³	0.38	0.40	1.03	1.03	0.40	0.43
Cigarettes	0.36	0.37	0.97	1.01	0.39	0.39
Smokeless Tobacco	0.12	0.13	0.38	0.30	0.14	0.14
Cigars	0.21	0.22	0.59	0.52	0.22	0.24
Pipe Tobacco ⁴	--	--	--	--	--	--
Alcohol	0.41	0.42	1.07	1.00	0.43	0.45
Binge Alcohol Use ⁵	--	--	--	--	--	--
Heavy Alcohol Use ⁵	--	--	--	--	--	--
PAST MONTH						
Tobacco Products ³	0.37	0.39	0.97	1.00	0.39	0.41
Cigarettes	0.36	0.35	0.93	0.98	0.38	0.37
Smokeless Tobacco	0.11	0.12	0.35	0.26	0.12	0.13
Cigars	0.15	0.16	0.37	0.34	0.16	0.18
Pipe Tobacco	0.07	0.07	0.23	0.18	0.07	0.07
Alcohol	0.44	0.44	1.09	1.09	0.46	0.48
Binge Alcohol Use ⁵	0.32	0.32	0.79	0.79	0.34	0.35
Heavy Alcohol Use ⁵	0.18	0.17	0.47	0.45	0.19	0.19

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year and past month use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

⁴ Information about past year use of pipe tobacco was not collected.

⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.25P Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
PAST YEAR						
Tobacco Products ³	0.3338		0.1579		0.6210	
Cigarettes	0.7220		0.0655		0.7226	
Smokeless Tobacco	0.1504		0.4738		0.0848	
Cigars	0.2591		0.6126		0.1461	
Pipe Tobacco ⁴	--		--		--	
Alcohol	0.0106		0.8339		0.0050	
Binge Alcohol Use ⁵	--		--		--	
Heavy Alcohol Use ⁵	--		--		--	
PAST MONTH						
Tobacco Products ³	0.5022		0.1190		0.9027	
Cigarettes	0.8377		0.0621		0.5951	
Smokeless Tobacco	0.1681		0.5063		0.0989	
Cigars	0.8356		0.9247		0.8594	
Pipe Tobacco	0.0864		0.4158		0.1292	
Alcohol	0.0039		0.6998		0.0012	
Binge Alcohol Use ⁵	0.9785		0.1582		0.6548	
Heavy Alcohol Use ⁵	0.1477		0.3664		0.2299	

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year and past month use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

⁴ Information about past year use of pipe tobacco was not collected.

⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.26C Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2004 and 2005

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
DEPENDENCE OR ABUSE³						
Illicit Drugs or Alcohol ⁴	432	421	202	205	391	391
Illicit Drugs ⁴	205	199	120	100	169	173
Alcohol	411	394	183	187	360	373
DEPENDENCE³						
Illicit Drugs or Alcohol ⁴	319	298	167	156	279	261
Illicit Drugs ⁴	164	169	106	86	132	145
Alcohol	296	260	147	137	248	227

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.26D Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2004 and 2005

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
DEPENDENCE OR ABUSE³						
Illicit Drugs or Alcohol ⁴	0.20	0.19	0.67	0.68	0.20	0.20
Illicit Drugs ⁴	0.10	0.09	0.42	0.35	0.09	0.09
Alcohol	0.19	0.18	0.63	0.63	0.19	0.19
DEPENDENCE³						
Illicit Drugs or Alcohol ⁴	0.15	0.14	0.58	0.54	0.14	0.14
Illicit Drugs ⁴	0.08	0.08	0.37	0.31	0.07	0.08
Alcohol	0.14	0.12	0.52	0.47	0.13	0.12

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.26P Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²	
			Received	Not Received
	2004	2005	2004	2005
DEPENDENCE OR ABUSE³				
Illicit Drugs or Alcohol ⁴	0.5766		0.1354	0.9568
Illicit Drugs ⁴	0.1926		0.0657	0.6194
Alcohol	0.8756		0.3099	0.7980
DEPENDENCE³				
Illicit Drugs or Alcohol ⁴	0.2059		0.1028	0.5512
Illicit Drugs ⁴	0.6009		0.1365	0.7981
Alcohol	0.2172		0.2129	0.4350

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.