

**Table 6.36C Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005**

Demographic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	93	92	60	59	64	66	63	52
<b>GENDER</b>								
Male	62	63	43	40	43	47	36	38
Female	66	65	42	41	46	49	48	40
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	83	81	55	50	53	59	57	46
White	86	82	44	41	47	45	50	43
Black or African American	51	49	28	26	23	33	27	22
American Indian or Alaska Native	6	5	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	17	25	12	10	8	*	10	9
Two or More Races	15	14	10	7	8	10	8	8
Hispanic or Latino	38	44	27	28	32	29	26	27

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.36D Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Demographic Characteristics: Standard Errors of Percentages, 2004 and 2005**

Demographic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.37	0.36	0.63	0.63	0.62	0.65	0.66	0.55
<b>GENDER</b>								
Male	0.49	0.49	0.90	0.85	0.84	0.93	0.81	0.81
Female	0.53	0.53	0.87	0.86	0.92	0.93	0.99	0.88
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	0.39	0.39	0.71	0.67	0.63	0.71	0.71	0.60
White	0.45	0.44	0.79	0.76	0.77	0.71	0.81	0.73
Black or African American	1.09	1.01	1.76	1.70	1.56	2.01	1.87	1.52
American Indian or Alaska Native	3.14	3.35	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	1.69	2.01	3.63	3.11	1.86	*	3.05	2.41
Two or More Races	2.48	2.84	4.86	3.97	4.73	5.05	5.13	5.59
Hispanic or Latino	0.91	1.01	1.55	1.63	1.95	1.71	1.68	1.71

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.36P Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Demographic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.1624		0.4135		0.6563		0.0295	
<b>GENDER</b>								
Male	0.8448		0.4987		0.9187		0.6926	
Female	0.0751		0.6106		0.6194		0.0016	
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	0.2432		0.2542		0.2344		0.0322	
White	0.0490		0.0601		0.6246		0.2693	
Black or African American	0.7227		0.1995		0.0871		0.0146	
American Indian or Alaska Native	0.2363		*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.1575		0.9683		*		0.7453	
Two or More Races	0.2925		0.7499		0.2182		0.4215	
Hispanic or Latino	0.4278		0.5999		0.2127		0.6211	

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.37C Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Geographic or Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005**

Geographic/Socioeconomic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	93	92	60	59	64	66	63	52
<b>GEOGRAPHIC REGION</b>								
Northeast	40	35	27	24	26	23	26	23
Midwest	37	37	24	20	21	27	25	23
South	52	57	36	37	35	41	39	33
West	53	50	32	34	41	40	35	28
<b>COUNTY TYPE</b>								
Large Metro	92	89	48	47	52	55	47	45
Small Metro	68	54	35	29	36	30	35	26
250K - 1 Mil. Pop.	54	46	28	23	29	26	26	21
< 250K Pop.	49	31	26	17	20	17	24	14
Nonmetro	38	42	18	22	21	23	24	20
Urbanized	24	28	12	15	12	15	15	14
Less Urbanized	29	29	14	15	17	16	17	13
Completely Rural	14	15	6	7	7	10	*	5
<b>FAMILY INCOME</b>								
Less Than \$20,000	51	52	26	26	32	32	29	25
\$20,000 - \$49,999	62	61	37	34	36	39	34	31
\$50,000 - \$74,999	44	44	26	22	26	28	25	25
\$75,000 or More	61	62	28	32	32	35	32	32
<b>GOVERNMENT ASSISTANCE<sup>1</sup></b>								
Yes	57	58	34	30	32	35	28	26
No	98	97	50	50	54	57	55	49

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>1</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.37D Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Geographic or Socioeconomic Characteristics: Standard Errors of Percentages, 2004 and 2005**

Geographic/Socioeconomic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.37	0.36	0.63	0.63	0.62	0.65	0.66	0.55
<b>GEOGRAPHIC REGION</b>								
Northeast	0.89	0.78	1.48	1.40	1.50	1.33	1.48	1.32
Midwest	0.66	0.65	1.14	0.98	1.01	1.21	1.16	1.07
South	0.58	0.63	1.08	1.11	0.95	1.16	1.10	1.01
West	0.89	0.83	1.41	1.50	1.58	1.46	1.57	1.28
<b>COUNTY TYPE</b>								
Large Metro	0.54	0.53	0.91	0.92	0.91	0.98	0.95	0.88
Small Metro	0.65	0.60	1.09	1.06	1.14	1.03	1.09	0.97
250K - 1 Mil. Pop.	0.77	0.74	1.32	1.28	1.41	1.27	1.26	1.15
< 250K Pop.	1.35	1.10	2.19	1.85	2.03	1.77	2.02	1.67
Nonmetro	0.73	0.77	1.20	1.35	1.20	1.37	1.46	1.18
Urbanized	1.16	1.33	2.19	2.37	1.76	2.24	2.16	2.16
Less Urbanized	1.02	1.02	1.71	1.75	1.84	1.72	1.99	1.55
Completely Rural	2.71	2.17	3.92	4.02	4.75	4.56	*	2.91
<b>FAMILY INCOME</b>								
Less Than \$20,000	0.91	0.92	1.36	1.51	1.70	1.73	1.64	1.44
\$20,000 - \$49,999	0.60	0.62	1.12	1.05	1.01	1.16	1.08	1.04
\$50,000 - \$74,999	0.76	0.79	1.50	1.33	1.33	1.42	1.40	1.43
\$75,000 or More	0.67	0.65	1.03	1.14	1.11	1.15	1.11	1.04
<b>GOVERNMENT ASSISTANCE<sup>1</sup></b>								
Yes	0.97	0.93	1.59	1.45	1.66	1.66	1.65	1.44
No	0.39	0.40	0.68	0.68	0.66	0.72	0.70	0.65

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>1</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.37P Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Geographic or Socioeconomic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Geographic/Socioeconomic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.1624		0.4135		0.6563		0.0295	
<b>GEOGRAPHIC REGION</b>								
Northeast	0.3602		0.3962		0.8161		0.5675	
Midwest	0.8105		0.6708		0.1831		0.6108	
South	0.7091		0.5548		0.6639		0.1053	
West	0.1119		0.2243		0.6814		0.2088	
<b>COUNTY TYPE</b>								
Large Metro	0.9565		0.7861		0.4931		0.5610	
Small Metro	0.0006		0.0915		0.0584		0.0124	
250K - 1 Mil. Pop.	0.0107		0.1138		0.0639		0.3228	
< 250K Pop.	0.0385		0.5616		0.5666		0.0042	
Nonmetro	0.2478		0.4126		0.0161		0.2419	
Urbanized	0.0480		0.5480		0.0102		0.7260	
Less Urbanized	0.8718		0.5349		0.1861		0.1070	
Completely Rural	0.5689		0.9690		0.8799		*	
<b>FAMILY INCOME</b>								
Less Than \$20,000	0.3800		0.5517		0.4894		0.0608	
\$20,000 - \$49,999	0.1806		0.4819		0.9007		0.1400	
\$50,000 - \$74,999	0.7959		0.4761		0.8904		0.9145	
\$75,000 or More	0.9547		0.6737		0.6878		0.3689	
<b>GOVERNMENT ASSISTANCE<sup>1</sup></b>								
Yes	0.5081		0.3819		0.2310		0.0681	
No	0.1457		0.5937		0.7126		0.1012	

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>1</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.38C Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005**

Demographic Characteristic	Total		ILLCIT DRUG USE <sup>1</sup>			
			Yes		No	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	93	92	59	55	91	94
<b>AGE</b>						
12-13	60	59	24	23	54	53
14-15	64	66	35	32	52	60
16-17	63	52	44	37	44	41
<b>GENDER</b>						
Male	62	63	34	36	63	62
Female	66	65	49	41	61	66
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	83	81	54	48	78	84
White	86	82	44	44	67	67
Black or African American	51	49	27	21	37	44
American Indian or Alaska Native	6	5	*	3	5	4
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	17	25	*	*	16	25
Two or More Races	15	14	*	6	13	13
Hispanic or Latino	38	44	25	27	43	42

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.38D Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Demographic Characteristics: Standard Errors of Percentages, 2004 and 2005**

Demographic Characteristic	Total		ILLCIT DRUG USE <sup>1</sup>			
			Yes		No	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.37	0.36	0.88	0.91	0.39	0.40
<b>AGE</b>						
12-13	0.63	0.63	2.46	2.48	0.64	0.64
14-15	0.62	0.65	1.55	1.66	0.65	0.73
16-17	0.66	0.55	1.30	1.20	0.72	0.65
<b>GENDER</b>						
Male	0.49	0.49	1.12	1.26	0.55	0.53
Female	0.53	0.53	1.31	1.29	0.55	0.58
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.39	0.39	0.93	0.98	0.42	0.43
White	0.45	0.44	0.99	1.11	0.49	0.48
Black or African American	1.09	1.01	2.75	2.28	1.05	1.14
American Indian or Alaska Native	3.14	3.35	*	4.88	4.12	3.76
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.69	2.01	*	*	1.85	2.14
Two or More Races	2.48	2.84	*	5.37	3.06	3.28
Hispanic or Latino	0.91	1.01	2.51	2.45	1.02	1.05

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.



**Table 6.38P Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Demographic Characteristic	Total		ILLCIT DRUG USE <sup>1</sup>			
			Yes		No	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.1624		0.1960		0.6007	
<b>AGE</b>						
12-13	0.4135		0.5741		0.4873	
14-15	0.6563		0.8927		0.2953	
16-17	0.0295		0.1937		0.1598	
<b>GENDER</b>						
Male	0.8448		0.6339		0.7034	
Female	0.0751		0.0348		0.7078	
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.2432		0.2540		0.7496	
White	0.0490		0.4912		0.1510	
Black or African American	0.7227		0.0284		0.1418	
American Indian or Alaska Native	0.2363		*		0.0176	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1575		*		0.2507	
Two or More Races	0.2925		*		0.2417	
Hispanic or Latino	0.4278		0.5622		0.6038	

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.39C Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Geographic or Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005**

Geographic/Socioeconomic Characteristic	Total		ILLCIT DRUG USE <sup>1</sup>			
			Yes		No	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	93	92	59	55	91	94
<b>GEOGRAPHIC REGION</b>						
Northeast	40	35	29	22	38	36
Midwest	37	37	23	23	35	35
South	52	57	33	32	52	58
West	53	50	32	32	53	53
<b>COUNTY TYPE</b>						
Large Metro	92	89	44	43	72	76
Small Metro	68	54	33	27	55	43
250K - 1 Mil. Pop.	54	46	24	23	44	37
< 250K Pop.	49	31	24	16	37	25
Nonmetro	38	42	24	25	27	33
Urbanized	24	28	13	17	19	22
Less Urbanized	29	29	20	14	19	23
Completely Rural	14	15	*	11	9	10
<b>FAMILY INCOME</b>						
Less Than \$20,000	51	52	27	25	40	44
\$20,000 - \$49,999	62	61	36	29	51	54
\$50,000 - \$74,999	44	44	25	24	35	38
\$75,000 or More	61	62	32	30	46	51
<b>GOVERNMENT ASSISTANCE<sup>2</sup></b>						
Yes	57	58	27	28	47	48
No	98	97	54	49	79	83

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.39D Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Geographic or Socioeconomic Characteristics: Standard Errors of Percentages, 2004 and 2005**

Geographic/Socioeconomic Characteristic	Total		ILLCIT DRUG USE <sup>1</sup>			
			Yes		No	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.37	0.36	0.88	0.91	0.39	0.40
<b>GEOGRAPHIC REGION</b>						
Northeast	0.89	0.78	2.19	2.01	0.92	0.85
Midwest	0.66	0.65	1.45	1.73	0.68	0.69
South	0.58	0.63	1.41	1.49	0.63	0.69
West	0.89	0.83	2.07	2.09	0.97	0.92
<b>COUNTY TYPE</b>						
Large Metro	0.54	0.53	1.29	1.37	0.54	0.58
Small Metro	0.65	0.60	1.53	1.41	0.72	0.63
250K - 1 Mil. Pop.	0.77	0.74	1.66	1.67	0.82	0.80
< 250K Pop.	1.35	1.10	2.90	2.72	1.50	1.09
Nonmetro	0.73	0.77	2.03	2.17	0.72	0.82
Urbanized	1.16	1.33	2.78	3.59	1.29	1.44
Less Urbanized	1.02	1.02	2.93	2.62	0.97	1.11
Completely Rural	2.71	2.17	*	5.85	2.32	2.28
<b>FAMILY INCOME</b>						
Less Than \$20,000	0.91	0.92	1.91	2.07	0.97	1.02
\$20,000 - \$49,999	0.60	0.62	1.57	1.45	0.68	0.71
\$50,000 - \$74,999	0.76	0.79	1.95	2.04	0.81	0.87
\$75,000 or More	0.67	0.65	1.78	1.74	0.66	0.67
<b>GOVERNMENT ASSISTANCE<sup>2</sup></b>						
Yes	0.97	0.93	1.78	1.90	1.11	1.00
No	0.39	0.40	1.04	1.06	0.42	0.44

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.39P Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Geographic or Socioeconomic Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Geographic/Socioeconomic Characteristic	Total		ILLCIT DRUG USE <sup>1</sup>			
			Yes		No	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.1624		0.1960		0.6007	
<b>GEOGRAPHIC REGION</b>						
Northeast	0.3602		0.8630		0.5442	
Midwest	0.8105		0.4288		0.8670	
South	0.7091		0.1169		0.7056	
West	0.1119		0.3314		0.3017	
<b>COUNTY TYPE</b>						
Large Metro	0.9565		0.2189		0.4113	
Small Metro	0.0006		0.2326		0.0019	
250K - 1 Mil. Pop.	0.0107		0.8274		0.0025	
< 250K Pop.	0.0385		0.0410		0.3142	
Nonmetro	0.2478		0.5398		0.2107	
Urbanized	0.0480		0.0306		0.2619	
Less Urbanized	0.8718		0.2848		0.2467	
Completely Rural	0.5689		*		0.4965	
<b>FAMILY INCOME</b>						
Less Than \$20,000	0.3800		0.3572		0.6337	
\$20,000 - \$49,999	0.1806		0.4385		0.4974	
\$50,000 - \$74,999	0.7959		0.1185		0.4654	
\$75,000 or More	0.9547		0.6279		0.8386	
<b>GOVERNMENT ASSISTANCE<sup>2</sup></b>						
Yes	0.5081		0.1177		0.9938	
No	0.1457		0.4352		0.4588	

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.40C Source of Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2004 and 2005**

Source of Treatment/Counseling	Total		AGE GROUP					
			12-13		14-15		16-17	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>RECEIVED TREATMENT/COUNSELING FROM ANY OF FOLLOWING 10 SOURCES<sup>1</sup></b>	116	110	60	59	64	66	63	52
Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor	73	71	37	36	44	42	43	39
School Counselor, School Psychologist, or Having Regular Meetings with a Teacher <sup>2</sup>	70	70	42	38	39	43	40	37
Mental Health Clinic or Center	40	35	20	17	25	21	21	21
In-Home Therapist, Counselor, or Family Preservation Worker	41	37	22	20	24	25	24	18
Pediatrician or Other Family Doctor	38	43	22	22	22	32	24	21
Overnight or Longer Stay in Any Type of Hospital	33	31	17	16	20	19	19	19
Special Education Services While in a Regular Classroom or in a Special Classroom or Placement in a Special Program or Special School <sup>2</sup>	46	43	25	24	25	27	27	24
Partial Day Hospital or Day Treatment Program	29	29	16	16	16	17	16	17
Overnight or Longer Stay in a Residential Treatment Center	24	22	13	9	15	16	13	12
Overnight or Longer Stay in Foster Care or in a Therapeutic Foster Care Home	18	17	9	8	12	10	10	11

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>1</sup> Respondents could indicate multiple sources; thus, these response categories are not mutually exclusive.

<sup>2</sup> Respondents who did not report their school enrollment status or who reported not being enrolled in school in the past 12 months were not asked about receipt of mental health treatment/counseling from this source.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.40D Source of Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2004 and 2005**

Source of Treatment/Counseling	Total		AGE GROUP					
			12-13		14-15		16-17	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>RECEIVED TREATMENT/COUNSELING FROM ANY OF FOLLOWING 10 SOURCES<sup>1</sup></b>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor	0.85	0.96	1.57	1.59	1.47	1.59	1.62	1.67
School Counselor, School Psychologist, or Having Regular Meetings with a Teacher <sup>2</sup>	0.85	0.94	1.55	1.55	1.46	1.68	1.65	1.67
Mental Health Clinic or Center	0.62	0.59	1.01	0.92	1.15	1.01	1.06	1.16
In-Home Therapist, Counselor, or Family Preservation Worker	0.67	0.62	1.13	1.07	1.09	1.13	1.26	1.06
Pediatrician or Other Family Doctor	0.61	0.71	1.14	1.18	1.01	1.39	1.21	1.18
Overnight or Longer Stay in Any Type of Hospital	0.54	0.53	0.90	0.86	0.92	0.90	0.99	1.08
Special Education Services While in a Regular Classroom or in a Special Classroom or Placement in a Special Program or Special School <sup>2</sup>	0.72	0.73	1.25	1.25	1.15	1.23	1.30	1.31
Partial Day Hospital or Day Treatment Program	0.48	0.51	0.84	0.89	0.78	0.82	0.84	0.96
Overnight or Longer Stay in a Residential Treatment Center	0.40	0.40	0.68	0.53	0.74	0.76	0.72	0.72
Overnight or Longer Stay in Foster Care or in a Therapeutic Foster Care Home	0.32	0.30	0.47	0.46	0.60	0.46	0.57	0.62

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>1</sup> Respondents could indicate multiple sources; thus, these response categories are not mutually exclusive.

<sup>2</sup> Respondents who did not report their school enrollment status or who reported not being enrolled in school in the past 12 months were not asked about receipt of mental health treatment/counseling from this source.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.40P Source of Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Source of Treatment/Counseling	Total		AGE GROUP					
			12-13		14-15		16-17	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>RECEIVED TREATMENT/COUNSELING FROM ANY OF FOLLOWING 10 SOURCES<sup>1</sup></b>	.		.		.		.	
Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor	0.0973		0.6753		0.5117		0.0730	
School Counselor, School Psychologist, or Having Regular Meetings with a Teacher <sup>2</sup>	0.5773		0.4689		0.2744		0.6595	
Mental Health Clinic or Center	0.4030		0.2691		0.2523		0.3918	
In-Home Therapist, Counselor, or Family Preservation Worker	0.8095		0.8955		0.9466		0.7445	
Pediatrician or Other Family Doctor	0.9221		0.7238		0.4143		0.4290	
Overnight or Longer Stay in Any Type of Hospital	0.4986		0.1938		0.2625		0.2418	
Special Education Services While in a Regular Classroom or in a Special Classroom or Placement in a Special Program or Special School <sup>2</sup>	0.8919		0.2749		0.8790		0.5021	
Partial Day Hospital or Day Treatment Program	0.5658		0.5851		0.5968		0.9470	
Overnight or Longer Stay in a Residential Treatment Center	0.0480		0.0249		0.3788		0.6238	
Overnight or Longer Stay in Foster Care or in a Therapeutic Foster Care Home	0.6748		0.6653		0.1888		0.3000	

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>1</sup> Respondents could indicate multiple sources; thus, these response categories are not mutually exclusive.

<sup>2</sup> Respondents who did not report their school enrollment status or who reported not being enrolled in school in the past 12 months were not asked about receipt of mental health treatment/counseling from this source.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.41C Reasons for Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2005**

Reason for Treatment/Counseling <sup>1</sup>	Total	AGE GROUP		
		12-13	14-15	16-17
<b>RECEIVED TREATMENT/ COUNSELING FROM ANY OF NINE SOURCES</b>	106	56	65	51
Thought about Killing Self or Tried to Kill Self	39	17	24	25
Felt Depressed	65	32	37	38
Felt Very Afraid or Tense	38	21	19	22
Breaking Rules or "Acting Out"	44	25	29	22
Eating Problems	25	14	14	15
Trouble Controlling Anger	35	22	24	14
Got into Physical Fights	23	14	15	8
Problems at Home/Family	44	23	26	25
Problems with Friends	37	22	24	17
Problems with Other Than Family/ Friends	30	17	18	14
Problems at School	47	23	32	24
Other Diagnosed Mental/Neurological Disorder <sup>2</sup>	12	6	8	5
Some Other Reason <sup>3</sup>	28	14	17	17

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. Youths who reported they only received treatment/counseling from special education services while in a regular classroom or in a special classroom, a special program, or in a special school were not asked for the reasons they received these services and were therefore excluded from this analysis.

NOTE: Due to questionnaire changes in 2005, these estimates are not comparable with those published in prior NSDUH reports.

<sup>1</sup> Respondents were asked the reasons for the last time they received mental health treatment/counseling from each reported source of treatment and could indicate multiple reasons for the last time they received treatment; thus, these response categories are not mutually exclusive. Respondents who did not indicate receiving mental health treatment/counseling for any of the reasons listed in this table were excluded from the reasons for receiving treatment.

<sup>2</sup> Respondent specified that he/she has been diagnosed with a mental/neurological disorder as a reason for having received mental health treatment/counseling. This reason is one of the most commonly reported other reasons for having received treatment/counseling.

<sup>3</sup> Respondents with unknown or invalid responses to the other-specify question on Some Other Reason for Receiving Mental Health Treatment/Counseling were classified as not having received treatment for Some Other Reason.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.



**Table 6.41D Reasons for Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2005**

Reason for Treatment/Counseling <sup>1</sup>	Total	AGE GROUP		
		12-13	14-15	16-17
<b>RECEIVED TREATMENT/ COUNSELING FROM ANY OF NINE SOURCES</b>	0.00	0.00	0.00	0.00
Thought about Killing Self or Tried to Kill Self	0.74	1.06	1.25	1.53
Felt Depressed	1.06	1.75	1.70	1.83
Felt Very Afraid or Tense	0.74	1.29	1.05	1.37
Breaking Rules or "Acting Out"	0.84	1.50	1.47	1.39
Eating Problems	0.52	0.93	0.78	1.01
Trouble Controlling Anger	0.70	1.38	1.26	0.96
Got into Physical Fights	0.47	0.94	0.83	0.55
Problems at Home/Family	0.81	1.43	1.30	1.50
Problems with Friends	0.72	1.32	1.27	1.12
Problems with Other Than Family/ Friends	0.61	1.09	0.99	0.96
Problems at School	0.85	1.41	1.50	1.46
Other Diagnosed Mental/Neurological Disorder <sup>2</sup>	0.24	0.41	0.44	0.33
Some Other Reason <sup>3</sup>	0.57	0.95	0.95	1.11

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. Youths who reported they only received treatment/counseling from special education services while in a regular classroom or in a special classroom, a special program, or in a special school were not asked for the reasons they received these services and were therefore excluded from this analysis.

NOTE: Due to questionnaire changes in 2005, these estimates are not comparable with those published in prior NSDUH reports.

<sup>1</sup> Respondents were asked the reasons for the last time they received mental health treatment/counseling from each reported source of treatment and could indicate multiple reasons for the last time they received treatment; thus, these response categories are not mutually exclusive. Respondents who did not indicate receiving mental health treatment/counseling for any of the reasons listed in this table were excluded from the reasons for receiving treatment.

<sup>2</sup> Respondent specified that he/she has been diagnosed with a mental/neurological disorder as a reason for having received mental health treatment/counseling. This reason is one of the most commonly reported other reasons for having received treatment/counseling.

<sup>3</sup> Respondents with unknown or invalid responses to the other-specify question on Some Other Reason for Receiving Mental Health Treatment/Counseling were classified as not having received treatment for Some Other Reason.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

**Table 6.42C Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2004 and 2005**

Drug	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>3</sup></b>	87	88	59	55	83	88
Marijuana and Hashish	77	75	48	46	68	69
Cocaine	27	28	18	18	21	22
Crack	10	10	8	7	7	7
Heroin	10	8	7	6	7	6
Hallucinogens	36	36	24	23	29	27
LSD	17	17	13	12	12	12
PCP	12	14	8	11	9	9
Ecstasy	23	21	15	15	19	15
Inhalants	45	47	29	28	38	38
Nonmedical Use of Psychotherapeutics <sup>4</sup>	59	61	40	35	48	51
Pain Relievers	56	56	37	32	46	46
OxyContin <sup>®</sup>	18	18	12	12	13	12
Tranquilizers	29	29	21	18	24	22
Stimulants	28	31	18	21	22	22
Methamphetamine	17	19	12	11	13	15
Sedatives	15	16	11	12	11	9
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>3</sup></b>	71	71	48	44	63	64

\*Low precision; no estimate reported.

<sup>1</sup> Estimates in the Total column represent past year use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>3</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>4</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.42D Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2004 and 2005**

Drug	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>3</sup></b>	0.34	0.35	0.81	0.86	0.38	0.38
Marijuana and Hashish	0.31	0.30	0.71	0.75	0.32	0.31
Cocaine	0.11	0.11	0.32	0.32	0.11	0.11
Crack	0.04	0.04	0.13	0.13	0.04	0.03
Heroin	0.04	0.03	0.12	0.10	0.04	0.03
Hallucinogens	0.14	0.14	0.40	0.41	0.14	0.13
LSD	0.07	0.07	0.22	0.21	0.06	0.06
PCP	0.05	0.06	0.15	0.19	0.05	0.05
Ecstasy	0.09	0.08	0.27	0.27	0.09	0.08
Inhalants	0.18	0.19	0.47	0.48	0.18	0.19
Nonmedical Use of Psychotherapeutics <sup>4</sup>	0.23	0.24	0.64	0.59	0.24	0.25
Pain Relievers	0.22	0.22	0.60	0.55	0.23	0.22
OxyContin <sup>®</sup>	0.07	0.07	0.22	0.22	0.07	0.06
Tranquilizers	0.12	0.11	0.36	0.33	0.12	0.11
Stimulants	0.11	0.12	0.31	0.38	0.11	0.11
Methamphetamine	0.07	0.07	0.21	0.20	0.07	0.08
Sedatives	0.06	0.06	0.20	0.22	0.05	0.05
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>3</sup></b>	0.28	0.28	0.74	0.72	0.29	0.29

\*Low precision; no estimate reported.

<sup>1</sup> Estimates in the Total column represent past year use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>3</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>4</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.42P Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling:  
P Values from Tests of Differences of Percentages, 2005 Versus 2004**

Drug	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>3</sup></b>	0.0202		0.0572		0.1638	
Marijuana and Hashish	0.0068		0.2853		0.0136	
Cocaine	0.7965		0.9872		0.6625	
Crack	0.4883		0.5827		0.9295	
Heroin	0.3627		0.7370		0.1565	
Hallucinogens	0.0707		0.8180		0.0401	
LSD	0.7374		0.7518		0.9734	
PCP	0.5166		0.2749		0.8519	
Ecstasy	0.0800		0.9800		0.0186	
Inhalants	0.6354		0.8657		0.7236	
Nonmedical Use of Psychotherapeutics <sup>4</sup>	0.1208		0.1740		0.4168	
Pain Relievers	0.0895		0.1668		0.3189	
OxyContin <sup>®</sup>	0.7539		0.8369		0.5636	
Tranquilizers	0.3483		0.9055		0.3000	
Stimulants	0.9143		0.3037		0.5729	
Methamphetamine	0.8029		0.5041		0.2591	
Sedatives	0.4779		0.6245		0.5116	
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>3</sup></b>	0.1172		0.1663		0.4357	

\*Low precision; no estimate reported.

<sup>1</sup> Estimates in the Total column represent past year use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>3</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>4</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.43C Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2004 and 2005**

Substance	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
<b>PAST YEAR</b>						
Tobacco Products <sup>3</sup>	96	99	59	53	88	93
Cigarettes	89	90	56	51	81	82
Smokeless Tobacco	45	46	22	22	40	40
Cigars	67	67	40	37	57	62
Pipe Tobacco <sup>4</sup>	--	--	--	--	--	--
Alcohol	103	105	67	63	118	125
Binge Alcohol Use <sup>5</sup>	--	--	--	--	--	--
Heavy Alcohol Use <sup>5</sup>	--	--	--	--	--	--
<b>PAST MONTH</b>						
Tobacco Products <sup>3</sup>	80	78	49	43	69	70
Cigarettes	76	71	46	40	65	62
Smokeless Tobacco	32	29	15	13	28	27
Cigars	44	46	27	26	38	37
Pipe Tobacco	19	17	11	12	16	12
Alcohol	81	81	47	46	81	80
Binge Alcohol Use <sup>5</sup>	73	65	41	35	65	58
Heavy Alcohol Use <sup>5</sup>	36	32	17	18	32	27

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Estimates in the Total column represent past year and past month use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.<sup>2</sup> Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.<sup>3</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.<sup>4</sup> Information about past year use of pipe tobacco was not collected.<sup>5</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.43D Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2004 and 2005**

Substance	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
<b>PAST YEAR</b>						
Tobacco Products <sup>3</sup>	0.38	0.39	0.86	0.85	0.40	0.42
Cigarettes	0.35	0.36	0.83	0.81	0.37	0.38
Smokeless Tobacco	0.18	0.18	0.39	0.40	0.20	0.20
Cigars	0.27	0.27	0.63	0.64	0.28	0.29
Pipe Tobacco <sup>4</sup>	--	--	--	--	--	--
Alcohol	0.41	0.41	0.83	0.89	0.47	0.47
Binge Alcohol Use <sup>5</sup>	--	--	--	--	--	--
Heavy Alcohol Use <sup>5</sup>	--	--	--	--	--	--
<b>PAST MONTH</b>						
Tobacco Products <sup>3</sup>	0.32	0.31	0.75	0.72	0.33	0.34
Cigarettes	0.30	0.28	0.72	0.68	0.31	0.31
Smokeless Tobacco	0.13	0.12	0.26	0.24	0.14	0.14
Cigars	0.18	0.18	0.45	0.45	0.19	0.19
Pipe Tobacco	0.08	0.07	0.20	0.22	0.08	0.06
Alcohol	0.32	0.32	0.72	0.76	0.37	0.36
Binge Alcohol Use <sup>5</sup>	0.29	0.26	0.67	0.61	0.32	0.28
Heavy Alcohol Use <sup>5</sup>	0.14	0.13	0.29	0.32	0.16	0.13

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Estimates in the Total column represent past year and past month use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>3</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>4</sup> Information about past year use of pipe tobacco was not collected.

<sup>5</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.43P Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Substance	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
<b>PAST YEAR</b>						
Tobacco Products <sup>3</sup>	0.1000		0.2087		0.2365	
Cigarettes	0.0259		0.1144		0.1282	
Smokeless Tobacco	0.8252		0.9269		0.8169	
Cigars	0.3034		0.4002		0.4944	
Pipe Tobacco <sup>4</sup>	--		--		--	
Alcohol	0.3763		0.3927		0.6275	
Binge Alcohol Use <sup>5</sup>	--		--		--	
Heavy Alcohol Use <sup>5</sup>	--		--		--	
<b>PAST MONTH</b>						
Tobacco Products <sup>3</sup>	0.0031		0.1502		0.0113	
Cigarettes	0.0069		0.2743		0.0180	
Smokeless Tobacco	0.2407		0.2177		0.4315	
Cigars	0.0242		0.6915		0.0167	
Pipe Tobacco	0.4228		0.5504		0.7670	
Alcohol	0.0190		0.8703		0.0121	
Binge Alcohol Use <sup>5</sup>	0.0027		0.9516		0.0004	
Heavy Alcohol Use <sup>5</sup>	0.1856		0.1925		0.0258	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Estimates in the Total column represent past year and past month use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>3</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>4</sup> Information about past year use of pipe tobacco was not collected.

<sup>5</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.44C Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2004 and 2005**

Dependence/Abuse	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
<b>DEPENDENCE OR ABUSE<sup>3</sup></b>						
Illicit Drugs or Alcohol <sup>4</sup>	61	60	39	40	52	50
Illicit Drugs <sup>4</sup>	46	47	30	32	37	35
Alcohol	50	51	32	31	43	43
<b>DEPENDENCE<sup>3</sup></b>						
Illicit Drugs or Alcohol <sup>4</sup>	43	45	29	28	34	35
Illicit Drugs <sup>4</sup>	36	37	24	23	26	28
Alcohol	31	35	21	21	24	27

\*Low precision; no estimate reported.

<sup>1</sup> Estimates in the Total column represent past year dependence/abuse for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>3</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>4</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.



**Table 6.44D Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2004 and 2005**

Dependence/Abuse	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2004	2005	2004	2005	2004	2005
<b>DEPENDENCE OR ABUSE<sup>3</sup></b>						
Illicit Drugs or Alcohol <sup>4</sup>	0.24	0.24	0.64	0.67	0.25	0.24
Illicit Drugs <sup>4</sup>	0.18	0.19	0.51	0.56	0.19	0.17
Alcohol	0.20	0.20	0.53	0.54	0.21	0.21
<b>DEPENDENCE<sup>3</sup></b>						
Illicit Drugs or Alcohol <sup>4</sup>	0.17	0.18	0.48	0.49	0.17	0.17
Illicit Drugs <sup>4</sup>	0.14	0.14	0.41	0.40	0.13	0.14
Alcohol	0.12	0.14	0.36	0.37	0.12	0.13

\*Low precision; no estimate reported.

<sup>1</sup> Estimates in the Total column represent past year dependence/abuse for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>3</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>4</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.44P Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Dependence/Abuse	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>	
			Received	Not Received
	2004	2005	2004	2005
<b>DEPENDENCE OR ABUSE<sup>3</sup></b>				
Illicit Drugs or Alcohol <sup>4</sup>	0.0187		0.4238	0.0305
Illicit Drugs <sup>4</sup>	0.0108		0.5126	0.0116
Alcohol	0.0993		0.8897	0.0712
<b>DEPENDENCE<sup>3</sup></b>				
Illicit Drugs or Alcohol <sup>4</sup>	0.4712		0.8953	0.5065
Illicit Drugs <sup>4</sup>	0.2245		0.5144	0.3582
Alcohol	0.8016		0.9063	0.5911

\*Low precision; no estimate reported.

<sup>1</sup> Estimates in the Total column represent past year dependence/abuse for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by drug or alcohol use. (See Table 6.40 for a list of the 10 sources.) Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

<sup>3</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>4</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.