

Table 6.3C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Use of Illicit Drugs: Standard Errors of Numbers in Thousands, 2004 and 2005

Drug	ILLCIT DRUG USE			
	Yes		No	
	2004	2005	2004	2005
ILLCIT DRUGS¹	371	229	668	463
Marijuana and Hashish	298	189	705	490
Cocaine	159	92	756	521
Crack	101	61	767	524
Heroin	*	*	773	530
Hallucinogens	86	52	772	527
LSD	*	17	776	529
PCP	*	*	778	530
Ecstasy	59	43	776	529
Inhalants	52	34	776	529
Nonmedical Use of Psychotherapeutics ²	247	162	731	493
Pain Relievers	205	142	751	499
OxyContin [®]	70	44	774	528
Tranquilizers	174	104	758	514
Stimulants	117	66	771	525
Methamphetamine	100	52	772	528
Sedatives	*	36	772	527
ILLCIT DRUGS OTHER THAN MARIJUANA¹	278	176	716	485

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.3D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Use of Illicit Drugs: Standard Errors of Percentages, 2004 and 2005

Drug	ILLCIT DRUG USE			
	Yes		No	
	2004	2005	2004	2005
ILLCIT DRUGS¹	1.03	0.66	0.35	0.24
Marijuana and Hashish	1.15	0.75	0.36	0.24
Cocaine	2.57	1.51	0.35	0.23
Crack	5.84	3.41	0.35	0.23
Heroin	*	*	0.35	0.23
Hallucinogens	2.49	1.53	0.35	0.23
LSD	*	3.89	0.35	0.23
PCP	*	*	0.35	0.23
Ecstasy	3.47	2.36	0.35	0.23
Inhalants	4.07	2.94	0.35	0.23
Nonmedical Use of Psychotherapeutics ²	1.73	1.03	0.35	0.23
Pain Relievers	1.91	1.20	0.35	0.23
OxyContin [®]	5.11	3.46	0.35	0.23
Tranquilizers	3.04	1.87	0.35	0.23
Stimulants	3.91	2.33	0.35	0.23
Methamphetamine	5.58	3.59	0.35	0.23
Sedatives	*	5.13	0.35	0.23
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1.47	0.87	0.35	0.23

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.3P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Use of Illicit Drugs: P Values from Tests of Differences of Percentages, 2005 Versus 2004

Drug	ILLCIT DRUG USE			
	Yes		No	
	2004	2005	2004	2005
ILLCIT DRUGS¹	0.0259		0.2394	
Marijuana and Hashish	0.0944		0.1367	
Cocaine	0.1114		0.0661	
Crack	0.2503		0.0433	
Heroin	*		0.0410	
Hallucinogens	0.5157		0.0425	
LSD	*		0.0404	
PCP	*		0.0326	
Ecstasy	0.8673		0.0333	
Inhalants	0.3979		0.0416	
Nonmedical Use of Psychotherapeutics ²	0.0104		0.0981	
Pain Relievers	0.0777		0.0552	
OxyContin [®]	0.2471		0.0386	
Tranquilizers	0.0514		0.0514	
Stimulants	0.6911		0.0472	
Methamphetamine	0.5488		0.0441	
Sedatives	*		0.0381	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0194		0.1132	

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.4C Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Numbers in Thousands, 2004 and 2005

Drug	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS²	734	541	371	229	674	497
Marijuana and Hashish	620	463	298	189	573	416
Cocaine	317	202	159	92	273	171
Crack	172	114	101	61	129	86
Heroin	88	42	64	27	61	35
Hallucinogens	207	140	86	52	188	125
LSD	68	45	33	17	60	40
PCP	30	15	6	10	29	11
Ecstasy	153	111	59	43	144	99
Inhalants	116	79	52	34	100	69
Nonmedical Use of Psychotherapeutics ³	465	341	247	162	401	298
Pain Relievers	390	300	205	142	343	263
OxyContin [®]	98	80	70	44	68	65
Tranquilizers	267	206	174	104	196	178
Stimulants	226	115	117	66	188	92
Methamphetamine	156	84	100	52	112	65
Sedatives	96	73	71	36	63	62
ILLCIT DRUGS OTHER THAN MARIJUANA²	560	393	278	176	500	346

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.4D Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Percentages, 2004 and 2005

Drug	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS²	0.34	0.25	1.22	0.82	0.34	0.25
Marijuana and Hashish	0.29	0.21	1.04	0.72	0.29	0.21
Cocaine	0.15	0.09	0.59	0.37	0.14	0.09
Crack	0.08	0.05	0.38	0.24	0.07	0.04
Heroin	0.04	0.02	0.24	0.11	0.03	0.02
Hallucinogens	0.10	0.06	0.33	0.22	0.10	0.06
LSD	0.03	0.02	0.13	0.07	0.03	0.02
PCP	0.01	0.01	0.02	0.04	0.02	0.01
Ecstasy	0.07	0.05	0.23	0.18	0.08	0.05
Inhalants	0.05	0.04	0.20	0.14	0.05	0.04
Nonmedical Use of Psychotherapeutics ³	0.22	0.16	0.89	0.61	0.21	0.15
Pain Relievers	0.18	0.14	0.76	0.54	0.18	0.13
OxyContin [®]	0.05	0.04	0.26	0.18	0.04	0.03
Tranquilizers	0.12	0.09	0.65	0.41	0.10	0.09
Stimulants	0.10	0.05	0.44	0.27	0.10	0.05
Methamphetamine	0.07	0.04	0.38	0.21	0.06	0.03
Sedatives	0.04	0.03	0.27	0.15	0.03	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.26	0.18	0.98	0.66	0.26	0.17

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.4P Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: P Values from Tests of Differences of Percentages, 2005 Versus 2004

Drug	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS²	0.2069		0.1322		0.7171	
Marijuana and Hashish	0.0854		0.1734		0.3151	
Cocaine	0.5444		0.2596		0.8819	
Crack	0.9905		0.6051		0.5359	
Heroin	0.5763		0.3646		0.7953	
Hallucinogens	0.5819		0.7440		0.7810	
LSD	0.4058		0.1461		0.9171	
PCP	0.3564		0.0894		0.1069	
Ecstasy	0.9076		0.4826		0.9634	
Inhalants	0.3911		0.3452		0.7255	
Nonmedical Use of Psychotherapeutics ³	0.1695		0.6075		0.0222	
Pain Relievers	0.1049		0.8413		0.0356	
OxyContin [®]	0.5616		0.8947		0.2475	
Tranquilizers	0.1554		0.9077		0.0247	
Stimulants	0.1786		0.4307		0.3829	
Methamphetamine	0.2884		0.4268		0.5863	
Sedatives	0.8400		0.7639		0.4701	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.4844		0.5163		0.1315	

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.5C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year and Past Month Tobacco Product and Alcohol Use: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	TOBACCO PRODUCT OR ALCOHOL USE			
	Yes		No	
	2004	2005	2004	2005
PAST YEAR				
Tobacco Products ¹	494	349	582	377
Cigarettes	471	330	601	393
Smokeless Tobacco	129	109	759	517
Cigars	267	144	717	505
Pipe Tobacco ²	--	--	--	--
Alcohol	634	408	447	303
Binge Alcohol Use ³	--	--	--	--
Heavy Alcohol Use ³	--	--	--	--
PAST MONTH				
Tobacco Products ¹	466	336	609	392
Cigarettes	448	317	621	408
Smokeless Tobacco	108	103	765	519
Cigars	144	110	764	514
Pipe Tobacco	44	49	774	527
Alcohol	563	359	542	367
Binge Alcohol Use ³	375	243	689	456
Heavy Alcohol Use ³	222	134	751	504

*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.5D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year and Past Month Tobacco Product and Alcohol Use: Standard Errors of Percentages, 2004 and 2005

Substance	TOBACCO PRODUCT OR ALCOHOL USE			
	Yes		No	
	2004	2005	2004	2005
PAST YEAR				
Tobacco Products ¹	0.59	0.41	0.41	0.26
Cigarettes	0.68	0.45	0.39	0.25
Smokeless Tobacco	1.37	1.09	0.36	0.24
Cigars	1.01	0.57	0.37	0.25
Pipe Tobacco ²	--	--	--	--
Alcohol	0.41	0.26	0.64	0.44
Binge Alcohol Use ³	--	--	--	--
Heavy Alcohol Use ³	--	--	--	--
PAST MONTH				
Tobacco Products ¹	0.65	0.45	0.40	0.25
Cigarettes	0.73	0.50	0.38	0.24
Smokeless Tobacco	1.55	1.32	0.36	0.23
Cigars	1.13	0.83	0.36	0.24
Pipe Tobacco	2.72	2.25	0.35	0.23
Alcohol	0.46	0.29	0.54	0.36
Binge Alcohol Use ³	0.66	0.42	0.41	0.27
Heavy Alcohol Use ³	1.23	0.79	0.37	0.24

*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.5P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year and Past Month Tobacco Product and Alcohol Use: P Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	TOBACCO PRODUCT OR ALCOHOL USE			
	Yes		No	
	2004	2005	2004	2005
PAST YEAR				
Tobacco Products ¹	0.4796		0.0240	
Cigarettes	0.6569		0.0182	
Smokeless Tobacco	0.9254		0.0329	
Cigars	0.0720		0.0986	
Pipe Tobacco ²	--		--	
Alcohol	0.0041		0.7770	
Binge Alcohol Use ³	--		--	
Heavy Alcohol Use ³	--		--	
PAST MONTH				
Tobacco Products ¹	0.6243		0.0218	
Cigarettes	0.6310		0.0262	
Smokeless Tobacco	0.9492		0.0319	
Cigars	0.9911		0.0317	
Pipe Tobacco	0.5908		0.0306	
Alcohol	0.0070		0.7760	
Binge Alcohol Use ³	0.0235		0.2234	
Heavy Alcohol Use ³	0.1501		0.0803	

*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.6C Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
PAST YEAR						
Tobacco Products ²	1,183	878	494	349	1,360	941
Cigarettes	1,111	811	471	330	1,238	829
Smokeless Tobacco	412	289	129	109	398	274
Cigars	655	478	267	144	628	476
Pipe Tobacco ³	--	--	--	--	--	--
Alcohol	1,306	908	634	408	2,267	1,598
Binge Alcohol Use ⁴	--	--	--	--	--	--
Heavy Alcohol Use ⁴	--	--	--	--	--	--
PAST MONTH						
Tobacco Products ²	1,126	839	466	336	1,259	873
Cigarettes	1,073	772	448	317	1,148	773
Smokeless Tobacco	387	264	108	103	383	248
Cigars	460	351	144	110	457	334
Pipe Tobacco	210	145	44	49	205	137
Alcohol	1,379	969	563	359	2,022	1,406
Binge Alcohol Use ⁴	1,048	701	375	243	1,081	749
Heavy Alcohol Use ⁴	593	378	222	134	552	362

*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

³ Information about past year use of pipe tobacco was not collected.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.6D Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Percentages, 2004 and 2005

Substance	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
PAST YEAR						
Tobacco Products ²	0.55	0.40	1.43	1.02	0.59	0.42
Cigarettes	0.52	0.37	1.41	1.00	0.55	0.38
Smokeless Tobacco	0.19	0.13	0.48	0.43	0.21	0.14
Cigars	0.30	0.22	0.94	0.57	0.32	0.24
Pipe Tobacco ³	--	--	--	--	--	--
Alcohol	0.61	0.42	1.40	0.96	0.65	0.45
Binge Alcohol Use ⁴	--	--	--	--	--	--
Heavy Alcohol Use ⁴	--	--	--	--	--	--
PAST MONTH						
Tobacco Products ²	0.52	0.39	1.41	1.01	0.56	0.40
Cigarettes	0.50	0.35	1.37	1.00	0.53	0.36
Smokeless Tobacco	0.18	0.12	0.41	0.41	0.20	0.12
Cigars	0.21	0.16	0.55	0.44	0.24	0.17
Pipe Tobacco	0.10	0.07	0.17	0.20	0.11	0.07
Alcohol	0.64	0.44	1.50	1.02	0.70	0.48
Binge Alcohol Use ⁴	0.49	0.32	1.29	0.86	0.52	0.34
Heavy Alcohol Use ⁴	0.28	0.17	0.82	0.52	0.28	0.18

*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

³ Information about past year use of pipe tobacco was not collected.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.6P Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹	
			Yes	No
	2004	2005	2004	2005
PAST YEAR				
Tobacco Products ²	0.9924		0.1889	0.8584
Cigarettes	0.7000		0.1937	0.5626
Smokeless Tobacco	0.5068		0.4861	0.6340
Cigars	0.8311		0.4863	0.5582
Pipe Tobacco ³	--		--	--
Alcohol	0.0640		0.4024	0.0261
Binge Alcohol Use ⁴	--		--	--
Heavy Alcohol Use ⁴	--		--	--
PAST MONTH				
Tobacco Products ²	0.6459		0.2589	0.5231
Cigarettes	0.4601		0.3786	0.3989
Smokeless Tobacco	0.6843		0.5204	0.8295
Cigars	0.8577		0.4871	0.7222
Pipe Tobacco	0.2028		0.1374	0.3484
Alcohol	0.1157		0.4372	0.0621
Binge Alcohol Use ⁴	0.9443		0.3413	0.7316
Heavy Alcohol Use ⁴	0.1348		0.2075	0.2876

*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

³ Information about past year use of pipe tobacco was not collected.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.