

**Table 6.45C Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 12 to 17, and Receipt of Treatment in Past Year for Depression among Persons Aged 12 to 17 with an MDE in Lifetime or Past Year, by Gender and Detailed Age Category: Standard Errors of Numbers in Thousands, 2004 and 2005**

Gender/Age Category	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	82	76	52	45	62	63	41	39
12	22	23	12	12	17	17	10	9
13	26	26	16	13	23	20	15	11
14	33	35	18	21	25	27	14	18
15	37	37	22	23	28	30	17	21
16	41	38	25	20	32	30	21	18
17	39	36	25	20	33	29	22	17
<b>MALE</b>	43	42	26	23	32	32	20	18
12	14	13	*	*	9	10	*	*
13	18	13	9	*	13	8	*	*
14	16	19	9	8	12	14	*	*
15	23	16	12	11	18	13	10	*
16	19	23	11	10	13	16	8	8
17	22	23	13	12	18	16	11	9
<b>FEMALE</b>	62	60	43	39	51	52	35	34
12	18	19	9	9	14	14	*	*
13	21	22	14	10	19	19	13	9
14	30	29	15	19	23	23	13	16
15	29	33	18	20	22	27	13	19
16	34	31	21	18	28	26	19	16
17	33	29	22	17	27	25	19	16

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.45D Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 12 to 17, and Receipt of Treatment in Past Year for Depression among Persons Aged 12 to 17 with an MDE in Lifetime or Past Year, by Gender and Detailed Age Category: Standard Errors of Percentages, 2004 and 2005**

Gender/Age Category	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.33	0.31	1.18	1.10	0.25	0.25	1.38	1.42
12	0.56	0.57	3.66	3.48	0.42	0.42	5.18	4.62
13	0.57	0.61	3.15	3.00	0.51	0.48	4.12	3.77
14	0.71	0.77	2.66	3.06	0.56	0.62	3.20	3.73
15	0.78	0.78	2.59	2.73	0.63	0.65	3.22	3.48
16	0.89	0.77	2.59	2.29	0.71	0.67	3.02	2.96
17	0.83	0.80	2.53	2.20	0.74	0.66	3.10	2.87
<b>MALE</b>	0.34	0.33	1.90	2.00	0.25	0.25	2.56	2.61
12	0.69	0.67	*	*	0.44	0.49	*	*
13	0.75	0.60	5.17	*	0.58	0.37	*	*
14	0.73	0.82	4.66	4.81	0.55	0.62	*	*
15	0.95	0.71	4.36	5.38	0.76	0.60	5.54	*
16	0.86	0.95	4.41	3.71	0.63	0.68	5.60	5.09
17	0.97	1.00	4.14	3.75	0.81	0.75	5.21	4.87
<b>FEMALE</b>	0.51	0.49	1.41	1.30	0.42	0.42	1.67	1.61
12	0.86	0.90	4.54	4.09	0.73	0.68	*	*
13	0.94	1.03	3.80	3.24	0.86	0.89	4.91	4.01
14	1.20	1.27	3.22	3.59	0.98	1.08	3.86	4.25
15	1.22	1.31	3.05	3.10	0.99	1.12	3.74	3.89
16	1.41	1.35	3.06	2.79	1.22	1.22	3.57	3.53
17	1.36	1.27	3.09	2.80	1.23	1.11	3.85	3.56

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.45P Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 12 to 17, and Receipt of Treatment in Past Year for Depression among Persons Aged 12 to 17 with an MDE in Lifetime or Past Year, by Gender and Detailed Age Category: P Values from Tests of Differences of Percentages, 2005 Versus 2004**

Gender/Age Category	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.5738		0.3676		0.6278		0.1981	
12	0.4268		0.6659		0.4459		0.5736	
13	0.1737		0.9126		0.2275		0.2888	
14	0.5815		0.1195		0.8409		0.2963	
15	0.8565		0.1518		0.6506		0.2239	
16	0.7991		0.0077		0.2884		0.0226	
17	0.8153		0.0786		0.6589		0.1432	
<b>MALE</b>	0.1655		0.6480		0.1205		0.3182	
12	0.7512		*		0.4604		*	
13	0.0248		*		0.0247		*	
14	0.4950		0.7518		0.9276		*	
15	0.0114		0.2104		0.0299		*	
16	0.3941		0.0824		0.7204		0.1995	
17	0.4241		0.3449		0.6145		0.1167	
<b>FEMALE</b>	0.7707		0.4026		0.6695		0.3227	
12	0.2116		0.4269		0.6775		*	
13	0.9846		0.5252		0.9354		0.3604	
14	0.9923		0.0529		0.7155		0.4591	
15	0.2506		0.3677		0.0798		0.3161	
16	0.5676		0.0425		0.2534		0.0698	
17	0.3251		0.1516		0.8189		0.4381	

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.46C Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 18 or Older, and Receipt of Treatment in Past Year for Depression among Persons Aged 18 or Older with an MDE in Lifetime or Past Year, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2004 and 2005**

Age Category	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	847	573	697	459	614	431	520	367
18	56	42	32	25	42	34	26	22
19	51	38	30	23	42	30	26	19
20	60	39	30	24	48	32	28	22
21	58	42	38	28	44	32	35	24
22	63	39	32	26	42	29	29	21
23	56	43	32	25	45	34	29	22
24	64	40	38	24	44	33	33	22
25	64	38	40	25	50	26	*	20
26-49	565	351	447	266	444	278	377	227
50 or Older	585	421	498	342	377	287	326	263

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older, while 2005 estimates are based on all respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.46D Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 18 or Older, and Receipt of Treatment in Past Year for Depression among Persons Aged 18 or Older with an MDE in Lifetime or Past Year, by Detailed Age Category: Standard Errors of Percentages, 2004 and 2005**

Age Category	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.40	0.26	1.46	0.97	0.29	0.20	1.75	1.24
18	1.10	0.85	4.01	2.98	0.85	0.71	5.00	3.89
19	1.24	0.81	4.13	2.95	1.06	0.67	5.31	3.67
20	1.37	0.88	3.67	2.83	1.12	0.73	4.87	3.54
21	1.32	0.96	4.68	3.29	1.05	0.75	5.47	4.28
22	1.43	0.85	4.02	3.18	1.02	0.66	5.65	3.78
23	1.33	0.98	3.93	2.98	1.11	0.83	5.34	3.98
24	1.45	0.95	4.10	2.89	1.03	0.81	5.17	4.03
25	1.50	0.90	4.62	2.97	1.22	0.66	*	3.92
26-49	0.57	0.35	1.77	1.22	0.45	0.28	2.28	1.65
50 or Older	0.71	0.50	3.79	2.23	0.46	0.34	4.33	2.74

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older, while 2005 estimates are based on all respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.46P Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 18 or Older, and Receipt of Treatment in Past Year for Depression among Persons Aged 18 or Older with an MDE in Lifetime or Past Year, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Age Category	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.2289		0.3696		0.0414		0.8275	
18	0.6445		0.7179		0.9306		0.6871	
19	0.3634		0.3316		0.3748		0.5677	
20	0.3651		0.1161		0.5313		0.1767	
21	0.6107		0.6595		0.6052		0.5731	
22	0.4551		0.0102		0.9813		0.2393	
23	0.6429		0.5531		0.8564		0.5455	
24	0.2512		0.7375		0.8926		0.8976	
25	0.2618		0.9847		0.4694		*	
26-49	0.0026		0.0810		0.0068		0.6847	
50 or Older	0.1175		0.7976		0.8871		0.6190	

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older, while 2005 estimates are based on all respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.47C Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 18 or Older, and Receipt of Treatment in the Past Year for Depression among Persons Aged 18 or Older with an MDE in Lifetime or Past Year, by Gender and Age Group: Standard Errors of Numbers in Thousands, 2004 and 2005**

Gender/Age	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	847	573	697	459	614	431	520	367
18-25	152	108	96	71	117	87	83	60
26-49	565	351	447	266	444	278	377	227
50 or Older	585	421	498	342	377	287	326	263
<b>MALE</b>	462	335	340	251	359	249	275	201
18-25	94	66	56	38	74	51	46	33
26-49	342	227	218	156	264	173	184	132
50 or Older	296	240	*	186	219	161	*	142
<b>FEMALE</b>	715	452	615	368	512	336	444	293
18-25	117	77	74	59	94	66	66	50
26-49	438	270	386	212	356	208	328	177
50 or Older	510	334	450	277	337	239	*	215

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older, while 2005 estimates are based on all respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.47D Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 18 or Older, and Receipt of Treatment in the Past Year for Depression among Persons Aged 18 or Older with an MDE in Lifetime or Past Year, by Gender and Age Group: Standard Errors of Percentages, 2004 and 2005**

Gender/Age	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.40	0.26	1.46	0.97	0.29	0.20	1.75	1.24
18-25	0.48	0.34	1.46	1.04	0.37	0.27	1.93	1.32
26-49	0.57	0.35	1.77	1.22	0.45	0.28	2.28	1.65
50 or Older	0.71	0.50	3.79	2.23	0.46	0.34	4.33	2.74
<b>MALE</b>	0.45	0.32	2.52	1.63	0.35	0.24	3.14	2.20
18-25	0.59	0.41	2.59	1.72	0.46	0.31	3.34	2.33
26-49	0.70	0.46	3.05	2.07	0.54	0.35	4.17	2.83
50 or Older	0.78	0.61	*	4.01	0.58	0.41	*	5.41
<b>FEMALE</b>	0.65	0.40	1.80	1.14	0.46	0.30	2.03	1.40
18-25	0.73	0.48	1.80	1.31	0.59	0.41	2.37	1.64
26-49	0.87	0.54	2.14	1.44	0.71	0.41	2.51	1.87
50 or Older	1.15	0.73	4.63	2.62	0.76	0.52	*	3.09

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older, while 2005 estimates are based on all respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.



**Table 6.47P Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 18 or Older, and Receipt of Treatment in the Past Year for Depression among Persons Aged 18 or Older with an MDE in Lifetime or Past Year, by Gender and Age Group: P Values from Tests of Differences of Percentages, 2005 Versus 2004**

Gender/Age	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.2289		0.3696		0.0414		0.8275	
18-25	0.1195		0.1011		0.3617		0.4694	
26-49	0.0026		0.0810		0.0068		0.6847	
50 or Older	0.1175		0.7976		0.8871		0.6190	
<b>MALE</b>	0.3320		0.4293		0.3588		0.9216	
18-25	0.2962		0.8287		0.4091		0.9249	
26-49	0.6384		0.5211		0.6041		0.7212	
50 or Older	0.1854		*		0.5861		*	
<b>FEMALE</b>	0.0356		0.8179		0.0657		0.7739	
18-25	0.2314		0.0276		0.6188		0.4454	
26-49	0.0000		0.2157		0.0024		0.6069	
50 or Older	0.3032		0.8168		0.6010		*	

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older, while 2005 estimates are based on all respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.48C Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 12 to 17, and Receipt of Treatment in the Past Year for Depression among Persons Aged 12 to 17 with an MDE in Lifetime or Past Year, by Hispanic Origin and Race: Standard Errors of Numbers in Thousands, 2004 and 2005**

Hispanic Origin and Race	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	82	76	52	45	62	63	41	39
Not Hispanic or Latino	71	69	47	42	56	56	38	36
White	67	65	40	35	53	53	33	30
Black or African American	35	38	16	23	26	27	13	19
American Indian or Alaska Native	4	3	*	*	3	3	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	21	16	*	*	17	12	*	*
Two or More Races	13	10	*	*	10	7	*	*
Hispanic or Latino	37	35	21	19	28	29	17	17

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.48D Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 12 to 17, and Receipt of Treatment in the Past Year for Depression among Persons Aged 12 to 17 with an MDE in Lifetime or Past Year, by Hispanic Origin and Race: Standard Errors of Percentages, 2004 and 2005**

Hispanic Origin and Race	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.33	0.31	1.18	1.10	0.25	0.25	1.38	1.42
Not Hispanic or Latino	0.34	0.33	1.27	1.23	0.27	0.27	1.48	1.59
White	0.39	0.38	1.37	1.36	0.31	0.32	1.62	1.74
Black or African American	0.83	0.86	2.81	3.68	0.63	0.64	3.60	4.67
American Indian or Alaska Native	2.80	1.90	*	*	2.03	1.44	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	2.01	1.40	*	*	1.64	1.17	*	*
Two or More Races	2.36	2.36	*	*	2.15	1.62	*	*
Hispanic or Latino	0.89	0.82	2.77	2.77	0.67	0.68	3.63	3.61

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.48P Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 12 to 17, and Receipt of Treatment in the Past Year for Depression among Persons Aged 12 to 17 with an MDE in Lifetime or Past Year, by Hispanic Origin and Race: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Hispanic Origin and Race	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.5738		0.3676		0.6278		0.1981	
Not Hispanic or Latino	0.8546		0.5862		0.5739		0.3581	
White	0.9595		0.0743		0.8264		0.0193	
Black or African American	0.6657		0.1568		0.9340		0.0778	
American Indian or Alaska Native	0.4258		*		0.4873		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.1125		*		0.2541		*	
Two or More Races	0.7466		*		0.6678		*	
Hispanic or Latino	0.3683		0.3147		0.9685		0.3251	

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.49C Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 18 or Older, and Receipt of Treatment in the Past Year for Depression among Persons Aged 18 or Older with an MDE in Lifetime or Past Year, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005**

Demographic Characteristic	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	847	573	697	459	614	431	520	367
<b>GENDER</b>								
Male	462	335	340	251	359	249	275	201
Female	715	452	615	368	512	336	444	293
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	799	530	669	440	569	392	489	348
White	830	536	618	405	531	351	434	313
Black or African American	266	195	194	147	197	158	163	129
American Indian or Alaska Native	31	33	*	*	30	21	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	165	100	*	*	119	62	*	*
Two or More Races	97	44	*	35	86	37	*	*
Hispanic or Latino	248	194	189	144	212	164	154	126
<b>EDUCATION</b>								
< High School	292	225	233	174	241	173	205	131
High School Graduate	451	300	315	227	335	226	252	194
Some College	492	326	360	228	367	224	300	185
College Graduate	570	349	450	268	315	233	283	207
<b>CURRENT EMPLOYMENT</b>								
Full-Time	721	435	516	299	468	281	357	224
Part-Time	319	199	247	150	207	142	186	118
Unemployed	156	129	108	105	105	108	*	95
Other <sup>3</sup>	464	370	390	293	360	277	330	247
<b>MARITAL STATUS</b>								
Married	658	425	477	317	437	291	381	251
Widowed	203	191	*	152	157	135	*	*
Divorced or Separated	412	287	305	212	308	204	247	171
Never Married	476	273	371	176	289	207	201	148

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older, while 2005 estimates are based on all respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

<sup>3</sup> The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.49D Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 18 or Older, and Receipt of Treatment in the Past Year for Depression among Persons Aged 18 or Older with an MDE in Lifetime or Past Year, by Demographic Characteristics: Standard Errors of Percentages, 2004 and 2005**

Demographic Characteristic	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.40	0.26	1.46	0.97	0.29	0.20	1.75	1.24
<b>GENDER</b>								
Male	0.45	0.32	2.52	1.63	0.35	0.24	3.14	2.20
Female	0.65	0.40	1.80	1.14	0.46	0.30	2.03	1.40
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	0.43	0.28	1.54	1.03	0.30	0.21	1.83	1.28
White	0.49	0.32	1.63	1.10	0.34	0.23	1.91	1.35
Black or African American	1.08	0.70	4.71	3.49	0.82	0.59	5.67	4.44
American Indian or Alaska Native	3.05	2.83	*	*	2.99	1.88	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	1.87	1.10	*	*	1.36	0.70	*	*
Two or More Races	4.21	2.05	*	6.05	4.18	1.68	*	*
Hispanic or Latino	0.94	0.70	4.65	3.14	0.80	0.60	5.97	4.00
<b>EDUCATION</b>								
< High School	0.79	0.60	3.43	2.67	0.66	0.48	3.90	3.05
High School Graduate	0.65	0.42	2.40	1.71	0.49	0.33	3.12	2.09
Some College	0.83	0.55	2.44	1.64	0.67	0.40	3.06	2.18
College Graduate	0.88	0.56	3.11	2.07	0.53	0.39	3.58	2.53
<b>CURRENT EMPLOYMENT</b>								
Full-Time	0.55	0.34	2.09	1.35	0.38	0.23	2.55	1.88
Part-Time	1.02	0.65	3.38	2.21	0.72	0.49	4.04	2.65
Unemployed	1.97	1.49	5.47	4.24	1.40	1.31	*	4.77
Other <sup>3</sup>	0.74	0.57	2.79	1.99	0.60	0.44	2.81	2.09
<b>MARITAL STATUS</b>								
Married	0.50	0.32	2.20	1.47	0.35	0.24	2.66	1.85
Widowed	1.62	1.33	*	5.54	1.27	0.96	*	*
Divorced or Separated	1.31	0.89	3.28	2.29	1.07	0.68	4.04	2.59
Never Married	0.80	0.46	2.56	1.37	0.52	0.37	2.66	1.79

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older, while 2005 estimates are based on all respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

<sup>3</sup> The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.49P Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 18 or Older, and Receipt of Treatment in the Past Year for Depression among Persons Aged 18 or Older with an MDE in Lifetime or Past Year, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2005 Versus 2004**

Demographic Characteristic	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.2289		0.3696		0.0414		0.8275	
<b>GENDER</b>								
Male	0.3320		0.4293		0.3588		0.9216	
Female	0.0356		0.8179		0.0657		0.7739	
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	0.1129		0.8026		0.0165		0.4227	
White	0.2913		0.8572		0.0429		0.2532	
Black or African American	0.3859		0.7832		0.5225		0.6455	
American Indian or Alaska Native	0.1662		*		0.7015		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.6056		*		0.3725		*	
Two or More Races	0.0188		*		0.0834		*	
Hispanic or Latino	0.3062		0.0402		0.6160		0.2927	
<b>EDUCATION</b>								
< High School	0.9779		0.7873		0.3829		0.4142	
High School Graduate	0.6902		0.7944		0.2997		0.0721	
Some College	0.0257		0.6118		0.0069		0.6754	
College Graduate	0.9180		0.2642		0.5274		0.4521	
<b>CURRENT EMPLOYMENT</b>								
Full-Time	0.0390		0.3160		0.0027		0.7943	
Part-Time	0.2197		0.6262		0.5092		0.7071	
Unemployed	0.5311		0.5854		0.1069		*	
Other <sup>3</sup>	0.2374		0.1516		0.9603		0.2919	
<b>MARITAL STATUS</b>								
Married	0.6076		0.2980		0.1693		0.9463	
Widowed	0.7468		*		0.4836		*	
Divorced or Separated	0.1587		0.8210		0.1892		0.3838	
Never Married	0.5151		0.5575		0.5204		0.9111	

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older, while 2005 estimates are based on all respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

<sup>3</sup> The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.50C Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 12 to 17, and Receipt of Treatment in the Past Year for Depression among Persons Aged 12 to 17 with an MDE in Lifetime or Past Year, by Geographic, Socioeconomic, and Health Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005**

Geographic/Socioeconomic/Health Characteristic	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	82	76	52	45	62	63	41	39
<b>GEOGRAPHIC REGION</b>								
Northeast	34	28	20	22	29	22	18	17
Midwest	33	29	20	19	28	24	16	16
South	47	46	30	26	36	37	26	23
West	47	46	32	27	30	39	21	24
<b>COUNTY TYPE</b>								
Large Metro	73	67	39	38	51	55	29	33
Small Metro	48	44	29	20	38	36	22	17
Nonmetro	31	36	19	19	25	27	16	17
<b>FAMILY INCOME</b>								
Less Than \$20,000	38	37	24	21	27	30	15	19
\$20,000 - \$49,999	51	50	31	25	40	38	26	21
\$50,000 - \$74,999	37	36	21	21	31	30	19	18
\$75,000 or More	49	46	25	25	37	34	22	22
<b>GOVERNMENT ASSISTANCE<sup>3</sup></b>								
Yes	40	42	26	25	27	33	18	21
No	85	74	45	39	64	59	37	34
<b>HEALTH INSURANCE<sup>4</sup></b>								
Private	70	68	39	36	55	51	32	31
Medicaid/CHIP <sup>5</sup>	42	46	27	25	34	36	22	21
Other <sup>6</sup>	20	19	10	*	15	16	7	*
No Coverage	24	26	12	12	16	19	8	11
<b>OVERALL HEALTH<sup>7</sup></b>								
Excellent	45	39	24	21	32	31	18	18
Very Good	58	54	31	31	45	44	27	28
Good	42	41	28	23	33	32	23	19
Fair/Poor	20	20	14	13	16	15	*	9

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

<sup>3</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

<sup>4</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>5</sup> CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

<sup>6</sup> Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

<sup>7</sup> Respondents with unknown health data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.



**Table 6.50D Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 12 to 17, and Receipt of Treatment in the Past Year for Depression among Persons Aged 12 to 17 with an MDE in Lifetime or Past Year, by Geographic, Socioeconomic, and Health Characteristics: Standard Errors of Percentages, 2004 and 2005**

Geographic/Socioeconomic/Health Characteristic	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.33	0.31	1.18	1.10	0.25	0.25	1.38	1.42
<b>GEOGRAPHIC REGION</b>								
Northeast	0.76	0.62	2.50	2.77	0.64	0.50	2.78	3.42
Midwest	0.58	0.52	1.87	2.19	0.50	0.43	2.21	2.69
South	0.53	0.51	1.94	1.80	0.41	0.42	2.52	2.46
West	0.80	0.76	3.03	2.40	0.51	0.65	3.27	3.14
<b>COUNTY TYPE</b>								
Large Metro	0.46	0.43	1.72	1.66	0.33	0.37	2.03	2.15
Small Metro	0.55	0.53	1.97	1.76	0.46	0.45	2.27	2.32
Nonmetro	0.66	0.72	2.54	2.44	0.57	0.56	3.08	3.11
<b>FAMILY INCOME</b>								
Less Than \$20,000	0.73	0.72	2.76	2.86	0.56	0.62	3.05	3.60
\$20,000 - \$49,999	0.54	0.55	2.01	1.79	0.45	0.43	2.49	2.34
\$50,000 - \$74,999	0.68	0.70	2.42	2.44	0.59	0.60	3.05	3.01
\$75,000 or More	0.60	0.52	2.03	2.05	0.47	0.41	2.61	2.84
<b>GOVERNMENT ASSISTANCE<sup>3</sup></b>								
Yes	0.75	0.74	2.66	2.77	0.55	0.60	3.05	3.47
No	0.37	0.33	1.30	1.20	0.28	0.27	1.57	1.55
<b>HEALTH INSURANCE<sup>4</sup></b>								
Private	0.38	0.36	1.38	1.29	0.30	0.28	1.66	1.73
Medicaid/CHIP <sup>5</sup>	0.57	0.65	2.41	2.40	0.49	0.53	2.85	2.99
Other <sup>6</sup>	1.63	1.69	4.68	*	1.30	1.49	5.74	*
No Coverage	1.09	1.20	3.91	3.68	0.77	0.92	4.87	4.88
<b>OVERALL HEALTH<sup>7</sup></b>								
Excellent	0.48	0.44	2.24	2.19	0.36	0.35	2.73	2.98
Very Good	0.50	0.48	1.71	1.78	0.40	0.41	2.10	2.35
Good	0.73	0.69	2.34	2.12	0.60	0.56	2.84	2.56
Fair/Poor	1.98	1.82	4.69	4.48	1.71	1.45	*	5.38

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

<sup>3</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

<sup>4</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>5</sup> CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

<sup>6</sup> Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

<sup>7</sup> Respondents with unknown health data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.50P Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 12 to 17, and Receipt of Treatment in the Past Year for Depression among Persons Aged 12 to 17 with an MDE in Lifetime or Past Year, by Geographic, Socioeconomic, and Health Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Geographic/Socioeconomic/Health Characteristic	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.5738		0.3676		0.6278		0.1981	
<b>GEOGRAPHIC REGION</b>								
Northeast	0.0839		0.3160		0.0622		0.6879	
Midwest	0.5829		0.8274		0.7008		0.6480	
South	0.7087		0.1874		0.7724		0.2208	
West	0.8541		0.5939		0.6306		0.4892	
<b>COUNTY TYPE</b>								
Large Metro	0.6148		0.7958		0.7992		0.7484	
Small Metro	0.1737		0.0162		0.0636		0.0302	
Nonmetro	0.1606		0.9151		0.3743		0.9611	
<b>FAMILY INCOME</b>								
Less Than \$20,000	0.4274		0.4874		0.8362		0.2426	
\$20,000 - \$49,999	0.7732		0.0248		0.9456		0.0283	
\$50,000 - \$74,999	0.6325		0.7379		0.6418		0.1733	
\$75,000 or More	0.3065		0.6216		0.1990		0.7980	
<b>GOVERNMENT ASSISTANCE<sup>3</sup></b>								
Yes	0.5371		0.7937		0.9842		0.8980	
No	0.7162		0.3635		0.5718		0.1592	
<b>HEALTH INSURANCE<sup>4</sup></b>								
Private	0.5027		0.3976		0.2210		0.2223	
Medicaid/CHIP <sup>5</sup>	0.6296		0.8464		0.9113		0.7594	
Other <sup>6</sup>	0.6002		*		0.9828		*	
No Coverage	0.3669		0.7461		0.1045		0.4959	
<b>OVERALL HEALTH<sup>7</sup></b>								
Excellent	0.5366		0.8927		0.6719		0.7731	
Very Good	0.6987		0.5988		0.5830		0.6207	
Good	0.6100		0.3924		0.9545		0.1107	
Fair/Poor	0.7725		0.2969		0.5829		*	

\*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

<sup>3</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

<sup>4</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>5</sup> CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

<sup>6</sup> Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

<sup>7</sup> Respondents with unknown health data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.51C Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 18 or Older, and Receipt of Treatment in the Past Year for Depression among Persons Aged 18 or Older with an MDE in Lifetime or Past Year, by Geographic, Socioeconomic, and Health Characteristics: Standard Errors of Numbers in Thousands, 2004 and 2005**

Geographic/Socioeconomic/Health Characteristic	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	847	573	697	459	614	431	520	367
<b>GEOGRAPHIC REGION</b>								
Northeast	316	251	268	212	278	187	225	156
Midwest	307	248	217	199	214	185	181	170
South	585	347	496	284	376	259	317	236
West	426	287	346	206	336	206	294	162
<b>COUNTY TYPE</b>								
Large Metro	746	474	559	329	489	333	389	264
Small Metro	483	362	368	257	341	227	293	192
Nonmetro	366	240	245	188	266	172	216	157
<b>FAMILY INCOME</b>								
Less Than \$20,000	388	300	301	227	312	243	259	198
\$20,000 - \$49,999	466	373	348	279	346	261	271	223
\$50,000 - \$74,999	377	259	307	185	294	176	259	145
\$75,000 or More	577	298	427	206	310	188	266	157
<b>GOVERNMENT ASSISTANCE<sup>3</sup></b>								
Yes	364	286	291	232	305	233	246	197
No	853	548	630	396	554	367	459	308
<b>HEALTH INSURANCE<sup>4</sup></b>								
Private	797	495	622	357	515	306	450	256
Medicaid/CHIP <sup>5</sup>	255	231	208	196	230	204	199	180
Other <sup>6</sup>	365	324	283	266	230	227	213	214
No Coverage	338	218	213	131	265	162	177	109
<b>OVERALL HEALTH<sup>7</sup></b>								
Excellent	420	245	308	148	270	141	245	104
Very Good	494	329	326	232	335	209	259	168
Good	530	302	437	218	335	214	272	178
Fair/Poor	406	318	318	279	348	269	288	241

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older, while 2005 estimates are based on all respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

<sup>3</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

<sup>4</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>5</sup> CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

<sup>6</sup> Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

<sup>7</sup> Respondents with unknown health data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.51D Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 18 or Older, and Receipt of Treatment in the Past Year for Depression among Persons Aged 18 or Older with an MDE in Lifetime or Past Year, by Geographic, Socioeconomic, and Health Characteristics: Standard Errors of Percentages, 2004 and 2005**

Geographic/Socioeconomic/Health Characteristic	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.40	0.26	1.46	0.97	0.29	0.20	1.75	1.24
<b>GEOGRAPHIC REGION</b>								
Northeast	0.78	0.62	3.04	2.34	0.68	0.46	3.72	2.66
Midwest	0.64	0.51	2.23	1.76	0.44	0.38	2.85	2.35
South	0.76	0.44	2.79	1.70	0.49	0.33	3.27	2.16
West	0.89	0.59	3.15	2.02	0.70	0.42	3.92	2.72
<b>COUNTY TYPE</b>								
Large Metro	0.57	0.37	2.20	1.40	0.40	0.28	2.51	1.80
Small Metro	0.68	0.48	2.35	1.70	0.51	0.33	3.11	2.12
Nonmetro	0.92	0.58	3.06	2.09	0.71	0.46	3.85	2.42
<b>FAMILY INCOME</b>								
Less Than \$20,000	0.87	0.65	2.89	1.96	0.72	0.55	3.45	2.20
\$20,000 - \$49,999	0.58	0.45	2.29	1.71	0.45	0.33	2.91	2.13
\$50,000 - \$74,999	0.91	0.61	3.22	2.17	0.73	0.44	4.36	2.58
\$75,000 or More	0.87	0.46	3.37	2.07	0.52	0.31	3.44	2.80
<b>GOVERNMENT ASSISTANCE<sup>3</sup></b>								
Yes	1.14	0.79	3.07	1.98	0.97	0.67	3.95	2.08
No	0.42	0.28	1.64	1.08	0.29	0.20	1.93	1.45
<b>HEALTH INSURANCE<sup>4</sup></b>								
Private	0.46	0.30	1.86	1.24	0.33	0.20	2.13	1.63
Medicaid/CHIP <sup>5</sup>	1.49	1.11	3.97	2.41	1.33	1.02	4.59	2.57
Other <sup>6</sup>	0.75	0.63	4.19	2.70	0.49	0.45	3.88	2.73
No Coverage	1.02	0.62	3.13	2.06	0.84	0.48	4.08	2.58
<b>OVERALL HEALTH<sup>7</sup></b>								
Excellent	0.72	0.46	3.22	2.12	0.48	0.27	4.38	3.23
Very Good	0.61	0.40	2.26	1.56	0.43	0.26	2.95	2.17
Good	0.91	0.48	2.80	1.73	0.60	0.36	3.03	2.08
Fair/Poor	1.29	1.06	3.28	2.19	1.14	0.91	3.82	2.28

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older, while 2005 estimates are based on all respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

<sup>3</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

<sup>4</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>5</sup> CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

<sup>6</sup> Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

<sup>7</sup> Respondents with unknown health data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.51P Had at Least One Major Depressive Episode (MDE) in Lifetime or Past Year among Persons Aged 18 or Older, and Receipt of Treatment in the Past Year for Depression among Persons Aged 18 or Older with an MDE in Lifetime or Past Year, by Geographic, Socioeconomic, and Health Characteristics: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Geographic/Socioeconomic/Health Characteristic	Lifetime MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Lifetime MDE <sup>2</sup>		Past Year MDE <sup>1</sup>		Receipt of Treatment in Past Year among Persons with Past Year MDE <sup>2</sup>	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	0.2289		0.3696		0.0414		0.8275	
<b>GEOGRAPHIC REGION</b>								
Northeast	0.3723		0.6565		0.1968		0.5406	
Midwest	0.1967		0.9323		0.8458		0.6713	
South	0.2690		0.6113		0.5207		0.5904	
West	0.2224		0.1188		0.0577		0.2653	
<b>COUNTY TYPE</b>								
Large Metro	0.2143		0.3990		0.0586		0.5685	
Small Metro	0.9122		0.0584		0.4372		0.8171	
Nonmetro	0.6074		0.0464		0.5779		0.2187	
<b>FAMILY INCOME</b>								
Less Than \$20,000	0.3361		0.4661		0.6605		0.4487	
\$20,000 - \$49,999	0.8783		0.5936		0.4918		0.5288	
\$50,000 - \$74,999	0.4491		0.4178		0.2622		0.8072	
\$75,000 or More	0.0344		0.2480		0.0085		0.1946	
<b>GOVERNMENT ASSISTANCE<sup>3</sup></b>								
Yes	0.4629		0.4886		0.7716		0.1611	
No	0.0850		0.1466		0.0095		0.5257	
<b>HEALTH INSURANCE<sup>4</sup></b>								
Private	0.0210		0.0401		0.0005		0.2853	
Medicaid/CHIP <sup>5</sup>	0.0372		0.4287		0.1127		0.6688	
Other <sup>6</sup>	0.0992		0.8137		0.1884		0.5190	
No Coverage	0.1675		0.4897		0.2158		0.1594	
<b>OVERALL HEALTH<sup>7</sup></b>								
Excellent	0.1155		0.1024		0.1830		0.1565	
Very Good	0.1559		0.9620		0.0028		0.3337	
Good	0.1823		0.0912		0.1148		0.5006	
Fair/Poor	0.0239		0.5721		0.0883		0.7073	

\*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older, while 2005 estimates are based on all respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Respondents with unknown data for the respective Lifetime or Past Year MDE measure were excluded.

<sup>2</sup> Treatment is defined as seeing or talking to a medical doctor or other professional or using prescription medication in the past year for depression. Respondents with unknown treatment data were excluded.

<sup>3</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

<sup>4</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>5</sup> CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

<sup>6</sup> Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

<sup>7</sup> Respondents with unknown health data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.