

Table 6.52C Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	Total ¹		PAST YEAR MAJOR DEPRESSIVE EPISODE ²			
			Yes		No	
	2004	2005	2004	2005	2004	2005
PAST YEAR USE						
Illicit Drugs ³	87	88	43	39	91	96
Marijuana and Hashish	77	75	35	32	76	76
Cocaine	27	28	11	12	25	25
Crack	10	10	6	5	8	8
Heroin	10	8	4	4	9	6
Hallucinogens	36	36	17	17	33	31
LSD	17	17	9	9	15	15
PCP	12	14	8	5	9	13
Ecstasy	23	21	12	11	21	18
Inhalants	45	47	21	21	44	44
Nonmedical Use of Psychotherapeutics ⁴	59	61	32	26	54	58
Pain Relievers	56	56	30	24	49	53
OxyContin [®]	18	18	9	8	15	15
Tranquilizers	29	29	14	12	27	26
Stimulants	28	31	14	15	26	27
Methamphetamine	17	19	8	9	15	17
Sedatives	15	16	7	8	13	13
Illicit Drugs Other Than Marijuana ³	71	71	36	32	73	72
PAST MONTH USE						
Daily Cigarette Use ⁵	42	33	17	14	35	31
Heavy Alcohol Use ⁶	36	32	12	12	34	31

*Low precision; no estimate reported.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.52D Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Standard Errors of Percentages, 2004 and 2005

Substance	Total ¹		PAST YEAR MAJOR DEPRESSIVE EPISODE ²			
			Yes		No	
	2004	2005	2004	2005	2004	2005
PAST YEAR USE						
Illicit Drugs ³	0.34	0.35	1.42	1.41	0.35	0.36
Marijuana and Hashish	0.31	0.30	1.29	1.25	0.31	0.30
Cocaine	0.11	0.11	0.47	0.53	0.11	0.11
Crack	0.04	0.04	0.26	0.22	0.04	0.03
Heroin	0.04	0.03	0.17	0.17	0.04	0.03
Hallucinogens	0.14	0.14	0.71	0.73	0.14	0.14
LSD	0.07	0.07	0.41	0.39	0.07	0.06
PCP	0.05	0.06	0.34	0.21	0.04	0.06
Ecstasy	0.09	0.08	0.53	0.48	0.09	0.08
Inhalants	0.18	0.19	0.86	0.91	0.18	0.18
Nonmedical Use of Psychotherapeutics ⁴	0.23	0.24	1.24	1.10	0.23	0.24
Pain Relievers	0.22	0.22	1.16	1.01	0.21	0.22
OxyContin [®]	0.07	0.07	0.40	0.38	0.07	0.07
Tranquilizers	0.12	0.11	0.61	0.53	0.12	0.11
Stimulants	0.11	0.12	0.62	0.68	0.11	0.12
Methamphetamine	0.07	0.07	0.36	0.39	0.07	0.07
Sedatives	0.06	0.06	0.31	0.36	0.06	0.06
Illicit Drugs Other Than Marijuana ³	0.28	0.28	1.34	1.26	0.29	0.28
PAST MONTH USE						
Daily Cigarette Use ⁵	0.17	0.13	0.75	0.61	0.15	0.13
Heavy Alcohol Use ⁶	0.14	0.13	0.52	0.54	0.15	0.13

*Low precision; no estimate reported.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 6.52P Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): P Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	Total ¹		PAST YEAR MAJOR DEPRESSIVE EPISODE ²			
			Yes		No	
	2004	2005	2004	2005	2004	2005
PAST YEAR USE						
Illicit Drugs ³	0.0202		0.1817		0.0296	
Marijuana and Hashish	0.0068		0.3297		0.0099	
Cocaine	0.7965		0.5922		0.9745	
Crack	0.4883		0.7756		0.5112	
Heroin	0.3627		0.9194		0.2026	
Hallucinogens	0.0707		0.3481		0.1086	
LSD	0.7374		0.8729		0.6696	
PCP	0.5166		0.6071		0.2780	
Ecstasy	0.0800		0.5167		0.0691	
Inhalants	0.6354		0.2907		0.2179	
Nonmedical Use of Psychotherapeutics ⁴	0.1208		0.1706		0.2334	
Pain Relievers	0.0895		0.0713		0.2678	
OxyContin [®]	0.7539		0.9599		0.5553	
Tranquilizers	0.3483		0.2578		0.4611	
Stimulants	0.9143		0.4660		0.7028	
Methamphetamine	0.8029		0.9825		0.6935	
Sedatives	0.4779		0.2629		0.1260	
Illicit Drugs Other Than Marijuana ³	0.1172		0.5085		0.1199	
PAST MONTH USE						
Daily Cigarette Use ⁵	0.0173		0.0151		0.1674	
Heavy Alcohol Use ⁶	0.1856		0.5220		0.2137	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 6.53C Substance Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Major Depressive Episode (MDE):
Standard Errors of Numbers in Thousands, 2004 and 2005**

Substance	Total ¹		PAST YEAR MAJOR DEPRESSIVE EPISODE ²			
			Yes		No	
	2004	2005	2004	2005	2004	2005
PAST YEAR USE						
Illicit Drugs ³	734	541	296	195	703	521
Marijuana and Hashish	620	463	233	156	602	439
Cocaine	317	202	121	84	284	181
Crack	172	114	85	57	129	96
Heroin	88	42	39	20	79	38
Hallucinogens	207	140	85	52	186	124
LSD	68	45	22	13	65	40
PCP	30	15	3	6	30	14
Ecstasy	153	111	66	46	134	97
Inhalants	116	79	43	23	105	76
Nonmedical Use of Psychotherapeutics ⁴	465	341	213	137	403	309
Pain Relievers	390	300	192	116	332	269
OxyContin [®]	98	80	43	33	84	71
Tranquilizers	267	206	146	78	215	192
Stimulants	226	115	80	58	207	97
Methamphetamine	156	84	70	44	135	69
Sedatives	96	73	59	34	72	61
Illicit Drugs Other Than Marijuana ³	560	393	241	157	493	362
PAST MONTH USE						
Daily Cigarette Use ⁵	898	628	324	217	906	625
Heavy Alcohol Use ⁶	593	378	171	107	587	378

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Therefore, 2004 estimates in the Total column may differ slightly from 2004 estimates (shown in other tables) for these same measures based on the full sample. Estimates for 2005 are based on all respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.53D Substance Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Major Depressive Episode (MDE): Standard Errors of Percentages, 2004 and 2005

Substance	Total ¹		PAST YEAR MAJOR DEPRESSIVE EPISODE ²			
			Yes		No	
	2004	2005	2004	2005	2004	2005
PAST YEAR USE						
Illicit Drugs ³	0.34	0.25	1.55	1.10	0.33	0.24
Marijuana and Hashish	0.29	0.21	1.30	0.92	0.29	0.21
Cocaine	0.15	0.09	0.70	0.52	0.14	0.09
Crack	0.08	0.05	0.50	0.36	0.07	0.05
Heroin	0.04	0.02	0.23	0.13	0.04	0.02
Hallucinogens	0.10	0.06	0.48	0.33	0.09	0.06
LSD	0.03	0.02	0.13	0.08	0.03	0.02
PCP	0.01	0.01	0.01	0.04	0.02	0.01
Ecstasy	0.07	0.05	0.37	0.29	0.07	0.05
Inhalants	0.05	0.04	0.25	0.15	0.05	0.04
Nonmedical Use of Psychotherapeutics ⁴	0.22	0.16	1.20	0.82	0.20	0.15
Pain Relievers	0.18	0.14	1.08	0.70	0.17	0.13
OxyContin [®]	0.05	0.04	0.25	0.21	0.04	0.04
Tranquilizers	0.12	0.09	0.84	0.49	0.11	0.09
Stimulants	0.10	0.05	0.47	0.37	0.11	0.05
Methamphetamine	0.07	0.04	0.41	0.28	0.07	0.03
Sedatives	0.04	0.03	0.34	0.21	0.04	0.03
Illicit Drugs Other Than Marijuana ³	0.26	0.18	1.30	0.93	0.24	0.17
PAST MONTH USE						
Daily Cigarette Use ⁵	0.42	0.29	1.64	1.17	0.43	0.29
Heavy Alcohol Use ⁶	0.28	0.17	0.97	0.65	0.29	0.18

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Therefore, 2004 estimates in the Total column may differ slightly from 2004 estimates (shown in other tables) for these same measures based on the full sample. Estimates for 2005 are based on all respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

**Table 6.53P Substance Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Major Depressive Episode (MDE):
P Values from Tests of Differences of Percentages, 2005 Versus 2004**

Substance	Total ¹		PAST YEAR MAJOR DEPRESSIVE EPISODE ²			
			Yes		No	
	2004	2005	2004	2005	2004	2005
PAST YEAR USE						
Illicit Drugs ³	0.2069		0.8295		0.2042	
Marijuana and Hashish	0.0854		0.9392		0.0517	
Cocaine	0.5444		0.8565		0.4954	
Crack	0.9905		0.7804		0.6214	
Heroin	0.5763		0.6530		0.7189	
Hallucinogens	0.5819		0.4682		0.7069	
LSD	0.4058		0.2533		0.4789	
PCP	0.3564		0.0397		0.1740	
Ecstasy	0.9076		1.0000		0.9954	
Inhalants	0.3911		0.3089		0.7059	
Nonmedical Use of Psychotherapeutics ⁴	0.1695		0.7517		0.0700	
Pain Relievers	0.1049		0.6872		0.0377	
OxyContin [®]	0.5616		0.4712		0.8003	
Tranquilizers	0.1554		0.5466		0.0366	
Stimulants	0.1786		0.6729		0.1346	
Methamphetamine	0.2884		0.7078		0.1870	
Sedatives	0.8400		0.9141		0.6434	
Illicit Drugs Other Than Marijuana ³	0.4844		0.9668		0.3808	
PAST MONTH USE						
Daily Cigarette Use ⁵	0.7058		0.9185		0.6880	
Heavy Alcohol Use ⁶	0.1348		0.2164		0.2125	

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Therefore, 2004 estimates in the Total column may differ slightly from 2004 estimates (shown in other tables) for these same measures based on the full sample. Estimates for 2005 are based on all respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.