

Table 6.66C Severity of Role Impairment as Measured by the Sheehan Disability Scale (SDS) (I), and Mean Number of Days in the Past Year in which Unable to Carry Out Normal Activities Due to Depression, by Overall Severity of Role Impairment (II), among Persons Aged 12 to 17 with Past Year Major Depressive Episode (MDE): Standard Errors of Numbers in Thousands, 2005

SDS Role Domains (I)	SEVERITY OF ROLE IMPAIRMENT					
	Total ¹	None	Mild	Moderate	Severe	Very Severe
Chores at Home	67	15	29	39	29	22
School or Work	67	13	28	37	35	21
Close Relationships with Family	68	13	22	35	38	26
Social Life	67	9	22	41	37	23
Overall ²	69	2	13	32	44	34
	OVERALL SEVERITY OF ROLE IMPAIRMENT ²					
	Total	None	Mild	Moderate	Severe	Very Severe
Mean Number of Days Unable to Carry Out Normal Activity in the Past Year (II)³	1.90	N/A	4.81	2.77	2.44	6.21

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: The Sheehan Disability Scale (SDS) measures the impact of a disorder on a person's life. The SDS asks youths to give 0 to 10 ratings of the amount of impairment caused by the disorder in each of four role domains: (1) chores at home, (2) school or work, (3) close relationships with family, and (4) social life.

¹ Respondents with unknown severity of SDS role impairment data were excluded.

² Overall role impairment is defined by the highest level of severity of role impairment reported across all four SDS role domains.

³ Respondents with unknown data on the mean number of days that they were unable to carry out normal activities in the past year were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 6.66D Severity of Role Impairment as Measured by the Sheehan Disability Scale (SDS) (I), and Mean Number of Days in the Past Year in which Unable to Carry Out Normal Activities Due to Depression, by Overall Severity of Role Impairment (II), among Persons Aged 12 to 17 with Past Year Major Depressive Episode (MDE): Standard Errors of Percentages, 2005

SDS Role Domains (I)	SEVERITY OF ROLE IMPAIRMENT					
	Total ¹	None	Mild	Moderate	Severe	Very Severe
Chores at Home	0.00	0.70	1.18	1.47	1.18	0.97
School or Work	0.00	0.60	1.18	1.45	1.35	0.93
Close Relationships with Family	0.00	0.59	0.96	1.37	1.42	1.07
Social Life	0.00	0.44	1.00	1.50	1.47	1.00
Overall ²	0.00	0.10	0.60	1.22	1.45	1.33
	OVERALL SEVERITY OF ROLE IMPAIRMENT ²					
	Total	None	Mild	Moderate	Severe	Very Severe
Mean Number of Days Unable to Carry Out Normal Activity in the Past Year (II)³	1.90	N/A	4.81	2.77	2.44	6.21

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: The Sheehan Disability Scale (SDS) measures the impact of a disorder on a person's life. The SDS asks youths to give 0 to 10 ratings of the amount of impairment caused by the disorder in each of four role domains: (1) chores at home, (2) school or work, (3) close relationships with family, and (4) social life.

¹ Respondents with unknown severity of SDS role impairment data were excluded.

² Overall role impairment is defined by the highest level of severity of role impairment reported across all four SDS role domains.

³ Respondents with unknown data on the mean number of days that they were unable to carry out normal activities in the past year were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 6.67C Severity of Role Impairment as Measured by the Sheehan Disability Scale (SDS) (I), and Mean Number of Days in the Past Year in which Unable to Carry Out Normal Activities Due to Depression, by Overall Severity of Role Impairment (II), among Persons Aged 18 or Older with Past Year Major Depressive Episode (MDE): Standard Errors of Numbers in Thousands, 2005

SDS Role Domains (I)	SEVERITY OF ROLE IMPAIRMENT					
	Total ¹	None	Mild	Moderate	Severe	Very Severe
Home Management	432	65	139	270	247	146
Work	401	87	198	217	210	129
Close Relationships with Others	425	75	163	255	237	113
Social Life	426	80	135	246	271	141
Overall ²	436	16	100	224	308	174
	OVERALL SEVERITY OF ROLE IMPAIRMENT ²					
	Total	None	Mild	Moderate	Severe	Very Severe
Mean Number of Days Unable to Carry Out Normal Activity in the Past Year (II)³	3.28	N/A	*	2.76	4.90	8.78

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: The Sheehan Disability Scale (SDS) measures the impact of a disorder on a person's life. The SDS asks adults to give 0 to 10 ratings of the amount of impairment caused by the disorder in each of four role domains: (1) home management, (2) work, (3) close relationships with others, and (4) social life.

¹ Respondents with unknown severity of SDS role impairment data were excluded.

² Overall role impairment is defined by the highest level of severity of role impairment reported across all four SDS role domains.

³ Respondents with unknown data on the mean number of days that they were unable to carry out normal activities in the past year were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 6.67D Severity of Role Impairment as Measured by the Sheehan Disability Scale (SDS) (I), and Mean Number of Days in the Past Year in which Unable to Carry Out Normal Activities Due to Depression, by Overall Severity of Role Impairment (II), among Persons Aged 18 or Older with Past Year Major Depressive Episode (MDE): Standard Errors of Percentages, 2005

SDS Role Domains (I)	SEVERITY OF ROLE IMPAIRMENT					
	Total ¹	None	Mild	Moderate	Severe	Very Severe
Home Management	0.00	0.42	0.85	1.35	1.32	0.90
Work	0.00	0.61	1.19	1.30	1.22	0.86
Close Relationships with Others	0.00	0.48	0.96	1.33	1.28	0.71
Social Life	0.00	0.51	0.84	1.35	1.39	0.87
Overall ²	0.00	0.10	0.61	1.23	1.40	1.02
	OVERALL SEVERITY OF ROLE IMPAIRMENT ²					
	Total	None	Mild	Moderate	Severe	Very Severe
Mean Number of Days Unable to Carry Out Normal Activity in the Past Year (II)³	3.28	N/A	*	2.76	4.90	8.78

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: The Sheehan Disability Scale (SDS) measures the impact of a disorder on a person's life. The SDS asks adults to give 0 to 10 ratings of the amount of impairment caused by the disorder in each of four role domains: (1) home management, (2) work, (3) close relationships with others, and (4) social life.

¹ Respondents with unknown severity of SDS role impairment data were excluded.

² Overall role impairment is defined by the highest level of severity of role impairment reported across all four SDS role domains.

³ Respondents with unknown data on the mean number of days that they were unable to carry out normal activities in the past year were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.