

Table 6.7C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Substance Dependence or Abuse: Standard Errors of Numbers in Thousands, 2004 and 2005

Dependence/Abuse	Total		DEPENDENCE ¹				DEPENDENCE OR ABUSE ¹			
			Yes		No		Yes		No	
	2004	2005	2004	2005	2004	2005	2004	2005	2004	2005
ILLICIT DRUGS²	750	505	183	101	744	518	212	116	726	512
Marijuana	750	505	141	63	755	525	171	76	737	522
ILLICIT DRUGS OTHER THAN MARIJUANA²	750	505	129	88	761	522	148	98	757	517
ALCOHOL	750	505	231	138	742	508	284	178	719	492
ILLICIT DRUGS OR ALCOHOL²	750	505	278	159	711	503	333	197	678	482
BOTH ILLICIT DRUGS AND ALCOHOL²	750	505	87	64	772	522	123	76	764	521

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.7D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Substance Dependence or Abuse: Standard Errors of Percentages, 2004 and 2005

Dependence/Abuse	Total		DEPENDENCE ¹				DEPENDENCE OR ABUSE ¹			
			Yes		No		Yes		No	
	2004	2005	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS²	0.35	0.23	3.27	1.96	0.34	0.23	2.73	1.64	0.34	0.23
Marijuana	0.35	0.23	4.63	2.50	0.34	0.23	3.60	2.00	0.34	0.23
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.35	0.23	4.22	2.87	0.34	0.23	3.42	2.41	0.34	0.23
ALCOHOL	0.35	0.23	2.57	1.56	0.35	0.23	1.50	0.92	0.36	0.23
ILLCIT DRUGS OR ALCOHOL²	0.35	0.23	2.07	1.27	0.34	0.23	1.42	0.86	0.34	0.23
BOTH ILLICIT DRUGS AND ALCOHOL²	0.35	0.23	5.59	3.66	0.35	0.23	3.10	2.31	0.35	0.23

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.7P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Substance Dependence or Abuse: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Dependence/Abuse	Total		DEPENDENCE ¹				DEPENDENCE OR ABUSE ¹			
			Yes		No		Yes		No	
	2004	2005	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS²	0.0348		0.5188		0.0578		0.2989		0.0935	
Marijuana	0.0348		0.4544		0.0530		0.3335		0.0744	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.0348		0.9404		0.0396		0.4543		0.0512	
ALCOHOL	0.0348		0.4735		0.0617		0.2502		0.0753	
ILLCIT DRUGS OR ALCOHOL²	0.0348		0.2291		0.1097		0.1187		0.1400	
BOTH ILLICIT DRUGS AND ALCOHOL²	0.0348		0.7854		0.0317		0.4108		0.0503	

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.8C Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Numbers in Thousands, 2004 and 2005

Dependence/Abuse	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
DEPENDENCE OR ABUSE²						
Illicit Drugs ³	354	199	212	116	297	161
Marijuana	285	137	171	76	241	116
Illicit Drugs Other Than Marijuana ³	233	156	148	98	179	120
Alcohol	667	394	284	178	621	369
Illicit Drugs or Alcohol ³	694	421	333	197	640	388
Both Illicit Drugs and Alcohol ³	214	135	123	76	163	112
DEPENDENCE²						
Illicit Drugs ³	267	169	183	101	214	132
Marijuana	191	116	141	63	158	96
Illicit Drugs Other Than Marijuana ³	204	136	129	88	158	100
Alcohol	449	260	231	138	394	222
Illicit Drugs or Alcohol ³	486	298	278	159	413	250
Both Illicit Drugs and Alcohol ³	136	88	87	64	101	59

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.8D Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Percentages, 2004 and 2005

Dependence/Abuse	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
DEPENDENCE OR ABUSE²						
Illicit Drugs ³	0.16	0.09	0.75	0.46	0.16	0.08
Marijuana	0.13	0.06	0.61	0.31	0.13	0.06
Illicit Drugs Other Than Marijuana ³	0.11	0.07	0.55	0.39	0.09	0.06
Alcohol	0.31	0.18	1.00	0.68	0.32	0.18
Illicit Drugs or Alcohol ³	0.32	0.19	1.11	0.74	0.33	0.19
Both Illicit Drugs and Alcohol ³	0.10	0.06	0.46	0.31	0.09	0.06
DEPENDENCE²						
Illicit Drugs ³	0.12	0.08	0.66	0.41	0.11	0.07
Marijuana	0.09	0.05	0.52	0.26	0.08	0.05
Illicit Drugs Other Than Marijuana ³	0.09	0.06	0.48	0.35	0.08	0.05
Alcohol	0.21	0.12	0.84	0.54	0.21	0.11
Illicit Drugs or Alcohol ³	0.23	0.14	0.97	0.62	0.21	0.13
Both Illicit Drugs and Alcohol ³	0.06	0.04	0.33	0.26	0.05	0.03

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.8P Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Dependence/Abuse	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
DEPENDENCE OR ABUSE²						
Illicit Drugs ³	0.0212		0.0806		0.2307	
Marijuana	0.0080		0.0455		0.1179	
Illicit Drugs Other Than Marijuana ³	0.3528		0.5221		0.7842	
Alcohol	0.6264		0.6976		0.9272	
Illicit Drugs or Alcohol ³	0.2051		0.2460		0.6152	
Both Illicit Drugs and Alcohol ³	0.2598		0.4002		0.6020	
DEPENDENCE²						
Illicit Drugs ³	0.1486		0.4059		0.4790	
Marijuana	0.0945		0.2552		0.4070	
Illicit Drugs Other Than Marijuana ³	0.4110		0.8894		0.5798	
Alcohol	0.2832		0.5305		0.5655	
Illicit Drugs or Alcohol ³	0.0798		0.1912		0.3971	
Both Illicit Drugs and Alcohol ³	0.9316		0.5781		0.7729	

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.9C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2004 and 2005

Gender/Age	Total		ILLCIT DRUG DEPENDENCE OR ABUSE ^{1,2}			
			Yes		No	
	2004	2005	2004	2005	2004	2005
TOTAL	750	505	212	116	726	512
18-25	159	105	80	50	168	125
26-49	516	334	168	93	522	334
50 or Older	481	332	*	*	472	340
MALE	454	301	175	78	447	302
18-25	113	70	64	38	107	73
26-49	319	207	127	65	302	203
50 or Older	310	193	*	*	306	198
FEMALE	606	390	118	87	594	402
18-25	122	76	46	32	133	90
26-49	410	246	*	70	433	254
50 or Older	372	275	*	*	376	282

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.9D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2004 and 2005

Gender/Age	Total		ILLCIT DRUG DEPENDENCE OR ABUSE ^{1,2}			
			Yes		No	
	2004	2005	2004	2005	2004	2005
TOTAL	0.35	0.23	2.73	1.64	0.34	0.23
18-25	0.50	0.32	2.20	1.39	0.49	0.34
26-49	0.52	0.33	4.02	3.06	0.51	0.33
50 or Older	0.58	0.39	*	*	0.57	0.39
MALE	0.44	0.29	3.29	1.88	0.43	0.29
18-25	0.70	0.43	2.73	1.79	0.67	0.43
26-49	0.65	0.42	4.80	3.60	0.63	0.42
50 or Older	0.81	0.49	*	*	0.78	0.49
FEMALE	0.54	0.35	4.50	2.84	0.54	0.34
18-25	0.76	0.47	3.48	2.28	0.76	0.48
26-49	0.81	0.49	*	5.03	0.80	0.48
50 or Older	0.83	0.60	*	*	0.83	0.59

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.9P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug Dependence or Abuse, Gender, and Age Group: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Gender/Age	Total		ILLCIT DRUG DEPENDENCE OR ABUSE ^{1,2}			
			Yes		No	
	2004	2005	2004	2005	2004	2005
TOTAL	0.0348		0.2989		0.0935	
18-25	0.0073		0.0735		0.0369	
26-49	0.0139		0.9581		0.0273	
50 or Older	0.7360		*		0.6253	
MALE	0.0504		0.0300		0.2838	
18-25	0.0269		0.0600		0.2093	
26-49	0.2866		0.4014		0.5404	
50 or Older	0.3518		*		0.6199	
FEMALE	0.2467		0.4564		0.2237	
18-25	0.1306		0.2444		0.1228	
26-49	0.0210		*		0.0215	
50 or Older	0.2424		*		0.3151	

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.10C Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2004 and 2005

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
TOTAL	354	199	212	116	297	161
18-25	123	83	80	50	100	69
26-49	268	146	168	93	210	121
50 or Older	168	92	106	44	131	81
MALE	256	158	175	78	208	139
18-25	100	65	64	38	88	59
26-49	204	123	127	65	155	110
50 or Older	153	68	*	21	112	66
FEMALE	200	121	118	87	162	82
18-25	60	47	46	32	41	34
26-49	174	89	110	70	138	57
50 or Older	68	62	*	39	68	48

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.5 of Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.10D Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: Standard Errors of Percentages, 2004 and 2005

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
TOTAL	0.16	0.09	0.75	0.46	0.16	0.08
18-25	0.38	0.26	1.09	0.76	0.37	0.25
26-49	0.27	0.15	1.14	0.71	0.24	0.14
50 or Older	0.20	0.11	1.80	0.71	0.17	0.10
MALE	0.25	0.15	1.67	0.88	0.22	0.14
18-25	0.62	0.40	2.01	1.42	0.62	0.39
26-49	0.41	0.25	2.41	1.39	0.35	0.24
50 or Older	0.40	0.17	*	1.05	0.31	0.18
FEMALE	0.18	0.11	0.71	0.54	0.17	0.09
18-25	0.38	0.29	1.10	0.82	0.33	0.27
26-49	0.34	0.18	1.17	0.84	0.33	0.13
50 or Older	0.15	0.13	*	0.93	0.16	0.11

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.5 of Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.10P Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
TOTAL	0.0212		0.0806		0.2307	
18-25	0.6232		0.4159		0.6599	
26-49	0.0143		0.3130		0.0513	
50 or Older	0.6610		0.4044		0.9665	
MALE	0.0089		0.0138		0.3344	
18-25	0.1058		0.0881		0.6535	
26-49	0.0775		0.1811		0.2664	
50 or Older	0.3623		*		0.8908	
FEMALE	0.5164		0.6560		0.3590	
18-25	0.1162		0.3774		0.0733	
26-49	0.0935		0.9201		0.0907	
50 or Older	0.4503		*		0.8736	

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.5 of Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.11C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2004 and 2005

Gender/Age	Total		ALCOHOL DEPENDENCE OR ABUSE ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
TOTAL	750	505	284	178	719	492
18-25	159	105	96	67	156	112
26-49	516	334	227	132	496	323
50 or Older	481	332	*	93	473	331
MALE	454	301	229	130	421	279
18-25	113	70	74	46	103	67
26-49	319	207	173	94	279	186
50 or Older	310	193	*	75	297	185
FEMALE	606	390	158	118	568	395
18-25	122	76	61	46	124	81
26-49	410	246	140	92	409	248
50 or Older	372	275	*	*	374	279

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.11D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2004 and 2005

Gender/Age	Total		ALCOHOL DEPENDENCE OR ABUSE ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
TOTAL	0.35	0.23	1.50	0.92	0.36	0.23
18-25	0.50	0.32	1.38	0.95	0.50	0.34
26-49	0.52	0.33	2.20	1.33	0.52	0.34
50 or Older	0.58	0.39	*	3.15	0.59	0.39
MALE	0.44	0.29	1.78	1.04	0.44	0.29
18-25	0.70	0.43	1.74	1.13	0.72	0.45
26-49	0.65	0.42	2.44	1.46	0.63	0.42
50 or Older	0.81	0.49	*	3.50	0.80	0.49
FEMALE	0.54	0.35	2.62	1.70	0.53	0.35
18-25	0.76	0.47	2.48	1.61	0.76	0.48
26-49	0.81	0.49	4.05	2.57	0.79	0.48
50 or Older	0.83	0.60	*	*	0.84	0.60

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.11P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Alcohol Dependence or Abuse, Gender, and Age Group: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Gender/Age	Total		ALCOHOL DEPENDENCE OR ABUSE ¹	
			Yes	No
	2004	2005	2004	2005
TOTAL	0.0348		0.2502	0.0753
18-25	0.0073		0.0301	0.0403
26-49	0.0139		0.2840	0.0290
50 or Older	0.7360		*	0.8194
MALE	0.0504		0.1196	0.1919
18-25	0.0269		0.0288	0.1920
26-49	0.2866		0.2554	0.6045
50 or Older	0.3518		*	0.3790
FEMALE	0.2467		0.9312	0.2128
18-25	0.1306		0.4045	0.1357
26-49	0.0210		0.5141	0.0193
50 or Older	0.2424		*	0.3414

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

¹ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.12C Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2004 and 2005

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
TOTAL	667	394	284	178	621	369
18-25	156	119	96	67	137	122
26-49	443	278	227	132	399	263
50 or Older	396	220	135	93	380	208
MALE	525	315	229	130	523	309
18-25	123	88	74	46	111	99
26-49	375	229	173	94	352	226
50 or Older	329	183	*	75	321	179
FEMALE	329	210	158	118	291	179
18-25	91	69	61	46	79	58
26-49	228	154	140	92	187	128
50 or Older	193	116	39	55	191	108

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.5 of Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.12D Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: Standard Errors of Percentages, 2004 and 2005

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
TOTAL	0.31	0.18	1.00	0.68	0.32	0.18
18-25	0.49	0.37	1.21	0.92	0.49	0.38
26-49	0.44	0.28	1.46	0.98	0.46	0.28
50 or Older	0.48	0.26	2.30	1.48	0.48	0.26
MALE	0.51	0.30	2.05	1.33	0.52	0.31
18-25	0.76	0.54	2.26	1.58	0.75	0.57
26-49	0.76	0.46	3.03	1.85	0.76	0.47
50 or Older	0.86	0.47	*	3.57	0.86	0.47
FEMALE	0.29	0.19	0.90	0.73	0.30	0.18
18-25	0.57	0.43	1.35	1.06	0.62	0.43
26-49	0.45	0.30	1.39	1.09	0.44	0.29
50 or Older	0.43	0.25	1.13	1.31	0.45	0.25

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.5 of Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.

Table 6.12P Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2004	2005	2004	2005	2004	2005
TOTAL	0.6264		0.6976		0.9272	
18-25	0.3855		0.8869		0.1312	
26-49	0.5214		0.7264		0.7868	
50 or Older	0.7686		0.7944		0.6640	
MALE	0.4223		0.3441		0.8207	
18-25	0.6791		0.4976		0.2488	
26-49	0.3378		0.2777		0.6224	
50 or Older	0.8248		*		0.8766	
FEMALE	0.7859		0.4783		0.8785	
18-25	0.3513		0.5167		0.3202	
26-49	0.7661		0.5694		0.6758	
50 or Older	0.8131		0.2134		0.4969	

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these 2004 and 2005 estimates are not comparable with 2004 and earlier estimates published in prior NSDUH reports. See Section B.4.5 of Appendix B of the *Results from the 2005 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample) and 2005.